

DOCTOR OF PHILOSOPHY

Investigating the role of Annexin A1 in obesity and associated inflammation

Sajid, Sehar

Award date:
2020

Awarding institution:
Coventry University

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of this thesis for personal non-commercial research or study
- This thesis cannot be reproduced or quoted extensively from without first obtaining permission from the copyright holder(s)
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Investigating the role of Annexin A1 in obesity and associated inflammation

By

Sehar Sajid

**A thesis submitted to the Centre of Sport, Exercise and
Life Sciences, Faculty of Health and Life Sciences,
Coventry University**

Vol II

July 2019

Investigating the role of Annexin A1 in obesity and associated inflammation

By

Sehar Sajid

**A thesis submitted in partial fulfilment of the
University's requirements for the Degree of Philosophy**

July 2019



Chapter 5: Annexin A1 regulates adipogenesis via differential regulation of MAPK intracellular pathways

The results observed in Chapter 4 show AC2-26 significantly regulates the expression of several genes that are involved in adipocyte differentiation. Interestingly, previous *in vivo* studies of Annexin A1 gene knockout in mice report conflicting findings. Therefore, the objective of this study was to investigate the role of Annexin A1 in adipocyte differentiation using receptor agonist and antagonists to further elucidate the changes in gene expression and protein secretion in adipocytes in obesity.

5.1 Abstract

Introduction: Adipose tissue expansion is mediated via two mechanisms; hypertrophy and hyperplasia. Dysregulated expansion observed in obesity is associated with local and systematic inflammation. Adipocyte differentiation is regulated by a range of intracellular and extracellular stimuli that induce the expression of genes involved in maturation of pre-adipocyte into a mature adipocyte phenotype. The results from chapter 4, indicate Annexin A1 (ANXA1) regulates the expression of these genes in mature obesogenic adipocytes, however, the effect of ANXA1 and/or its receptors during adipocyte differentiation is unknown.

Aim: The aim of this study was to investigate the role of ANXA1 and its receptors in adipocyte differentiation.

Methodology: Simpson Golabi Behmel Syndrome (SGBS) preadipocytes were differentiated with receptor agonist (AC2-26) and antagonists (WRW4 and BOC-2) over 14 days to reach a mature adipocyte phenotype. Morphological changes (lipid accumulation, number of lipid droplets and size of lipid droplets), mRNA expressions of adipogenic nuclear factors, lipogenic and lipolytic enzymes and adipokines, glucose uptake and activation of proteins involved in intracellular signalling pathways were investigated. The results were statistically analysed using GraphPad prism 5.0. Statistical significance at 95% was determined using One-tailed unpaired students T-test

Results: Mature cells differentiated with WRW4, AC2-26 or WRW4 and AC2-26 accumulated significantly less lipids and altered morphology of lipid droplets compared to vehicle control. The mRNA expression of adipogenic nuclear factors (*PPARG* and *SREBP*) lipolytic enzymes (*ACC*, *FASN* and *GPAM*), lipolytic enzymes (*ATGL*, *HSL* and *MGL*), adipokines (*LEP*, *RETN*, *NAMPT*, *ADIPOQ* and *ANXA1*) and glucose uptake

transporters (*SLC2A1* and *SLC2A4*) were significantly downregulated and glucose uptake was significantly reduced in mature cells differentiated with WRW4, AC2-26 or WRW4 and AC2-26, compared to vehicle control. Furthermore, intracellular activation of p38 mitogen activated protein (p38 MAPK) were significantly decreased in mature cells differentiated with WRW4, AC2-26 or WRW4 and AC2-26, compared to vehicle control. However, mature cells differentiated with low concentrations of BOC-2 accumulated less lipids and mature cells differentiated with high concentration of BOC-2 accumulated more lipids, compared to vehicle control.

Conclusion: This effect was only apparent during later stages of adipocyte differentiation. ANXA1 may regulate the rate of adipogenesis through differential regulation of MAPK intracellular pathways. Activation of Formyl Peptide Receptor 1 (FPR1) by AC2-26/ANXA1 decreases p38 MAPK activation thus, decreasing lipid accumulation in mature adipocytes through downregulation of genes associated with adipogenesis.

5.2 Introduction

Adipose tissue is a complex, multidepot organ, producing and secreting bioactive molecules which influence a range of physiological functions including; appetite, energy expenditure, insulin sensitivity, vascular function and immunity (Osborn, Olefsky 2012). The importance of adipose tissue for health has been demonstrated through mouse models of lipoatrophy (Gavrilova, Marcus-Samuels et al. 2000, Shimomura, Hammer et al. 1999). These mice are genetically altered to express a dominant-negative protein, A-ZIP/F. It prevents the DNA binding of B-ZIP transcription factors of both CCAAT-enhancer-binding protein (C/EBP) and JUN families, inhibiting adipogenesis, therefore, resulting in very little or no white fat tissue (Moitra, Mason et al. 1998). Transplantation of adipose tissue from healthy mice, drastically reverses hyperglycaemia, improves insulin sensitivity, decreases circulating triacylglyceride, decreases hepatic gluconeogenesis and decreases ectopic fat accumulation in these mice (Gavrilova et al. 2000). However, obese individuals carrying excess amounts of adipose tissue display similar characteristics to lipoatrophic mice and humans and are at a higher risk of developing several serious chronic diseases, such as type 2 diabetes mellitus (T2DM), cardiovascular disease, hypertension, hypercholesterolemia and hypertriglyceridemia (Greenberg, Obin 2006). This suggests, an optimal amount of adipose tissue is required for normal health, too little or too much will result in ill health.

Adipose tissue expands and remodels itself by two mechanisms; hypertrophy; increase in cell size and by hyperplasia; increase in cell number to accommodate for excess nutrients (Ali, Hochfeld et al. 2013). Dysregulated expansion of the adipose tissue through these mechanisms is associated with intracellular abnormalities such as, endoplasmic reticulum and mitochondrial stress, leading to adipocyte insulin resistance, production of

adipokines, free fatty acids and inflammatory mediators; reflecting clinical manifestations of obesity (Figure 5.1) (de Ferranti, Mozaffarian 2008). In an adult, hypertrophy of adipocytes is the most common form of storing excess lipids, whereas, hyperplasia is most common in the prepubertal age (Gustafson, Gogg et al. 2009). Although, recruitment of new adipocytes in adults is less common, a recent study demonstrated a continuous turnover of ~10%/year of new adipocytes in adults (Spalding, Arner et al. 2008). Increase in fat mass in this manner, results in a metabolically healthy, insulin sensitive obese phenotype, thus hyperplastic growth results in functionally healthy smaller adipocytes. It is estimated that 20-30% of the obese population are characterised as ‘metabolically healthy but obese’ (Karelis, St-Pierre et al. 2004).

Adipocyte size positively correlates with whole body insulin resistance independent of body mass index, reflective of hypertrophic obesity carrying a greater risk of metabolic complications compared to hyperplastic obesity. The reduced rate of adipocyte formation in hypertrophic obesity is due to i) inadequate preadipocyte number, as observed in obese versus (vs) lean subjects or ii) attenuated differentiation capacity of preadipocytes, thus inducing hypertrophic growth resulting in enlarged swollen adipocytes predisposed to inflammation and insulin resistance (Hirsch, Fried et al. 1989, Danforth 2000, Heilbronn, Smith et al. 2004). Failure of subcutaneous preadipocytes to differentiate has been linked to central obesity (Permana, Nair et al. 2004).

However, adipocyte hypertrophy and hyperplasia are not separate phenomena, it has been suggested they co-exist and equally contribute to the metabolic dysfunctions in obesity. Hyperplasia is associated with greater severity and less reversibility of metabolic consequences (de Ferranti, Mozaffarian 2008) . Enlargement of the adipocytes alters the phenotype of these cells, resulting in major inflammatory responses in the tissue with

increased expression and secretion of pro-inflammatory in parallel to decreased expression and secretion of anti-inflammatory cytokines and adipokines (Divella, De Luca et al. 2016) . Conclusions from *in vitro* studies suggest factors such as tumour necrosis factor α (TNF α) and insulin growth factor-1 secreted by hypertrophied adipocytes stimulate hyperplasia of themselves and adjacent cells leading to altered adipocyte size (Avram, Avram et al. 2007).

The expansion of the adipose tissue is a tightly regulated process and requires communication between the cells and communication between the cells and their surrounding environment (Ali et al. 2013). Mitogens, glucocorticoids, hormones, lipids, synthetic drugs such as, thiazolidinediones and those that lead to an increase in cyclic adenosine monophosphate induce adipogenesis (Ali et al. 2013). Glucocorticoids exert complex actions on adipocytes as chronic treatment promotes adipogenesis through induction of key transcription factors such as, peroxisome proliferator activated receptor γ (PPAR γ) thus inducing lipogenesis, whereas, acute treatment promotes lipolysis (John, Marino et al. 2016).




Figure 5.1: The consequences of energy imbalance. Increased energy consumption and decreased energy expenditure affect energy homeostasis. Energy imbalance disrupts the normal function of β cells, adipocytes and hepatocytes. Chronic expansion of adipose tissue results in cell dysfunction and increased release of adipokines, free fatty acids and inflammatory mediators which affect skeletal muscle, cardiac muscle and endothelium, increasing the risk of developing Type 2 diabetes mellitus, Cardiovascular Disease and Atherosclerosis.

ER; Endoplasmic reticulum

Adapted from (de Ferranti, Mozaffarian 2008)

Previous *in vivo* studies of Annexin A1 (ANXA1) gene knockout in mice report conflicting findings. Decrease in number of fat cells thus, decreased weight was observed in *ANXA1*^{-/-} mice compared to WT by (Warne, John et al. 2006), whereas, (Akasheh et al. 2013) in their study report significant increase in adiposity, blunted lipolysis and elevated hepatic lipogenesis. Furthermore, results from Chapter 4 suggest ANXA1 is involved in regulating lipid accumulation, secretion of adipokines and insulin activity and, therefore differentiation of adipocytes.

5.3 Aim, objectives and hypothesis

The aim of this study was to investigate the role of ANXA1 in the differentiation of preadipocytes into mature adipocytes.

- To assess lipid accumulation in Simpson Golabi Behmel Syndrome (SGBS) preadipocytes differentiated with and without formyl peptide receptor 2 (FPR2/ALX) antagonist and agonist
- To assess the morphology of the lipid droplets in mature SGBS adipocytes differentiated with and without FPR2/ALX antagonist and agonist
- To identify the intracellular signalling pathways activated in mature SGBS adipocytes differentiated with and without FPR2/ALX antagonist and agonist
- To assess *de-novo* lipogenesis, lipolysis and glucose uptake in mature SGBS adipocytes differentiated with and without FPR2/ALX antagonist and agonist
- To assess the gene expression of adipokines in mature SGBS adipocytes differentiated with and without FPR2/ALX antagonist and agonist

It is hypothesised that ANXA1 regulates lipid homeostasis through modulating PPAR γ and is positively involved in regulating the differentiation process.

5.4 Materials and Methods

5.4.1 Differentiation of SGBS preadipocytes with FPR2/ALX receptor agonist and antagonists

WRWWWW (WRW4); a trp-rich hexapeptide, is a potent and specific antagonist for FPR2/ALX as it binds to and interferes with ligand-receptor interactions (IC_{50} : 230nM for agonist; WKYMVm). The antagonistic effects include, blocking ligand activated calcium increase, extracellular signal-regulated kinase (ERK) activation, chemotactic migration and internalisation in human macrophages. Similarly, a non-specific antagonist BOC-FLFLFL (BOC-2), antagonises all members of the FPR receptor family and blocks the ligand-receptor interaction inhibiting ligand induced activation of downstream signalling pathways.

5.4.1.1 Methodology

SGBS cells were differentiated using the methodology described in section 3.1.1.1.1 with the addition of 10 μ M WRW4 (Millipore#344220) and/or 10 μ M AC2-26 or dimethylsulfoxide (DMSO) (vehicle control). 10 μ M of WRW4 is enough to antagonise the receptor without triggering toxic effects. (Bae, Park et al. 2009). Similarly, SGBS cells were differentiated with the addition of 1 μ M, 10 μ M or 100 μ M BOC-2 (2BScientific#RP12950) or vehicle control. WRW4 and/or AC2-26 or BOC-2 and vehicle control were added to the cells at day 0, day 4, day 8 and day 12, coinciding with media changes. Cells were fixed and stained for analysis using the methodology described in section 3.1.2.1.1 at day 0, day 4, day 8 and day 14, or the cells were homogenised (section 3.2.1.1 (RNA) or section 5.4.5.1.1 (protein)) and the cell media were stored at -80°C, for analysis.

5.4.2 Staining and analysis of lipid droplets in SGBS cells

Staining is a technique used in microscopy to enhance contrast in the microscopic image. Stains and dyes are used to highlight structures in biological tissues to define and examine bulk tissues, cell populations or organelles within individual cells. Lipid staining dyes include, Nile Red A; a photostable, lipophilic stain with a bright red fluorescence, Sudan III A lysochrome and Oil Red O (ORO); fat soluble dyes predominantly used for staining triglycerides in animal tissues with a maximal absorption at 518nm, stains neutral lipids and cholesteryl esters but not biological membranes (Ramirez-Zacarias, Castro-Munozledo et al. 1992).

5.4.2.1 Methodology

Prior to staining, the cells were fixed using paraformaldehyde. It causes covalent cross-links between molecules, effectively gluing them together into an insoluble meshwork. The cells were washed phosphate buffered saline (PBS) twice and incubated with fixative solution (4% paraformaldehyde and 1.5% Methanol in PBS) for 30 minutes at room temperature. The cells were washed with PBS twice and refrigerated or stained for lipid droplets.

A stock solution of ORO (Sigma#O0625) was prepared; 1mg/mL in 60% Triethyl Phosphate. The stock solution was mixed to allow the crystals to dissolve for an hour on a roller shaker at room temperature. A working solution of the dye of 60:40 in distilled water was prepared and filtered through Whatman filter paper. The cells were incubated with the dye in the dark for 2-3 hours at room temperature, washed twice with PBS to remove unbound dye and quantified for adipogenesis or stored at 4°C.

Adipogenic differentiation of SGBS preadipocytes for lipid quantification were carried out in 6 well plates. Approximately 8-10 images were taken per well using a light microscope at x10. The amount of ORO stain was proportional to the amount of lipids and size of adipocytes. The images were analysed using Image J following the methodology described by (Mehlem, Hagberg et al. 2013). Briefly, the images were changed from a Red-Green-Blue (RGB) into 8-bit binary images. The threshold was adjusted manually based on saturation and intensity to minimise any background staining and identify areas of interest using a control image. The threshold used in this chapter was set as, Lower limit: 33 and Upper limit: 103. The % area red was analysed per picture, as it corresponded to lipid accumulation (Chapter 5, Figure 5.2). Furthermore, the average size of lipid droplets, the average number of lipid droplets, and the average number of lipid droplets of varying sizes were analysed using the 'analyse particles' function. The sizes of the particles were manually entered as 0-10, 0-100, 100-1000 or 1000-infinity.

5.4.3 Cell viability assay

Cellular viability and toxicity can be measured by assessing various indicators including; plasma membrane integrity, DNA synthesis, DNA content, enzyme activity, presence of adenosine triphosphate (ATP) and cellular reducing conditions. Cell viability reagents such as, AlamarBlueTM use the reducing power of living cells to quantitatively measure proliferation of various cells types to establish relative cytotoxicity.

The main active ingredient in AlamarBlueTM; Resazurin is a non-toxic, cell permeable compound that is blue in colour and non-fluorescent. Live cells reduce compounds in their cytosol. Upon entering the cell, resazurin is reduced to resorufin, a compound red in

colour and highly fluorescent. Viable cells continue to convert resazurin, increasing the overall fluorescence and colour of the media surrounding the cells.

5.4.3.1 Methodology

The cellular viability was assessed using AlamarBlue™ (Invitrogen#DAL1025) by following the protocol provided by the manufacture. Briefly, SGBS preadipocytes were differentiated in 96 well plates with agonist, and/or antagonists, as described in section 5.4.1.1. On day 14, 10% AlamarBlue™ was added per well (i.e. 10µL in 100µL sample) to treated cells, positive and negative controls and incubated for 4 hours at 37°C (95% O₂/5% CO₂). The absorbance of the reagent was measured using a spectrophotometer at 562nm, using 600nm as a reference wavelength. The standard curve generated was used to calculate % cellular viability.

5.4.4 Gene expression analysis

Cells homogenised in section were used to isolate ribonucleic acid (RNA) using methodology described in and synthesise complementary deoxyribonucleic acid (cDNA) using the methodology described in to analyse the gene expression of an array of different genes listed below:

10X *PPARG* QuantiTect Primer Assay (Qaigen#QT00029841), 10X *SREBP* QuantiTect Primer Assay (Qaigen#QT00036897), 10X *ACC* QuantiTect Primer Assay (Qaigen#QT00033761), 10X *FAS* QuantiTect Primer Assay (Qaigen#QT00014588), 10X *GPAM* QuantiTect Primer Assay (Qaigen#QT00009324), 10X *ATGL* QuantiTect Primer Assay (Qaigen#QT00019754), 10X *HSL* QuantiTect Primer Assay (Qaigen#QT00016093), 10X *MGL* QuantiTect Primer Assay (Qaigen#QT00039837), 10X *SLC2A1* QuantiTect Primer Assay (Qaigen#QT00068957), 10X *SLC2A4* QuantiTect

Primer Assay (Qaigen#QT00097902), 10X *LEP* QuantiTect Primer Assay (Qaigen#QT00030261), 10X *RET* QuantiTect Primer Assay (Qaigen#QT00210483), 10X *NAMPT* QuantiTect Primer Assay (Qaigen#QT00087920), 10X *ADIPOQ* QuantiTect Primer Assay (Qaigen#QT00014091) and 10X *ANXA1* QuantiTect Primer Assay (Qaigen#QT00078197).

5.4.4.1 Methodology

The methodology described in section was used to assess the expression of genes and the data obtained was normalised against the housekeeping gene (18s RNA) and relative gene expression was calculated using the methodology described in section 3.2.5.

5.4.5 Intracellular protein expression analysis

5.4.5.1 Protein extraction

To quantify the expression of intracellular proteins, present within the cells, the proteins need to be extracted. In general, the cell membrane is lysed by detergents, enzymes or physical agitation, sequentially precipitated and re-suspended in a suitable buffer. The concentration of the protein sample can be determined by using fluorescent assays or colorimetric assays. Extracted proteins can then be used for downstream applications such as western blots to probe against and quantify proteins of interest.

5.4.5.1.1 Methodology

Intracellular proteins were extracted using radioimmunoprecipitation assay buffer (RIPA) cell lysis buffer (Pierce#89900) following the protocol provided by the manufacturer. Briefly, SGBS cells were differentiated in 6 well plates as described in section 5.4.1.1 and on day 14, the media was decanted, and mature adipocytes were washed with ice cold

PBS twice. 100µL RIPA buffer mix (1 SigmaFast protease inhibitor cocktail tablet (Sigma#S8830) and 10 PhosSTOP tablets (Roche#04906845001) in 100mL of RIPA buffer) per well was added, and the cells were incubated for 5 minutes on ice, with gentle agitation occasionally. The cells were scrapped, and the cell suspension was centrifuged for 15 minutes at 14,000xg, 4°C. The supernatants were transferred into new tubes and used for protein quantification (as described in section 5.4.5.2.1). Following quantification, the samples were in diluted equal volumes of Laemmli buffer (250mM Tris-hydrochloric acid-pH 6.8, 10% glycerol, 0.006% bromophenol blue, 4% sodium dodecyl sulfate, β-mercaptoethanol-pH 6.8), incubated at 100°C for 5 minutes and stored at -20°C.

5.4.5.2 Bicinchoninic acid assay

Protein quantification is important for standardisation of data. The basic used test for protein quantification is the biuret reaction; a chemical test used for detecting the presence of peptide bonds. In the presence of peptides, cupric form of copper is reduced to cuprous form of copper forming a violet-coloured coordination complexes in an alkaline solution. The darker the colour, the higher the protein concentration. The Pierce™ Bicinchoninic acid (BCA) is a colorimetric assay which based on the same principles. It is a two-step, high precision and a sensitive test with a high linearity range (20-2000µg).

The first step of the assay is the binding of copper to a protein in an alkaline environment containing sodium potassium tartrate to form a light blue complex; a cuprous cation (biuret reaction). In the second step, BCA reacts with the cuprous cation formed in step one. The intense purple-colour reaction product results from binding of two BCA

molecules to one cuprous ion. The BCA/copper complex is water-soluble and exhibits a strong linear absorbance at 562nm with increasing protein concentrations.

5.4.5.2.1 Methodology

The Pierce BCA protein assay kit (Thermofisher#23227) was used to quantify protein concentrations by following the protocol provided by the manufacturer. Briefly, bovine serum albumin (BSA) was used to prepare a serial dilution (0-2000 μ L/mL) of known standards as described in the protocol. The BCA working reagent (WR) was prepared following a 50:1 ratio of reagent A and reagent B. 10 μ L of each standard and unknown sample was added in replicates into a 96 well microplate. 200 μ L of WR was added to each well (sample to WR ratio=1:20) and the plate was mixed thoroughly on a plate shaker for 30 seconds. The plate was covered with aluminium foil and incubated at 37°C for 30 minutes, cooled to room temperature and measured the absorbance at 562nm on a spectrophotometer. The average absorbance measurement of blank standard was subtracted from the average absorbance measurements of all individual standards and unknown samples. A standard curve was generated, and the trend line equation used to determine the protein concentration of each unknown sample.

5.4.5.3 Western blotting

Western Blotting is an important technique used in cellular and molecular biology. It was introduced by (Towbin, Staehelin et al. 1979) and is now is used routinely for protein analysis. The specificity of the antibody-antigen interaction enables the researchers to identify specific proteins from a mixture of proteins extracted from cells.

The first step in a western blotting is separation of macromolecules in the sample using gel electrophoresis based on their molecular weight. Separated molecules are transferred

or blotted onto a second matrix; a nitrocellulose or polyvinylidene difluoride membrane. The membrane is blocked using 5% BSA or Milk powder to prevent any nonspecific binding of antibodies to the surface. The transferred protein is then probed using antibody specific to the protein of interest (primary antibody) and antibody specific to the host species of the primary antibody (secondary antibody) labelled with a conjugate molecule; biotin, fluorescent probes and enzymes. The most commonly used enzymes are horseradish peroxidase (HRP), and alkaline phosphatase (AP). The HRP-conjugated antibodies are better than AP-conjugated antibodies because HRP is smaller in size, higher activity rate and lower cost. These conjugated molecules produce a detectable signal with combined with an appropriate substrate; chromogenic, fluorescent or chemiluminescent. Use of a chemiluminescent substrate results in the production of light as a by-product (when enzyme reacts with substrate) which can be captured using film or digital imaging instruments (cameras). The data can then be quantified using densitometry software's such as, Image J.

5.4.5.3.1 Gel electrophoresis

The protein samples prepared in section 5.4.5.1.1 were further diluted with Laemmli buffer to 40µg or 60µg to a total volume of 20µL per well. The samples and precision plus protein western blotting standard (Bio-rad#1610376) were centrifuged at 1200xg for 2 minutes at 4°C and loaded into 4-20% Precast TGX™ (Tris/glycine) gradient gels (Bio-Rad, UK) in a Mini-PROTEAN™ vertical electrophoresis units filled with 1x running buffer (Pierce #28362). The gels were run at 150V for 90 minutes using Power-Pac 3000 (Bio-Rad, UK).

5.4.5.3.2 Protein transfer

The protein containing polyacrylamide gels were carefully removed and sandwiched into Trans-Blot® Turbo™ transfer packs consisting of filter paper, buffer and a polyvinylidene difluoride membrane. The assembled cassettes were loaded into the Trans-Blot system (Bio-Rad, UK) and run for mixed molecular weight transfer for 7 minutes. Electrophoretic transfer is commonly used because of its speed and transfer efficiency.

5.4.5.3.3 Antibody probing

The antibodies used in this study were; anti p-ERK/total-ERK (CST#9101S/#9102S) at a concentration of 1:2000, anti-p-p38/total-p38 (NEB#9211s) at a concentration of 1:1000, anti p-ACC/total-ACC (NEB#3661s/Sigma#SAB4501395) at a concentration of 2:1000 and anti-rabbit HRP IgG (CST#7074) at a concentration of 1:10,000.

The membranes were blocked in a blocking buffer (5% w/v milk powder in tris-buffered saline (Fisher Scientific #10776834) with 1% tween 20 (TBS/T)) and incubated at room temperature for 1 hour on an orbital shaker, with the protein side facing downwards. The membranes were washed with TBS/T three times for 5 minutes each and incubated with primary antibody (5% w/v BSA in TBS/T) at 4°C overnight on a roller shaker. The membranes were then washed with TBS/T three times for 5 minutes each to remove any unbound primary antibody and incubated with secondary antibody (5% w/v milk powder in TBS/T) at room temperature for 1 hour on an orbital shaker. The membranes were washed with TBS/T up to five times for 5 minutes each before visualising.

5.4.5.3.4 Visualisation, densitometry and quantification

The membranes were placed protein side up onto acetate sheets and 1mL of the substrate solution (1:1 solutions of SuperSignal West Femto Kit (Thermo Scientific#34095)) were

poured onto the surface of each membrane to amplify the signals. The membranes were visualised using a ChemiDoc transilluminator for imaging (Bio-Rad, UK) and visualised using the ImageLab software. The membranes were auto exposed to detect dense bands corresponding to the protein of interest, captured and analysed using Image J. The image was opened in image J and boxes were drawn around the bands of interest. The density of the band corresponds to the peak on the plot, thus the area under the peak was measured and used for data analysis.

5.4.6 Lipid peroxidation assay

Lipid peroxidation, also known as lipid oxidation, is a process of oxidative degradation of lipids in which free radicals are take electrons from lipids resulting in cell damage. Quantification of lipid peroxidation is used to assess oxidative stress. The end-product of the reaction, malondialdehyde (MDA) forms adduct with Thiobarbituric Acid (TBA) can be detected and quantified to assess the rate of lipid peroxidation.

5.4.6.1 Methodology

The lipid peroxidation assay (Abcam#136955) was carried out following the protocol provided by the manufacturer. Briefly, SGBS cells were differentiated in 96 well plates as described in section 5.4.1.1 and on day 14, mature adipocytes were washed with ice cold PBS, homogenised with 303µL lysis buffer (300µL MDA lysis buffer + 3µL butylhydroxytoluene (100X)) on ice, centrifuged at 13,000xg for 10 minutes and the supernatants were transferred into fresh Eppendorf tubes. The MDA standards (0-20 nmol) were prepared, 600µL of TBA reagent was added to each Eppendorf containing 200µL of standards and samples, incubated at 95°C for 60 minutes and cooled to room temperature on ice for 10 minutes. 200µL of each standard and samples were plated in

triplicates in 96 well plates and absorbance were immediately measured using a spectrophotometer at optical density 532nm. The absorbance of all samples was corrected by subtracting the mean absorbance value of the blank from all standards and samples. The standard curve was generated, and the trend line equation was used to calculate the rate of lipid peroxidation in the samples.

5.4.7 Glucose uptake assay

Glucose uptake is an important biological function that can be used to study insulin activity. Glucose analog, 2-deoxyglucose (2-DG) is widely used for measuring glucose uptake because it is structurally like glucose. Like glucose, 2-DG can be taken up by glucose transporters and metabolised to 2-DG-6 phosphate (2-DG6P). However, 2-DG6P cannot be further metabolised and thus accumulates in the cells. The accumulated 2-DG6P is oxidized to generate nicotinamide adenine dinucleotide phosphate (NAPDH) resulting in an oxidized substrate, which can be measured at optical density of 412nm. The absorbance value is directly proportional to 2-DG (or glucose) uptake by the cells.

5.4.7.1 Methodology

The glucose uptake assay (Abcam#136955) was carried out following the protocol supplied by the manufacturer. Briefly, SGBS cells were differentiated in 96 well plates as described in section 5.4.1.1 and maintained for further 4 days. The cells were incubated in 100µL serum free media (Dulbecco's modified Eagle medium/Nutrient mixture F-12 + 1% anti-anti) per well overnight at 37°C (95% O₂/5% CO₂). They were washed twice with PBS and starved for glucose by incubating in 100µL Krebs-Ringer-Phosphate-Hepes F(20mM HEPES, 5mM potassium phosphate monobasic, 1mM magnesium sulfate, 1mM calcium chloride, 136mM sodium chloride and 4.7 mM potassium chloride; pH 7.4) +

2% BSA buffer per well for 40 minutes, at 37°C (95% O₂/5% CO₂). The cells were stimulated with 1µM insulin, 10µM WRW4, 10µM AC2-26 or vehicle controls and incubated at 37°C (95% O₂/5% CO₂) for 20 minutes, followed by 10mM 2-DG incubation at 37°C (95% O₂/5% CO₂) for a further 20 minutes. A 2-DG6P standard curve (0-100pmol/well) was prepared and NADPH was generated from 2DG6P. Endogenous NAD(P) was degraded to reduce background signals and a recycling amplification reaction was carried out and the uptake of 2-DG was measured at 412nm using a spectrophotometer in a kinetic mode, every 5 minutes for 20 minutes. The absorbance of all samples was corrected by subtracting the mean absorbance value of the blank from all standards and samples. The standard curve was generated, and the trend line equation was used to calculate the amount of 2-DG6P in the samples.

5.4.8 Data and statistical analysis

The data in this study is presented as Mean \pm Standard Error of the Mean (SEM). One tailed, unpaired student's T-test was used to statistically analyse the data and to assess the differences in the data sets, a p value of $p < 0.05$ was considered statistically significant. GraphPad Prism 5 was used to statistically analyse the data and produce the graphs.

5.5 Results

5.5.1 Lipid accumulation in SGBS cells differentiated with and without FPR2/ALX receptor antagonists and/or agonist

SGBS cells were seeded and induced to differentiate for 14 days to reach a mature adipocyte phenotype. A key phenotypic hallmark of adipocyte differentiation is the accumulation of lipid droplets. Cells at Day 0, 4, 8 and 14 were stained with ORO and assessed for lipid accumulation. Figure 5.2a displays a pictorial representation of the accumulation of lipid droplets over the 14-day period. As expected, small lipid droplets were first observed at day 4 indicative of adipogenic capacity of these cells. At day 0, 20.5% \pm 2.8, n=3 of the area analysed showed lipid staining which significantly increased at day 4; 57.9% \pm 4.7, $p < 0.0001$, n=3. Lipids continue to significantly accumulate through to day 8; 69.6% \pm 4.1, $p < 0.001$, n=3 and significantly increased at day 14; 100% \pm 5.1, $p < 0.0001$, n=6, as shown in Figure 5.2b.

SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 or DMSO as vehicle control. Cells at Day 0, 4, 8 and 14 were stained with ORO and assessed for lipid accumulation. A significant increase in lipid accumulation at day 4 was observed in comparison to day 0; 50.2% \pm 4.0 vs 20.5% \pm 2.8 respectively, $p < 0.0001$, n=3, as shown in Figure 5.3b. Similarly, a significant increase was observed at day 8; 70.8% \pm 3.4, $p = 0.0001$, n=3, however, no significant difference was observed between day 8 and day 14. Both the vehicle control and WRW4 differentiated cells showed similar lipid accumulation pattern and phenotype at day 14. Interestingly, no significant difference in lipid accumulation between vehicle control and WRW4 differentiated cells was observed during the early stages of differentiation (Days 0-8), however, lipid accumulation

significantly decreased in WRW4 differentiated cells compared to vehicle control; 78.0% \pm 4.7 vs 100% \pm 5.1, $p=0.0016$, $n=6$, respectively at day 14, as shown in Figure 5.3b.

Inhibiting the activity of FPR2/ALX using a specific receptor antagonist demonstrated a significant decrease in lipid accumulation therefore, it was thought FPR2/ALX activation using the receptor agonist may increase lipid accumulation, suggesting it plays a positive role in adipocyte differentiation. SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M AC2-26 or vehicle control. A significant increase in lipid accumulation at day 4 was observed in comparison to day 0; 48.9% \pm 4.2 vs 20.5% \pm 2.8 respectively, $p<0.0001$, $n=3$, as shown in Figure 5.4b. Similarly, a significant increase was observed at day 8; 70.3% \pm 3.6, $p=0.0001$, $n=3$, however, no significant difference was observed between day 8 and day 14. Like WRW4 differentiated cells, no significant difference in lipid accumulation between vehicle control or AC2-26 differentiated cells were observed during the early stages of differentiation (Days 0-8). However, lipid accumulation significantly decreased in AC2-26 differentiated cells compared to vehicle control; 79.6% \pm 3.6 vs 100% \pm 5.1, $p=0.0013$, $n=6$, respectively at day 14, as shown in Figure 5.4b.

Although ANXA1 is a specific agonist for FPR2/ALX, its *N* terminal peptide AC2-26 can also bind to and activate other members of the FPR family such as FPR1 (Woloszynek, Hu et al. 2012). Antagonising FPR2/ALX may redirect the binding of endogenous AC2-26 to FPR1 and instigating a decrease in lipid accumulation, as shown in Figure 5.28. Thereby, FPR1 activation with the treatment of AC2-26 may override the effects of FPR2/ALX activation presenting equivalent results. Thus, activating FPR1 and inhibiting FPR2/ALX simultaneously, may support to identify if FPR1 influences lipid accumulation, shown in Figure 5.28. Thus, SGBS cells were seeded and induced to

differentiate with 10 μ M WRW4 and 10 μ M AC2-26 or vehicle control over 14 days. The cells were stained with ORO and assessed for lipid accumulation. A significant increase in lipid accumulation at day 4 was observed in comparison to day 0; 54.5% \pm 4.6 vs 20.5% \pm 2.8 respectively, $p < 0.0001$, $n=3$, as shown in Figure 5.5b. Similarly, a significant increase was observed at day 8; 73.2% \pm 3.7, $p=0.0030$, $n=3$, however, no significant difference was observed between day 8 and day 14. No significant difference in lipid accumulation between vehicle control and WRW4 + AC2-26 differentiated cells was observed during the early stages of differentiation (Days 0-8). However, WRW4 + AC2-26 differentiated cells accumulated significantly less lipids than vehicle control; 77.2% \pm 5.2 vs 100% \pm 5.1, $p=0.0011$, $n=6$, respectively at day 14, as shown in 5.5b.




Figure 5.2: Lipid accumulation during differentiation of SGBS cells. SGBS cells were differentiated and lipid droplets were stained using Oil Red O dye at Day 0, 4, 8 and 14. a) Visual representation of lipid droplets indicated by white arrows (magnification x10). b) Oil Red O staining was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. Significant increase in lipid accumulation was observed at day 14 in comparison to day 0 ($p < 0.0001$, $n = 6$). Data represents as Mean \pm SEM normalised to day 14. P value was set at $P < 0.05$ and is donated by * = Day 0 vs, # = Day 4 vs and \$ = Day 8 vs.




Figure 5.3: Lipid accumulation during differentiation of SGBS cells treated with FPR2/ALX antagonist; WRW4. SGBS cells were differentiated with and without 10 μ M WRW4 and lipid droplets were stained using Oil Red O at Day 0, 4, 8 and 14. a) Visual representation of lipid droplets indicated by white arrows (magnification x10). b) Oil Red O staining was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. Significant increase in lipid accumulation was observed at day 14 in comparison to day 0 ($p < 0.0001$, $n = 6$) in WRW4 differentiated cells. No significant difference in lipid accumulation was observed between vehicle control and WRW4 differentiated cells between day 0 and 8. However, at Day 14, lipid accumulation was significantly less in WRW4 differentiated cells compared to vehicle control ($p = 0.0367$, $n = 6$). Data represents as Mean \pm SEM normalised to vehicle control. P value was set at $P < 0.05$ and is denoted by ***= Day 0 vs, ####= 4 vs. Histogram: Black= control, Grey= treated




Figure 5.4: Lipid accumulation during differentiation of SGBS cells treated with FPR2/ALX agonist; AC2-26. SGBS cells were differentiated with and without 10 μ M AC2-26 and lipid droplets were stained using Oil Red O at Day 0, 4, 8 and 14. a) Visual representation of lipid droplets indicated by white arrows (magnification x10). b) Oil Red O staining was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. Significant increase in lipid accumulation was observed at day 14 in comparison to day 0 ($p < 0.0001$, $n = 6$) in AC2-26 differentiated cells. No significant difference in lipid accumulation was observed between vehicle control and AC2-26 differentiated cells between day 0 and 8. However, at Day 14, lipid accumulation was significantly less in AC2-26 differentiated cells compared to vehicle control ($p = 0.0210$, $n = 6$). Data represents as Mean \pm SEM normalised to vehicle control. P value was set at $P < 0.05$ and is denoted by ***= Day 0 vs, ####= 4 vs. Histogram: Black= control, Grey= treated




Figure 5.5: Lipid accumulation during differentiation of SGBS cells treated with FPR2/ALX antagonist; WRW4 and agonist; AC2-26. SGBS cells were differentiated with and without 10 μ M WRW4 + AC2-26 and lipid droplets were stained using Oil Red O at Day 0, 4, 8 and 14. a) Visual representation of lipid droplets indicated by white arrows (magnification x10). b) Oil Red O staining was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. Significant increase in lipid accumulation was observed at day 14 in comparison to day 0 ($p < 0.001$, $n = 6$) in WRW4 + AC2-26 differentiated cells. No significant difference in lipid accumulation was observed between vehicle control and WRW4 + AC2-26 differentiated cells between day 0 and 8. However, at Day 14, lipid accumulation was significantly less in WRW4 + AC2-26 differentiated cells compared to vehicle control ($p = 0.0109$, $n = 6$). Data represents as Mean \pm SEM normalised to vehicle control. P value was set at $P < 0.05$ and is denoted by ***= Day 0 vs, ####= 4 vs. Histogram: Black= control. Grey= treated

Lipid accumulation in mature adipocytes differentiated with WRW4, AC2-26 or WRW4 + AC2-26, was not significantly different between each treatment, as shown in Figure 5.6.




Figure 5.6: Comparison of lipid accumulation in mature adipocytes differentiated with FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 + AC2-26 and lipid droplets were stained using Oil Red O at Day 14. Oil Red O staining was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. No Significant difference was observed between mature adipocytes differentiated with WRW4, AC2-26 or WRW4 + AC2-26. Data represents as Mean \pm SEM

To further understand the role of FPR1 in adipogenesis, a non-selective FPR antagonist; BOC-2 was used. It has a higher affinity towards FPR1, therefore at lower concentrations it binds and inhibits signalling through FPR1, however, at higher concentrations it binds and inhibits signalling through FPR2/ALX also thus, completely inhibiting ANXA1 signalling (Woloszynek et al. 2012). SGBS cells were seeded and induced to differentiate for 14 days with 0-100 μ M BOC-2. At day 14, the cells were stained with ORO and assessed for lipid accumulation, as shown in Figure 5.7a. No significant difference in lipid

accumulation was observed in cells differentiated with 1 μ M BOC-2 at day 14, as shown in Figure 5.7b. However, lipid accumulation in cells differentiated with 10 μ M BOC-2 significantly increased compared to vehicle control; 120.6 \pm 4.3 vs 100% \pm 6.5 respectively, $p=0.0089$, $n=5$ at day 14. Furthermore, lipid accumulation in cells differentiated with 100 μ M BOC-2 was not significantly altered compared vehicle control (105.8 \pm 4.8, $p=0.02364$, $n=5$) at day 14.




Figure 5.7: Lipid accumulation in mature adipocytes differentiated with FPR antagonist; BOC-2. SGBS cells were differentiated with 0-100μM BOC-2 and lipid droplets were stained using Oil Red O at Day 14. Oil Red O staining was analysed using image J One tailed unpaired student's T-test was used to statistically analyse the data. a) Visual representation of lipid droplets indicated by white arrows (magnification x10). B) Mature adipocytes differentiated with 10μM accumulated significantly more lipids compared to vehicle control differentiated mature adipocytes ($p=0.0089$, $n=5$). Yet, lipid accumulation in adipocytes differentiated with 100μM did not differ compared to vehicle control differentiated mature adipocytes. Data represents as Mean \pm SEM normalised to 0μM BOC-2. P value was set at $P<0.05$ and is denoted by * vs 0.

5.5.2 Morphology of lipid droplets in mature adipocytes differentiated with and without FPR2/ALX receptor antagonist and/or agonist

The average number of lipid droplets in WRW4 differentiated mature adipocytes was not significantly different compared to vehicle control (96.9 ± 11.5 vs 100 ± 4.0 , $p=0.1595$, $n=4$, respectively), as shown in Figure 5.8. Though the average size of lipid droplets was decreased, it was not significantly different ($84.47 \text{ pixel}^2 \pm 7.0$ vs $100 \text{ pixel}^2 \pm 6.9$, $p=0.0638$, $n=4$, respectively), as shown in Figure 5.9. However, further analysis revealed, on average, that these cells accumulated a significantly increased number of smaller lipid




Figure 5.8: Average number of Lipid droplets in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and agonist; AC2-26. SGBS cells were differentiated with $10\mu\text{M}$ WRW4 and/or $10\mu\text{M}$ AC2-26 and lipid droplets were stained using Oil Red O at Day 14. 10 images per well of a 6 well plate was taken, and the average number of lipid droplets were analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. WRW4 differentiated mature adipocytes did not show a difference in the average number of lipid droplets compared to vehicle control ($p=0.1595$), whereas, those differentiated with AC2-26 had significantly less lipid droplets compared to vehicle control ($p=0.0094$, $n=4$). Similarly, WRW4 + AC2-26 differentiated mature adipocytes contained significantly less number of lipid droplets compared to vehicle control ($p=0.0045$, $n=4$). No statistical difference was observed between treated cells. Data represents as Mean \pm SEM normalised to control. P value was set at $P<0.05$ and denoted by * vs control. Histogram: Black= control, Grey= treated

droplets (between 100-1000 pixel²) (2.3 ± 0.2 vs 1.9 ± 0.1 , $p=0.0462$, $n=4$) and a significantly decreased number of larger lipid droplets (1000+ pixel²) (0.5 ± 0.1 vs 0.6 ± 0.1 , $p=0.0200$, $n=4$) compared to vehicle control (Figure 5.10).




Figure 5.9: Average size of lipid droplets in mature adipocytes differentiated with or without FPR2/ALX antagonist; WRW4 and agonist; AC2-26 SGBS cells were differentiated with 10 μ M WRW4 and/or 10 μ M AC2-26 and lipid droplets were stained using Oil Red O at Day 14. 10 images per well of a 6 well plate was taken, and the average size of lipid droplets were analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. WRW4 differentiated mature adipocytes showed a decrease in the average size of lipid droplets compared to vehicle control, however, it was not statically different ($p=0.0638$, $n=4$). AC2-26 differentiated mature adipocytes showed no difference in the average size of lipid droplets compared to vehicle control. Similarly, WRW4 + AC2-26 differentiated mature adipocytes showed a decrease in the average size of the lipid droplets compared to vehicle control; however, it was not statically different ($p=0.0575$, $n=4$). Data represents as Mean \pm SEM normalised to control. Histogram: Black= control, Grey= treated

However, the average number of lipid droplets in AC2-26 differentiated mature adipocytes were significantly reduced compared to vehicle control (87.1 ± 3.1 vs 100 ± 3.9 , $p=0.0094$, $n=4$, respectively), as shown in Figure 5.8, but the average size was not different in these adipocytes compared to vehicle control ($94.6 \text{ pixel}^2 \pm 7.9$ vs $100 \text{ pixel}^2 \pm 6.9$, $p=0.3048$, $n=4$, respectively), as shown in Figure 5.9. Furthermore, on average, no

difference was observed in the different sizes of lipid droplets between these adipocytes and vehicle control (Figure 5.11).

Similarly, the average number of lipid droplets were significantly reduced in WRW4 + AC2-26 differentiated mature adipocytes compared to vehicle control (76.9 ± 7.2 vs 100 ± 3.9 , $p=0.0045$, $n=4$, respectively), as shown in Figure 5.8. The average size of lipid droplets was reduced in these adipocytes compared to vehicle control, however, it was not statistically significant (85.3 ± 5.7 vs 100 ± 6.9 , $p=0.0575$, $n=4$, respectively), as shown




Figure 5.10: Number of lipid droplets of different sizes in mature adipocytes differentiated with FPR2/ALX antagonist; WRW4. SGBS cells were differentiated with 10 μ M WRW4 and lipid droplets were stained using Oil Red O at Day 14. 10 images per well of a 6 well plate was taken, and the lipid droplet sizes were categorically analysed and normalised against the total count per image. One tailed unpaired student's T-test was used to statistically analyse the data. WRW4 differentiated mature adipocytes contained significantly increased number smaller lipid droplets (100-1000 pixel²) compared to vehicle control ($p=0.0462$, $n=4$) and significantly decreased number of larger lipid droplets (1000+ pixel²); $p=0.0200$, $n=4$. Data represented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs control. Histogram: Black= control, Grey= treated

in Figure 5.9. However, on average, these cells accumulated a significantly increased number of smaller lipid droplets (10-100 pixel²) (24.2 ± 1.1 vs 20.9 ± 1.1 , $p=0.0216$, $n=4$)

and a significantly decreased number of larger lipid droplets (1000+ pixel²) (0.41 ± 0.1 vs 0.2 ± 0.1 , $p=0.0407$, $n=4$) compared to vehicle control, as shown in Figure 5.12.




Figure 5.11: Number of lipid droplets of different sizes in mature adipocytes differentiated with FPR2/ALX agonist; AC2-26. SGBS cells were differentiated with 10 μ M AC2-26 and lipid droplets were stained using Oil Red O at Day 14. 10 images per well of a 6 well plate was taken, and the lipid droplet sizes were categorically analysed and normalised against the total count per image. One tailed unpaired student's T-test was used to statistically analyse the data. No difference in the size of lipid droplets was observed between AC2-26 differentiated mature adipocytes and vehicle control ($n=4$). Data represented as Mean \pm SEM. Histogram: Black= control, Grey= treated




Figure 5.12: Average number of lipid droplets of different sizes in mature adipocytes differentiated with FPR2/ALX antagonist; WRW4 and agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and 10 μ M AC2-26 and lipid droplets were stained using Oil Red O at Day 14. 10 images per well of a 6 well plate was taken, and the lipid droplet sizes were categorically analysed and normalised against the total count per image. One tailed unpaired student's T-test was used to statistically analyse the data. WRW4 + AC2-26 differentiated mature adipocytes contained significantly increased number smaller lipid droplets (10-100 pixel²) compared to vehicle control ($p=0.0216$, $n=4$) and significantly decreased number of larger lipid droplets (1000+ pixel²); $p=0.0407$, $n=4$. Data represented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs control. Histogram: Black= control, Grey= treated

Moreover, on average, AC2-26 only differentiation of mature adipocytes resulted in a significantly increased number of smaller lipid droplets (0-10 pixel²) compared to WRW4 + AC2-26 differentiated mature adipocytes; $p=0.0436$. Whereas, WRW4 + AC2-26 differentiated mature adipocytes showed increased number of 10-100 pixel² lipid droplets compared to AC2-26 only differentiated mature adipocytes ($p= 0.0165$). The average number of lipid droplets between 100-1000+ were significantly altered in all conditions, as shown in Figure 5.13. AC2-26 only differentiated mature adipocytes and WRW4 + AC2-26 differentiated mature adipocytes accumulated significantly less 100-1000 pixel²

lipid droplets ($p=0.0143$ and $p<0.0001$, $n=4$, respectively) compared to WRW4 only differentiated mature adipocytes. Similarly, WRW4 + AC2-26 differentiated mature adipocytes contained significantly fewer 100-1000 pixel² lipid droplets compared to AC2-26 only differentiated mature adipocytes; $p=0.0277$, $n=4$. AC2-26 only differentiated adipocytes contained significantly increased number of larger lipid droplets (1000+ pixel²) compared to WRW4 only differentiated adipocytes ($p=0.0169$, $n=4$) and WRW4 + AC2-26 differentiated adipocytes ($p<0.0001$, $n=4$). Whereas, WRW4 + AC2-26 differentiated mature adipocytes contained significantly fewer large lipid droplets




Figure 5.13: Comparison of average number of lipid droplets of different sizes in mature adipocytes differentiated with or without FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and/or 10 μ M AC2-26 and lipid droplets were stained using Oil Red O at Day 14. 10 images per well of a 6 well plate was taken, and the lipid droplet sizes were categorically analysed using Image J and normalised against the total count per image. One tailed unpaired student's T-test was used to statistically analyse the data. Data represented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by *.

(1000+ pixel²) compared to WRW4 only differentiated mature adipocytes (p=0.0015, n=4).

5.5.3 Cytotoxicity of antagonist and agonist compounds in mature adipocytes

To ensure the results observed were a result of the receptor antagonist and agonist and not due to cellular cytotoxicity of the compounds, cell viability was assessed using the AlamarBlue cell viability assay, as described in section 5.4.3.




Figure 5.14: Cell viability in mature adipocytes differentiated with and without FPR2/ALX antagonist: WRW4 and/or agonist: AC2-26. SGBS cells were differentiated with 10μM WRW4 and/or 10μM AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, the cells were incubated with 10% AlamarBlue for 4 hours at 37C. The absorbance was measured using a spectrophotometer at 562nm and 600nm as a reference wavelength. Cell viability was normalised against control and is presented as %. One tailed unpaired student's T-test was used to statistically analyse the data. No significant difference in % cell viability is observed between vehicle control and WRW4 differentiated mature adipocytes (p=0.1385, n=4), vehicle control and AC2-26 differentiated mature adipocytes (p=0.3622, n=4) and vehicle control and WRW4 + AC2-26 differentiated mature adipocytes (0.2688, n=4). Similarly, no difference in % cell viability as observed between treated mature adipocytes. Data represents as Mean ± SEM normalised to control. Histogram: Black= control, Grey= treated

SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 and/or 10 μ M AC2-26 or vehicle control and cellular viability was assessed, as shown in Figure 5.14. No significant difference in percentage of cell viability was observed between vehicle control and WRW4 differentiated mature adipocytes; 100% \pm 5.6 vs 82.7% \pm 15.1, $p=0.01385$, $n=4$, respectively. Similarly, no difference in percentage of cell viability was observed between vehicle control and AC2-26 differentiated mature adipocytes; 100% \pm 5.6 vs 103.5% \pm 8.3, $p=0.3622$, $n=4$, respectively. Furthermore, no difference in percentage of cell viability was observed between vehicle control and WRW4 + AC2-26 differentiated mature adipocytes; 100% \pm 5.6 vs 93.6% \pm 8.3, $p=0.2688$, $n=4$, respectively. No difference in percentage of cell viability was observed between treatment differentiated mature adipocytes.

5.5.4 Intracellular signalling pathway activation in Mature Adipocytes differentiated with and without FPR2/ALX receptor antagonist and/or agonist

To investigate whether endogenous AC2-26 binds to FPR1 upon FPR2/ALX antagonist and agonist treatment, intracellular signalling pathways of FPR1 and FPR2/ALX were analysed. ANXA1/AC2-26 binding to and activating FPR2/ALX is thought to activate ERK1/2 MAPK whereas, AC2-26 binding to and activating FPR1 is thought to activate p38 mitogen activated protein kinase (p38 MAPK). SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 and/or 10 μ M AC2-26 or vehicle control and phosphorylation of ERK1/2 MAPK and p38 MAPK were analysed.

The percentage of phosphorylated ERK1/2 MAPK (%p-ERK1/2) was slightly decreased in mature adipocytes differentiated with WRW4 and/or AC2-26, however, it was not statistically significant compared to vehicle control, as shown in Figure 5.15. %p-ERK1/2

was decreased to $92.8\% \pm 14.8$ in WRW4 differentiated mature adipocytes compared to vehicle control; $100\% \pm 8.3$, $p=0.3409$, $n=5$. Similarly, %p-ERK1/2 was decreased to $95.2\% \pm 13.9$ in AC2-26 differentiated mature adipocytes compared to vehicle control; $100\% \pm 8.3$, $p=0.3879$, $n=5$. Furthermore, %p-ERK1/2 was decreased to $97.4\% \pm 14.4$ in WRW4 + AC2-26 differentiated mature adipocytes compared to vehicle control; $100\% \pm 8.3$, $p=0.4369.16$, $n=5$. Consequently, no difference was found between differently treated mature adipocytes.

The percentage of phosphorylated p38 MAPK (%p-p38) was significantly decreased in WRW4 differentiated mature adipocytes ($64.8\% \pm 15.1$ vs $100\% \pm 6.3$, $p=0.0317$, $n=5$), in AC2-26 differentiated mature adipocytes ($71.4\% \pm 4.7$ vs $100\% \pm 6.3$, $p=0.0024$, $n=6$) and in WRW4 + AC2-26 differentiated mature adipocytes ($67.1\% \pm 15.1$ vs $100\% \pm 6.3$, $p=0.0394$, $n=5$) compared to vehicle control, as shown in Figure 5.16. However, no significant difference was observed between differently treated mature adipocytes.




Figure 5.15: ERK1/2 MAPK activation in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, the cells were lysed using RIPA buffer and intracellular proteins were extracted and the concentration determined using BCA assay. 40 μ g of protein per well was loaded into SDS gels along with the sample buffer and run for 90 minutes at 150V. The separated proteins were transferred onto a PVDF membrane. The membrane was incubated in 5% milk for an hour, washed with TBST x3 and incubated with 1:2000 primary anti-p-ERK1/2 antibody overnight. The membrane was washed with TBST x3 and incubated with 1:10,000 secondary anti-IgG antibody for 2 hours. It was washed with TBST x3 and visualised using the SuperSignal west femto kit. The antibodies were stripped the membrane using stripping buffer and the processes was repeated for ERK1/2 and β Actin. The density of the bands was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. % phosphorylated p-ERK MAPK was slightly decreased in mature adipocytes differentiated with WRW4 and/or AC2-26, however, it should no statistical difference; WRW4 differentiated mature adipocytes; $p=0.3192$, $n=5$, AC2-26 differentiated mature adipocytes; $p=0.3700$, $n=5$, and WRW4 + AC2-26 differentiated mature adipocytes; $p=0.4216$, $n=4$, compared to vehicle control. Data presented as Mean \pm SEM normalised to control. Histogram: Black= control, Grey= treated
ERK MAPK= Extracellular signal regulated kinase and MAPK=Mitogen activated protein kinase




Figure 4.16: p38 MAPK activation in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, the cells were lysed using RIPA buffer and intracellular proteins were extracted and the concentration determined using BCA assay. 40 μ g of protein per well was loaded into SDS gels along with the sample buffer and run for 90 minutes at 150V. The separated proteins were transferred onto a PVDF membrane. The membrane was incubated in 5% milk for an hour, washed with TBST x3 and incubated with 1:1000 primary anti-p38 antibody overnight. The membrane was washed with TBST x3 and incubated with 1:10,000 secondary anti-IgG antibody for 2 hours. It was washed with TBST x3 and visualised using the SuperSignal west femto kit. The antibodies were stripped the membrane using stripping buffer and the processes was repeated for p38 and β Actin. The density of the bands was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. % phosphorylated p-38 MAPK was significantly decreased in WRW4 differentiated mature adipocytes ($p=0.0317$, $n=5$), in AC2-26 differentiated mature adipocytes ($p=0.0024$, $n=6$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p=0.0394$, $n=5$) compared to vehicle control. However, no significant difference was found between treated cells. Data presented as Mean \pm SEM normalised to control. * vs control. Histogram: Black= control, Grey= treated
p38 MAPK= p38 mitogen activated protein kinase

5.5.5 *De novo* lipogenesis in mature adipocytes differentiated with and without FPR2/ALX receptor antagonist and/or agonist

Active p38 MAPK phosphorylates several transcription factors involved in lipogenesis including transcriptional factors and enzymes through *de-novo* lipogenesis. Therefore, transcriptional factors and enzymes involved in de-novo lipogenesis were assessed.




Figure 5.17: mRNA expression of transcription factors involved in lipogenesis in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist: AC2-26. SGBS cells were differentiated with 10μM WRW4 and/or 10μM AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, mRNA was extracted using Trisure and cDNA was synthesised. 80ng of cDNA was used in SYBR green mediated quantitative PCR. Ct values were obtained and used to calculate normalised gene expression using the Livak 2^{-ΔΔCT} equation (as described in Chapter 3). One tailed unpaired student's T-test was used to statistically analyse the data. mRNA expression of PPARγ (PPARG) was significantly upregulated in mature adipocytes compared to preadipocytes (p=0.0098, n=7) and significantly downregulated in WRW4 differentiated mature adipocytes (p<0.0001, n=5), in AC2-26 differentiated mature adipocytes (p=0.0008, n=5) and in WRW4 + AC2-26 differentiated mature adipocytes (0.0103, n=4) compared to vehicle control. Similarly, mRNA expression of SREB (SREBP) was significantly upregulated in mature adipocytes compared to preadipocytes (p=0.0467, n=8) and significantly downregulated in WRW4 differentiated mature adipocytes (p=0.0002, n=4), in AC2-26 differentiated mature adipocytes (p<0.0001, n=4) and in WRW4 + AC2-26 differentiated mature adipocytes (p=0.0037, n=4) compared to vehicle control. Preadipocytes and vehicle control normalised to 1. Data presented as Mean±SEM. P value was set at P<0.05 and is denoted by * vs 1. Histogram: Dark Grey= control, Grey= treated. Housekeeping gene:18s RNA PPARγ= Peroxisome proliferator activated receptor γ and SREBP= Sterol regulatory element-binding protein

SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 and/or 10 μ M AC2-26 and mRNA expression levels were studied.

As expected, mRNA expression of transcriptional factors; *PPARG* and *SREBP* were significantly upregulated in mature adipocytes compared to preadipocytes; 41.8 \pm 10.6-fold, $p=0.0098$, $n=7$ and 6.1 \pm 1.9-fold, $p=0.0467$, $n=8$ vs 1, $n=8$ respectively, as shown in Figure 5.17. Similarly, mRNA expression levels of lipogenic enzymes were significantly upregulated in mature adipocytes in comparison to preadipocytes, as shown in Figure 5.18. Gene expression levels of *ACC*, *FAS* and *GPAT* were significantly upregulated by 16.2 \pm 2.6-fold, $p=0.0013$, $n=8$, 70.2 \pm 20.5-fold, $p=0.0173$, $n=7$ and 80.0 \pm 34.7-fold, $p=0.0424$, $n=5$ respectively, compared to preadipocytes (1, $n=8$).

Interestingly, mRNA expression of PPAR γ (*PPARG*) and SREBP (*SREBP*) were significantly downregulated in WRW4 differentiated mature adipocytes (0.49 \pm 0.08-fold, $p<0.0001$, $n=5$ and 0.54 \pm 0.08-fold, $p=0.0002$, $n=4$, respectively), in AC2-26 differentiated mature adipocytes (0.38 \pm 0.1-fold, $p=0.0008$, $n=5$ and 0.36 \pm 0.1-fold, $p<0.0001$, $n=4$, respectively) and in WRW4 + AC2-26 differentiated mature adipocytes (0.64 \pm 0.1, $p=0.0103$, $n=4$ and 0.52 \pm 0.1, $p=0.0037$, $n=4$, respectively), compared to vehicle control (1, $n=5$), as shown in Figure 5.17.

Similarly, mRNA expressions of lipogenesis enzymes were dysregulated, as shown in Figure 5.18. mRNA expression of acetyl CoA carboxylase (*ACC*) was significantly downregulated in WRW4 differentiated mature adipocytes (0.73 \pm 0.1, $p=0.0154$, $n=4$), in AC2-26 differentiated mature adipocytes (0.20 \pm 0.1, $p<0.0001$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes (0.62 \pm 0.1, $p=0.0213$, $n=6$) compared to vehicle control (1, $n=6$). mRNA expression of fatty acid synthase (*FASN*) was significantly

downregulated in WRW4 differentiated mature adipocytes (0.62 ± 0.1 , $p=0.0081$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes (0.80 ± 0.1 , $p=0.0071$, $n=5$) compared to vehicle control (1, $n=5$). However, no significant difference was observed in the mRNA expression of *FAS* in AC2-26 differentiated mature adipocytes (1.10 ± 0.2 , $p=0.3540$, $n=4$) compared to vehicle control (1, $n=4$). mRNA expression of glycerol-3-phosphate acyltransferase (*GPAM*) was significantly upregulated in WRW4 differentiated mature adipocytes (2.69 ± 0.6 , $p=0.0270$, $n=7$), significantly downregulated in AC2-26 differentiated mature adipocytes (0.69 ± 0.1 , $p=0.0031$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes (0.67 ± 0.1 , $p=0.0144$, $n=4$) compared to vehicle control (1, $n=7$).





Figure 5.18: mRNA expression of enzymes involved in lipogenesis in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist: AC2-26. SGBS cells were differentiated with 10μM WRW4 and/or 10μM AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, mRNA was extracted using Trisure and cDNA was synthesised. 80ng of cDNA was used in SYBR green mediated quantitative PCR. Ct values were obtained and used to calculate normalised gene expression using the Livak 2^{-ΔΔCt} equation (as described in Chapter 3). One tailed unpaired student's T-test was used to statistically analyse the data. mRNA expression of ACC (ACC) was significantly upregulated in mature adipocytes compared to preadipocytes (p=0.0013, n=8) and significantly downregulated in WRW4 differentiated mature adipocytes (p=0.0154, n=4), in AC2-26 differentiated mature adipocytes (p<0.0001, n=5) and in WRW4 + AC2-26 differentiated mature adipocytes (0.0213, n=6) compared to vehicle control. Similarly, mRNA expression of FAS (FASN) was significantly upregulated in mature adipocytes compared to preadipocytes (p=0.0173, n=7) and significantly downregulated in WRW4 differentiated mature adipocytes (p=0.0081, n=5) and in WRW4 + AC2-26 differentiated mature adipocytes (p=0.0071, n=5) compared to vehicle control. However, no significant difference was observed in the mRNA expression of FASN in AC2-26 differentiated mature adipocytes compared to vehicle control (p=0.3540, n=4). mRNA expression of GPAT (GPAM) was significantly upregulated in mature adipocytes compared to preadipocytes (p=0.0424, n=5) and in WRW4 differentiated mature adipocytes (p=0.0270, n=7), whereas it was significantly downregulated in AC2-26 differentiated mature adipocytes (p=0.0031, n=5) and in WRW4 + AC2-26 differentiated mature adipocytes (p=0.0144, n=4) compared to vehicle control. Preadipocytes and vehicle control normalised to 1. Data presented as Mean±SEM. P value was set at P<0.05 and is denoted by * vs 1. Dark Grey= control, Grey= treated. Housekeeping gene: 18s RNA
ACC= acetyl CoA carboxylase, FAS= Fatty acid synthase and GPAT= Glycerol-3-phosphate acyltransferase 1

A similar pattern was observed at the proteomic level for ACC. The total percentage of ACC was significantly reduced in WRW4 differentiated mature adipocytes ($70.2\% \pm 9.5$, $p=0.00343$, $n=3$), in AC2-26 differentiated mature adipocytes ($86.2\% \pm 4.9$, $p=0.0146$, $n=4$) and in WRW4 + AC2-26 mature adipocytes ($63.8\% \pm 2.1$, $p<0.0001$, $n=3$) compared to vehicle control ($100\% \pm 15$, $n=4$, as shown in Figure 5.19. Furthermore, percentage phosphorylated ACC (%pACC) was also significantly decreased in WRW4 differentiated mature adipocytes ($70.2\% \pm 9.5$, $p=0.0343$, $n=5$), in AC2-26 differentiated mature adipocytes ($59.2\% \pm 10.2$, $p=0.0130$, $n=6$) and in WRW4 + AC2-26 differentiated mature adipocytes ($58.8\% \pm 13.1$, $p=0.0265$, $n=6$) compared to vehicle control ($100\% \pm 10$, $n=4$), as shown in Figure 5.19.






Figure 5.19: % protein expression of ACC in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, the cells were lysed using RIPA buffer and intracellular proteins were extracted and the concentration determined using BCA assay. 60 μ g of protein per well was loaded into SDS gels along with the sample buffer and run for 90 minutes at 150V. The separated proteins were transferred onto a PVDF membrane. The membrane was incubated in 5% milk for an hour, washed with TBST x3 and incubated with 2:1000 primary anti-p-ACC antibody overnight. The membrane was washed with TBST x3 and incubated with 1:10,000 secondary anti-IgG antibody for 2 hours. It was washed with TBST x3 and visualised using the SuperSignal west femto kit. The antibodies were stripped the membrane using stripping buffer and the processes was repeated for total ACC and β Actin. The density of the bands was analysed using Image J. One tailed unpaired student's T-test was used to statistically analyse the data. b) % of total protein was significantly decreased in WRW4 differentiated mature adipocytes ($p=0.0024$, $n=3$), in AC2-26 differentiated mature adipocytes ($p=0.0146$, $n=4$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p<0.0001$, $n=3$), compared to vehicle control. c) % phosphorylated p-ACC was significantly decreased in mature adipocytes differentiated with WRW4 and/or AC2-26, however, it should no statistical difference; WRW4 differentiated mature adipocytes ($p=0.0343$, $n=5$), in AC2-26 differentiated mature adipocytes ($p=0.0130$, $n=6$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p=0.0265$, $n=6$) compared to vehicle control. Data presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs control. Black= control, Grey= treated

5.5.6 Lipolysis in mature adipocytes differentiated with and without FPR2/ALX receptor antagonist and/or agonist

In addition to lipogenesis, lipolysis was also investigated. Triglycerides stored in lipid droplets in mature adipocytes are hydrolysed to produce glycerol and free fatty acids. Triglyceride and fatty acid recycling is important in metabolic regulation and is tightly regulated by enzymes such as phosphoenolpyruvate carboxykinase and lipases; adipose triglyceride lipase (*ATGL*), hormone sensitive lipase (*HSL*) and monoacylglycerol lipase (*MGL*).

SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 and/or 10 μ M AC2-26 and mRNA expression levels were studied. mRNA expression levels of *ATGL*, *HSL* and *MGL* were significantly upregulated in mature adipocytes; 5.45 \pm 1.8-fold, $p=0.0334$, $n=5$, 112.3 \pm 31.4-fold, $p=0.0109$, $n=6$ and 4.54 \pm 1.2-fold, $p=0.0175$, $n=5$ respectively compared to preadipocytes (1, $n=6$), as shown in Figure 5.20. However, mRNA expressions of these lipolytic enzymes were significantly downregulated in mature adipocytes differentiated with antagonist and/or agonist, as shown in Figure 5.20. mRNA expression of *ATGL* was significantly downregulated in WRW4 differentiated mature adipocytes (0.29 \pm 0.1-fold, $p<0.0001$, $n=5$), in AC2-26 differentiated mature adipocytes (0.38 \pm 0.1-fold, $p=0.0006$, $n=6$) and in WRW4 + AC2-26 differentiated mature adipocytes (0.50 \pm 0.1-fold, $p=0.0009$, $n=4$), compared to vehicle control (1, $n=6$). mRNA expression of *HSL* was significantly downregulated in WRW4 differentiated mature adipocytes (0.41 \pm 0.03-fold, $p<0.0001$, $n=5$) and in AC2-26 differentiated mature adipocytes (0.63 \pm 0.1-fold, $p=0.0205$, $n=6$) compared to vehicle control (1, $n=6$). Whereas, the mRNA expression levels of *HSL* were upregulated in WRW4 + AC2-26 differentiated mature adipocytes compared to vehicle control, but the difference was not

statistically significant (1.14 ± 0.2 -fold, $p=0.2451$, $n=6$). Furthermore, mRNA expression of *MGL* was significantly downregulated in WRW4 differentiated adipocytes (0.42 ± 0.1 -fold, $p=0.0004$, $n=7$) and in AC2-26 differentiated mature adipocytes (0.63 ± 0.1 -fold, $p=0.0205$, $n=6$) compared to vehicle control (1, $n=7$). Furthermore, mRNA expression levels were downregulated in WRW4 + AC2-26 differentiated mature adipocytes compared to vehicle control; however, the difference was not statistically significant (0.96 ± 0.2 -fold vs 1, $p=0.4046$, $n=4$).




Figure 5.20: mRNA expression of enzymes involved in lipolysis in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist: AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and/or 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, mRNA was extracted using Trisure and cDNA was synthesised. 80ng of cDNA was used in SYBR green mediated quantitative PCR. Ct values were obtained and used to calculate normalised gene expression using the Livak 2^{-ΔΔCt} equation (as described in Chapter 3). One tailed unpaired student's T-test was used to statistically analyse the data. mRNA expression levels of ATGL, HSL and MGL were significantly upregulated in mature adipocyte; $p=0.0334$, $n=5$, $p=0.0109$, $n=6$ and $p=0.0175$, $n=5$ respectively compared to preadipocytes. However, mRNA expression of these lipolytic enzymes were significantly downregulated in treated mature adipocytes. mRNA expression of ATGL was significantly downregulated in WRW4 differentiated mature adipocytes ($p<0.0001$, $n=5$), in AC2-26 differentiated mature adipocytes ($p=0.0006$, $n=6$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p=0.0009$, $n=4$), compared to DMSO control. mRNA expression of HSL was significantly downregulated in WRW4 differentiated mature adipocytes ($p<0.0001$, $n=5$) and in AC2-26 differentiated mature adipocytes ($p=0.0205$, $n=6$) compared to vehicle control. Whereas, the mRNA expression levels of HSL were upregulated in WRW4 + AC2-26 differentiated mature adipocytes compared to vehicle control, but the difference was not statistically significant ($p=0.2451$, $n=6$). Furthermore, mRNA expression of MGL was significantly downregulated in WRW4 differentiated adipocytes ($p=0.0004$, $n=7$) and in AC2-26 differentiated mature adipocytes ($p=0.0205$, $n=6$) compared to vehicle control. Furthermore, mRNA expression levels were downregulated in WRW4 + AC2-26 differentiated mature adipocytes compared to vehicle control, however, the difference was not statistically significant ($p=0.4046$, $n=4$). Preadipocytes and vehicle control normalised to 1. Data presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by *vs 1. Dark Grey= control, Grey= treated. Housekeeping gene: 18s RNA
ATGL= adipose triglyceride lipase, HSL= hormone sensitive lipase and MGL= monoacylglycerol lipase

MDA is a by-product of lipid peroxidation and its quantification is used as a marker of lipid peroxidation. In line with the downregulation of mRNA expression of lipolytic enzymes, lipid peroxidation was decreased in mature adipocytes differentiated with WRW4 and/or AC2-26, as shown in Figure 5.21.




Figure 5.21: Lipid peroxidation analysis in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist: AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and/or 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, the cells were lysed using lysis buffer and protein samples were incubated with Thiobarbituric acid for 60 minutes at 95C and the absorbance was measured using a spectrophotometer at 532nm. The data was normalised to control. One tailed unpaired student's T-test was used to statistically analyse the data. Significantly less MDA was present in WRW4 differentiated mature adipocytes ($p=0.0450$, $n=4$), in AC2-26 differentiated mature adipocytes ($p=0.0085$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p=0.0198$, $n=5$), compared to vehicle control. MDA concentration is proportionate to the level lipid peroxidation in mature adipocytes. Data normalised to control and presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs control. Black= control, Grey= treated. MDA= malondialdehyde

Significantly less MDA was present in WRW4 differentiated mature adipocytes ($0.84 \pm 0.1 \text{ nmol}$, $p=0.0450$, $n=4$), in AC2-26 differentiated mature adipocytes ($0.81 \pm 0.04 \text{ nmol}$, $p=0.0298$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes ($0.75 \pm 0.1 \text{ nmol}$, $p=0.0198$, $n=5$), compared to vehicle control ($1 \pm 0.1 \text{ nmol}$, $n=5$).

5.5.7 Glucose uptake in mature adipocytes differentiated with and without FPR2/ALX receptor antagonist and/or agonist

ANXA1 is thought to regulate insulin secretion and sensitivity, therefore it was speculated whether insulin mediated glucose uptake was affected in mature adipocytes differentiated with WRW4 and/or AC2-26 and whether it was a contributing factor towards decreased lipid accumulation. Therefore, mRNA expression levels of glucose transporters and glucose uptake were assessed.




Figure 4.22: mRNA expression of receptors involved in glucose uptake in mature adipocytes differentiated with and without FPR2/ALX antagonist; WRW4 and/or agonist: AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and/or 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, mRNA was extracted using Trisure and cDNA was synthesised. 80ng of cDNA was used in SYBR green mediated quantitative PCR. Ct values were obtained and used to calculate normalised gene expression using the Livak 2^{-ΔΔCt} equation (as described in Chapter 3). mRNA expression levels of GLUT1 (SLC2A1) and GLUT4 (SLC2A4) were significantly upregulated in mature adipocyte; $p=0.0059$, $n=4$ and $p=0.0049$, $n=5$ respectively compared to preadipocytes. However, mRNA expression of these receptors was significantly downregulated in treated mature adipocytes. mRNA expression of GLUT1 was significantly downregulated in WRW4 differentiated mature adipocytes ($p<0.0001$, $n=6$), in AC2-26 differentiated mature adipocytes ($p=0.0316$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p=0.0212$, $n=5$), compared to vehicle control. mRNA expression of GLUT4 was significantly downregulated in WRW4 differentiated mature adipocytes ($p<0.0001$, $n=7$), in AC2-26 differentiated mature adipocytes ($p=0.0001$, $n=5$) and in WRW4 + AC2-26 differentiated mature adipocytes ($p=0.267$, $n=5$) compared to vehicle control. Preadipocytes and vehicle control normalised to 1. Data presented as Mean \pm SEM. * vs 1. Dark Grey= control, Grey= treated, Housekeeping gene: 18s RNA
GLUT= Glucose transporter

SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 and/or 10 μ M AC2-26 and mRNA expression levels were studied. mRNA expression of glucose transporters; GLUT1 (*SLC2A1*) and GLUT4 (*SLC2A4*) were significantly upregulated in mature adipocytes (9.57 \pm 2.4-fold, p=0.0059, n=4 and 9.18 \pm 2.1-fold, p=0.0049, n=5, respectively) compared to preadipocytes (1, n=5), as shown in Figure 5.22. However, mRNA expression levels were significantly downregulated in mature adipocytes differentiated with WRW4 and/or AC2-26, as shown in Figure 5.21. mRNA expression of GLUT1 was significantly downregulated in WRW4 differentiated mature




Figure 5.23: Glucose uptake in mature adipose treated with insulin. SGBS cells were differentiated over 14 days to reach a mature phenotype and maintained for further 4 days. The cells were starved of glucose for 40 minutes and then stimulated with insulin for 20 minutes followed by 2-DG for 20 minutes. The uptake was measured using a spectrophotometer at 412nm. 2-DG is taken up and metabolised to 2DG6P, therefore, the uptake of 2-DG is equivalent to 2-DG6P accumulation. One tailed unpaired student's T-test was used to statistically analyse the data. Glucose uptake significantly increased in adipocytes treated with insulin compared to untreated vehicle control (p=0.0032, n=3). Data presented as Mean \pm SEM. P value was set at P<0.05 and is denoted by * vs Glucose. 2-DG= 2-deoxyglucose and 2DG6P= 2-DG-6 phosphate

adipocytes (0.26 ± 0.1 -fold, $p < 0.0001$, $n=6$), in AC2-26 differentiated mature adipocytes (0.52 ± 0.2 -fold, $p=0.0316$, $n=5$), and in WRW4 + AC2-26 differentiated mature adipocytes (0.64 ± 0.1 -fold, $p=0.0212$, $n=5$) compared to vehicle control (1, $n=6$). Similarly, mRNA expression of GLUT4 were significantly downregulated in WRW4 differentiated mature adipocytes (0.34 ± 0.1 -fold, $p < 0.0001$, $n=7$), in AC2-26 differentiated mature adipocytes (0.33 ± 0.1 -fold, $p=0.0001$, $n=5$), and in WRW4 + AC2-26 differentiated mature adipocytes (0.72 ± 0.1 -fold, $p=0.0267$, $n=5$) compared to vehicle control (1, $n=6$).

Acute treatment of WRW4 and/or AC2-26 decreased glucose uptake in mature adipocytes compared to vehicle control, as shown in Figure 5.27. Glucose uptake significantly increased with insulin treatment in mature adipocytes compared to non-insulin stimulated cells (20.0 ± 3.2 pmol vs 3.1 ± 0.4 pmol, $p=0.0032$, $n=3$), as shown in Figure 4.23.

Interestingly, 2-DG uptake was significantly increased in non-insulin stimulated adipocytes treated with WRW4 compared to 2DG6P only adipocytes (10.3 ± 0.1 pmol vs 7.0 ± 0.4 pmol, $p=0.0006$, $n=3$), as shown in Figure 5.24. Furthermore, this was increased in insulin stimulated adipocytes treated with WRW4 compared to non-insulin stimulated WRW4 treated adipocytes (13.8 ± 1.4 pmol vs 10.3 ± 0.1 pmol, $p=0.0328$, $n=3$) and 2DG6P only adipocytes (13.8 ± 1.4 pmol vs 7.0 ± 0.4 , $p=0.0045$, $n=3$).




Figure 5.24: Glucose uptake in mature adipocytes treated with insulin and FPR2ALX antagonist; WRW4
 SGBS cells were differentiated over 14 days to reach a mature phenotype and maintained for further 4 days. The cells were starved of glucose for 40 minutes and then stimulated with insulin and 10 μ M WRW4 for 20 minutes followed by 2-DG for 20 minutes. The uptake was measured using a spectrophotometer at 412nm. 2-DG is taken up and metabolised to 2DG6P, therefore, the uptake of 2-DG is equivalent to 2-DG6P accumulation. One tailed unpaired student's T-test was used to statistically analyse the data. Glucose uptake significantly increased in adipocytes treated with WRW4 compared to 2DG6P only adipocytes ($p=0.0006$, $n=3$) and in adipocytes treated with WRW4 and stimulated with insulin compared to unstimulated and WRW4 treated control ($p=0.0328$, $n=3$ and $p=0.0045$, $n=3$, respectively). Data presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs 2DG6P only and # vs WRW4 treated. 2-DG= 2-deoxyglucose and 2DG6P= 2-DG-6 phosphate

A similar pattern of 2-DG uptake was observed in adipocytes treated with AC2-26, as shown in Figure 5.25. Uptake of 2-DG was slightly but not significantly increased in non-insulin stimulated adipocytes treated with AC2-26 compared to 2DG6P only adipocytes (7.8 ± 1.1 pmol vs 7.0 ± 0.4 pmol, $p=0.2636$, $n=3$). Furthermore, uptake was significantly increased in insulin stimulated AC2-26 treated adipocytes compared to non-insulin

stimulated AC2-26 treated adipocytes (12.1 ± 1.1 pmol vs 7.8 ± 1.1 pmol, $p=0.0226$, $n=3$) and 2DG6P adipocytes (12.1 ± 1.1 pmol vs 7.0 ± 0.4 pmol, $p=0.0051$, $n=3$).




Figure 5.25: Glucose uptake in mature adipocytes treated with insulin and FPR2ALX agonist; AC2-26
 SGBS cells were differentiated over 14 days to reach a mature phenotype and maintained for further 4 days. The cells were starved of glucose for 40 minutes and then stimulated with insulin and $10\mu\text{M}$ AC2-26 for 20 minutes followed by 2-DG for 20 minutes. The uptake was measured using a spectrophotometer at 412nm. One tailed unpaired student's T-test was used to statistically analyse the data. 2-DG is taken up and metabolised to 2DG6P, therefore, the uptake of 2-DG is equivalent to 2-DG6P accumulation. Glucose uptake significantly increased in adipocytes stimulated with insulin and treated with AC2-26 compared to unstimulated AC2-26 only treated control ($p=0.0226$, $n=3$) and 2DG6P glucose only ($p=0.0051$, $n=3$). Data presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs 2GD6P and # vs AC2-26 treated. 2-DG= 2-deoxyglucose and 2DG6P= 2-DG-6 phosphate

Moreover, a similar pattern of 2-DG uptake was observed in adipocytes treated with WRW4 + AC2-26, as shown in Figure 5.26. Uptake of 2-DG was significantly increased in non-insulin stimulated adipocytes treated with WRW4 + AC2-26 compared to 2GD6P only adipocytes (10.6 ± 0.4 pmol vs 7.0 ± 0.4 pmol, $p=0.003$, $n=2/3$). Furthermore, this was

increased in insulin stimulated adipocytes treated with WRW4 + AC2-26 compared to non-insulin stimulated WRW4 + AC2-26 adipocytes (16.7 ± 1.5 pmol vs 10.9 ± 0.4 pmol, $p=0.0293$, $n=2/3$) and 2DG6P only adipocytes (16.7 ± 1.5 pmol vs 7.0 ± 0.4 pmol, $p=0.0017$, $n=3$).




Figure 5.26: Glucose uptake in mature adipocytes treated with insulin and FPR2ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated over 14 days to reach a mature phenotype and maintained for further 4 days. The cells were starved of glucose for 40 minutes and then stimulated with insulin, $10\mu\text{M}$ WRW4 and $10\mu\text{M}$ AC2-26 for 20 minutes followed by 2-DG for 20 minutes. The uptake was measured using a spectrophotometer at 412nm. One tailed unpaired student's T-test was used to statistically analyse the data. 2-DG is taken up and metabolised to 2DG6P, therefore, the uptake of 2-DG is equivalent to 2-DG6P accumulation. Glucose uptake significantly increased in adipocytes treated with WRW4 + AC2-26 compared to 2DG6P only ($p=0.003$, $n=2/3$) and in adipocytes stimulated with insulin and treated with WRW4 + AC2-26 compared to WRW4 + AC2-26 only treated and 2DG6P only adipocytes ($p=0.0293$, $n=2/3$ and $p=0.0017$, $n=3$, respectively). Data presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs untreated and # vs AC2-26 treated. 2-DG= 2-deoxyglucose and 2DG6P= 2-DG-6 phosphate

However, 2-DG uptake was significantly less in insulin stimulated treated adipocytes (WRW; $p=0.0222$, $n=3$, AC2-26; $p=0.0126$, $n=3$ and WRW4 + AC2-26; $p=0.0567$, $n=3$)

compared to insulin stimulated vehicle control adipocytes, as shown in Figure 5.27. Additionally, no significant difference between insulin stimulated treated adipocytes was observed.




Figure 5.27: Comparison of glucose uptake in adipocytes treated with FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. One tailed unpaired student's T-test was used to statistically analyse the data. 2-DG was significantly less in treated cells (WRW; $p=0.0222$, $n=3$, AC2-26; $p=0.0126$, $n=3$ and WRW4 + AC2-26; $p=0.0567$, $n=3$) compared to adipocytes treated with insulin only and no significant difference between treated adipocytes was observed. Data presented as Mean \pm SEM. P value was set at $P<0.05$ and is denoted by * vs 2-DG and insulin only

5.5.8 mRNA expression of adipokines in mature adipocytes differentiated with and without FPR2/ALX antagonist and/or agonist

Lipid accumulation and expression and secretion of adipokines are positively correlated with adipocyte differentiation. It was questioned whether decreased lipid accumulation was associated with an immature adipocyte phenotype, therefore gene expression of adipokines was investigated. SGBS cells were seeded and induced to differentiate for 14 days with 10 μ M WRW4 and/or 10 μ M AC2-26 and mRNA expression levels were studied.




Figure 5.28: mRNA expression of adipokines in mature adipocytes differentiated with FPR2/ALX antagonist; WRW4 and/or agonist; AC2-26. SGBS cells were differentiated with 10 μ M WRW4 and/or 10 μ M AC2-26 over 14 days to reach a mature adipocyte phenotype. On day 14, mRNA was extracted using Trisure and cDNA was synthesised. 80ng of cDNA was used in SYBR green mediated quantitative PCR. Ct values were obtained and used to calculate normalised gene expression using the Livak 2^{-ΔΔCT} equation (as described in Chapter 3). One tailed unpaired student's T-test was used to statistically analyse the data. mRNA expression levels of Leptin (p=.0030, n=5), Adiponectin (p=0.0323, n=4) and Annexin A1 (p=0.0438, n=3) were significantly upregulated in mature adipocytes compared to preadipocytes. Similarly, Resistin (p=0.841, n=4) and Visfatin (p=0.2151, n=3) were upregulated in mature adipocytes compared to preadipocytes, however were not statistically significance. mRNA expression levels of these adipokines were significantly downregulated in WRW4 differentiated mature adipocytes; Leptin (p=0.0019, n=4), Resistin (p<0.0001, n=9), Visfatin (p=0.0072, n=5), Adiponectin (p=0.0074, n=4) and Annexin A1 (p=0.0023, n=8) compared to vehicle control. Similarly, mRNA expression levels of these adipokines were significantly downregulated in AC2-26 differentiated mature adipocytes Leptin (p=0.0092, n=5), Resistin (p<0.0001, n=3), Visfatin (p=0.0064, n=5), Adiponectin (p=0.0069, n=4) and Annexin A1 (p=0.0002, n=6) compared to vehicle control. Likewise, mRNA expression levels of these adipokines were significantly downregulated in WRW4 + AC2-26 differentiated mature adipocytes; Leptin (p=0.0379, n=3), Resistin (p=0.0045, n=3) and Annexin A1 (p=0.0484, n=3) compared to vehicle control. mRNA expression levels of Adiponectin (p=0.0770, n=3) were downregulated and Visfatin were upregulated (p=0.2064, n=3) in these cells compared to vehicle control, however, the differences were not statistically significant. Preadipocytes and DMSO control normalised to 1. Data presented as Mean±SEM. P value was set at P<0.05 and is denoted by * vs 1. Dark Gray=

mRNA expression levels of adipokines were upregulated in mature adipocytes compared to preadipocytes, as shown on Figure 5.28. mRNA expression levels of Leptin (*LEP*) (26.0 ± 5.7 -fold, $p=0.0030$, $n=5$), Adiponectin (*ADIPOQ*) (15.4 ± 6.4 -fold, $p=0.0323$, $n=4$) and Annexin A1 (*ANXA1*) (1.5 ± 0.3 -fold, $p=0.0438$, $n=3$) were significantly upregulated in mature adipocytes compared to preadipocytes (1 , $n=4$). Similarly, Resistin (*RETN*) (1.8 ± 0.5 -fold; $p=0.841$, $n=4$) and Visfatin (*NAMPT*) (1.2 ± 0.3 -fold, $p=0.2151$, $n=3$) were upregulated in mature adipocytes compared to preadipocytes, however were not statistically significant.

mRNA expression levels of these adipokines were downregulated in WRW4 differentiated mature adipocytes, as shown in Figure 5.28. mRNA expression levels of Leptin (0.71 ± 0.1 -fold, $p=0.0019$, $n=4$), Resistin (0.25 ± 0.1 -fold, $p<0.0001$, $n=9$), Visfatin (0.50 ± 0.1 -fold, $p=0.0072$, $n=5$), Adiponectin (0.72 ± 0.1 -fold, $p=0.0074$, $n=4$) and Annexin A1 (0.62 ± 0.1 -fold, $p=0.0023$, $n=8$) were significantly downregulated compared to vehicle control (1 , $n=9$).

Similarly, mRNA expression levels of these adipokines were downregulated in AC2-26 differentiated mature adipocytes, as shown in Figure 5.28. mRNA expression levels of Leptin (0.46 ± 0.2 -fold, $p=0.0092$, $n=5$), Resistin (0.12 ± 0.1 -fold, $p<0.0001$, $n=3$), Visfatin (0.50 ± 0.2 -fold, $p=0.0064$, $n=5$), Adiponectin (0.41 ± 0.2 -fold, $p=0.0069$, $n=4$) and Annexin A1 (0.35 ± 0.1 -fold, $p=0.0002$, $n=6$) were significantly downregulated compared to vehicle control (1 , $n=6$).

Likewise, mRNA expression levels of these adipokines were downregulated in WRW4 + AC2-26 differentiated mature adipocytes, as shown in Figure 5.28. mRNA expression levels of Leptin (0.61 ± 0.2 -fold, $p=0.0379$, $n=3$), Resistin (0.64 ± 0.1 -fold, $p=0.0045$, $n=3$)

and Annexin A1 (0.92 ± 0.05 -fold, $p=0.0484$, $n=3$) were significantly downregulated compared to vehicle control (1, $n=3$). mRNA expression levels of Adiponectin (0.88 ± 0.1 -fold, $p=0.0770$, $n=3$) were downregulated and Visfatin were upregulated (1.28 ± 0.4 -fold, $p=0.2064$, $n=3$) in these cells compared to vehicle control (1, $n=3$), however, the differences were not statistically significant.

5.6 Discussion

In the current study, the role of ANXA1 in adipocyte differentiation was investigated. The two main *in vivo* studies in murine models published in the area of ANXA1 and adipogenesis reported opposing results; (Warne et al. 2006) conclude ANXA1 may be required for early stages of adipogenesis, namely differentiation from MSC to preadipocytes, whereas (Akasheh et al. 2013) conclude ANXA1 may be involved in regulation of lipolysis in mature adipocytes. In the current study, inhibition of ANXA1 activity did not affect the early stages of adipogenesis in a human primary cell, however, mature adipocytes differentiated with ANXA1 receptor agonist and antagonist accumulated less lipid suggesting, ANXA1 may be involved in modulating the rate of adipogenesis and maintaining a mature adipocyte phenotype. Furthermore, the role of FPR receptors, FPR1 and FPR2/ALX in adipogenesis were investigated as recent studies report. FPR1 agonist fMLF has shown to promote differentiation of hMSC into osteoblasts and suppress their differentiation into preadipocytes (Peacey, Wright et al. 2012, Shin, Jang et al. 2011). This observation suggests FPR's expression is upregulated in preadipocytes compared to their precursor cells and continues to upregulate in mature adipocytes as reported by (Kosicka et al. 2013). In this study, differential activation of these receptors resulted in different effect on lipid accumulation in mature adipocytes.

5.6.1 Lipid accumulation

In the current study, differentiating adipocytes with a specific ANXA1 antagonist; WRW4, reduced total lipid accumulation in mature adipocytes, which reflects previous reports describing the reduced fat mass in *ANXA1*^{-/-} mice as reported by (Warne et al. 2006). Interestingly, specifically inhibiting ANXA1 activity or exogenous treatment with

AC2-26 resulted in decreased lipid accumulation in mature adipocytes, but the morphology of the lipid droplets differed. Although, on average, the size of the lipid droplets in mature adipocytes differentiated with specific ANXA1 inhibition was unchanged, morphological analysis revealed increased number of smaller lipid droplets; a phenomenon previously reported in lung fibroblasts, where *ANXA1*^{-/-} mice contained lipid droplets but, *ANXA1*^{+/+} mice did not (Croxtall, Gilroy et al. 2003). Similarly, exogenous treatment with AC2-26 did not result in increased number of smaller lipid droplets. However, on average the size of the lipid droplets was significantly smaller suggesting, ANXA1 may play a role in regulating the size of the lipid droplets and in turn regulating the overall lipid content of mature adipocytes mediating a positive effect as the size and the number of lipid droplets positively correlates with obesity (Engin 2017).

5.6.2 Intracellular Pathways

Interestingly specific inhibition of ANXA1 activity or exogenous treatment with AC2-26 display similar patterns of lipid accumulation in mature adipocytes. Therefore, the roles of ANXA1 receptors in adipogenesis using receptor agonists and antagonists were explored. Previous studies have investigated the role of ANXA1 in adiposity through knockout mice models, however, little evidence is present to suggest ANXA1 or its receptors are involved in adipogenesis. Full length ANXA1 displays specific affinity towards binding and activating the FPR2/ALX receptor, whereas, cleaved peptides such as AC2-26 have shown to bind and activate all members of the FPR family (Dalli, Montero-Melendez et al. 2012). It is interesting to note that, the cleaved peptides of ANXA1 are believed to terminate the anti-inflammatory properties of this protein; binding of ANXA1 to its receptor FPR2/ALX inhibits neutrophil migration, whereas, activation of FPR1 or FPR2/ALX by AC2-26 stimulates neutrophil migration during

inflammation (Dalli, Montero-Melendez et al. 2012, Gavins, Hickey 2012, Hazeldine, Hampson et al. 2015). However, activation of FPR1 and FPR2/ALX have been shown to activate different intracellular signalling cascades, which could be a contributing factor in the differential responses elicited by these receptors.

In the current study, it was hypothesised that blocking the FPR2/ALX receptor may force endogenous AC2-26 to bind to the FPR1 receptor and differentially activate downstream signalling pathways (Figure 5.29a). Adipocytes were differentiated with WRW4 + AC2-26 showed decreased lipid accumulation and an increased number of smaller lipid droplets in mature adipocytes, like that observed in mature adipocytes differentiated with WRW4 only, supporting the hypothesis. Overexpressing AC2-26 may have led to FPR1




Figure 5.29: Activation of FPR receptors. a) Specifically blocking FPR2/ALX receptor using WRW4 may force endogenous AC2-26 (green) to bind to FPR1 receptor. b) Activation of both FPR2/ALX and FPR1 receptors using exogenous AC2-26 (yellow) ((Woloszynek, Hu et al. 2012)). c) Specifically blocking FPR2/ALX receptor using WRW4 and overexpression of exogenous AC2-26 to test the hypothesis. FPR= Formyl peptide receptor, FPR1= Formyl peptide receptor 1 and FPR2/ALX= Formyl peptide receptor 2

activation along with activation of FPR2/ALX, thus displaying similar effects on lipid accumulation as specifically inhibiting FPR2/ALX receptor alone (Figure 5.29b). Furthermore, blocking FPR1 receptor using lower concentrations of BOC-2 resulted in opposing effects compared to specifically blocking FPR2/ALX receptor suggesting, FPR1 and FPR2/ALX both may act together to regulate and achieve lipid homeostasis in mature adipocytes through heterodimerization or induce crosstalk between intracellular pathways (Kosicka et al. 2013, Dalli et al. 2012). Interestingly, complete inhibition of ANXA1/AC2-26 activity using higher concentrations of BOC-2 showed no difference suggesting, ANXA1 may be involved in regulating adipogenesis in conditions where its dysregulated such as in obesity, (similar to the *ANXA1*^{-/-} mice) not displaying any phenotypic differences unless challenged with an inflammatory stimulus (Akasheh et al. 2013, Warne et al. 2006, Damazo, Yona et al. 2006, Y. H. Yang, Morand et al. 2004, Babbitt, Laukoetter et al. 2008).

FPR1 and FPR2/ALX mediate their effects through MAPK signalling. Which specific intracellular MAPK cascades are activated by the individual FPR's is not clear or conclusive further reinforcing the complexity of these signalling pathways. FPR2/ALX activation by ANXA1/AC2-26 is reported to induce ERK1/2 MAPK phosphorylation but not p38 MAPK or JUN MAPK pathways (Bena, Brancalione et al. 2012), (Dalli et al. 2012, Gavins, Hickey 2012), whereas, others report FPR2/ALX homodimerization by ANXA1/AC2-26 activates the p38 MAPK pathway (Filep 2013) and AC2-26 induced heterodimerisation of FPR1 and FPR2/ALX activates JUN MAPK pathway (Cooray, Gobbetti et al. 2013). Furthermore, ERK1/2 and p38 MAPK predominantly influence chemotaxis of neutrophils, a response exhibited by AC2-26 induced activation of FPR1 (Rabiet, Huet et al. 2007). The contrasting findings of MAPK activation may be cell type

specific and therefore, in this project ERK1/2 MAPK was attributed to FPR2/ALX and p38 MAPK was attributed to FPR1 activation (Figure 5.30) because specific inhibition of FPR2/ALX or exogenous treatment with AC2-26 or both (Figure 5.29) decreased p38 MAPK activity in mature adipocytes. Previously, *ANXA1*^{-/-} mice showed increased p38 MAPK signalling, whereas, treatment with ANXA1 inhibited p38 MAPK expression, further supporting the association of FPR1 and p38 MAPK activity (Y. H. Yang, Toh et al. 2006).




Figure 5.30: Intracellular signalling cascades associated with FPR receptors. FPR2/ALX is thought to modulate the activity of ERK1/2 MAPK whereas FPR1 is thought to modulate the activity of p38MAPK. ERK1/2 MAPK= FPR1= Formyl peptide receptor 1, FPR2/ALX= Formyl peptide receptor 2, Extracellular signal-regulated kinase mitogen activated protein kinase 1/2, PPAR γ Peroxisome proliferator activated receptor γ and C/EBP= CCAAT-enhancer binding proteins

5.6.2.1 p38 MAPK and adipogenesis

p38 MAPK activity is involved in adipogenesis, however, whether it positively or negatively regulates this process is yet to be decided (Figure 5.31). Phosphorylation of p38 MAPK was decreased in all 3 experimental conditions in the current study, thus reducing its activity.




Figure 5.31: Role of p38 MAPK in adipogenesis. Activated p38 MAPK increases PPAR γ and C/EBP activity thus increasing adipogenesis, however, activated p38 MAPK phosphorylates NFATC4; an activator of PPAR γ expression in the nucleus. Phosphorylation of NFATC4 inhibits its translocation into the nucleus thus inhibiting adipogenesis.

GPCR= G-protein coupled receptor, p38 MAPK= p38 mitogen activated protein kinase, PPAR γ Peroxisome proliferator activated receptor γ , C/EBP= CCAAT-enhancer binding proteins and NFATC4= Nuclear Factor of activated T cells 4

The first study demonstrating a positive role of p38 MAPK in adipogenesis used 3T3-L1 preadipocytes and reported a decrease in adipocyte formation in the presence of p38 MAPK inhibitors (early on) and associate this effect with decreased C/EBP β phosphorylation (Engelman, Lisanti et al. 1998). Acute overexpression of mitogen-activated protein kinase 6; an upstream activator of p38 MAPK stimulates adipogenesis, whereas prolonged activation result in adipocyte cell death (Engelman et al. 1998,

Engelman, Berg et al. 1999). Similarly, other authors report decrease in PPAR γ activity in the presence of p38 MAPK inhibitors resulting in decreased adipogenesis (Hata, Nishimura et al. 2003, Takenouchi, Takayama et al. 2004), suggesting, p38 MAPK acts as an activator of PPAR γ (Hata et al. 2003). However, in human preadipocytes treated with p38 MAPK inhibitors, a smaller but significant decrease in adipogenesis is reported compared to the murine cell line. AC2-26 may act as a p38 MAPK inhibitor as human preadipocytes differentiated with or without p38 MAPK inhibitor, result in decreased lipid accumulation, lipogenesis and leptin secretion in mature adipocytes; similar to the phenotypes observed in the current study (Figure 5.31) (Patel, Holder et al. 2003, Xu, Ju et al. 2014, Aouadi, Laurent et al. 2006, Aouadi, Jager et al. 2007). Furthermore, (Xu et al. 2014) report that not only does p38 MAPK positively regulate gene expression of PPAR γ , it also regulates the expression of genes involved in the formation of lipid droplets such as C/EBP, thus decreasing p38 MAPK phosphorylation could be a contributing factor in the decreased number of large lipid droplets observed.

Other authors report a negative role of p38 MAPK in adipogenesis. Phosphorylation of Nuclear Factor of Activated T cells 4, an activator of PPAR γ expression in the nucleus, by p38 MAPK inhibits its translocation into the nucleus thus inhibiting adipogenesis (T. Yang, Xiong et al. 2002, Ho, Kim et al. 1998). Others report, impaired adipogenesis in 3T3-L1 cells without affecting p38 MAPK activity (S. Liu, Xu et al. 2014). These opposing reports regarding the role of p38 MAPK in adipogenesis suggest, p38 MAPK may have differential roles in the process of adipogenesis depending on the stage of

adipogenesis, differentiation condition and/ or may vary in different cell models (Xu et al. 2014).




Figure 5.32: Activation of FPR1 decreases p38 MAPK phosphorylation. Phosphorylated p38 MAPK upregulates the expression of adipogenic factors such as PPAR γ and C/EBP thus positively regulating adipogenesis. However, activation of FPR1 decreases p38 MAPK phosphorylation therefore decrease the expression of PPAR γ and C/EBP thus adipogenesis.
FPR1= Formyl peptide receptor 1, p38 MAPK= p38 kinase mitogen activated protein kinase, PPAR γ Peroxisome proliferator activated receptor γ and C/EBP= CCAAT-enhancer binding proteins

Levels of ANXA1 increase in obesity and p38 MAPK activity decreases in both obese human patients and obese animals (Aouadi et al. 2006) (He, Zhu et al. 2013). The results of the current study are the first to link ANXA1 and p38 MAPK activity in adipocyte and may indicate a link between increase in ANXA1 activity, decrease in p38 MAPK activity and dysregulation in adipocyte size in obesity.

5.6.2.2 ERK1/2 MAPK and adipogenesis

Reports investigating the role of ERK1/2 MAPK in adipogenesis also present contradictory results due to ERK1/2 MAPK having multiple roles and each isoform

having distinct biological functions, all contributing to its complex nature (Figure 5.33) (Bost, Aouadi, Caron, and Binetruy 2005). Earlier reports conclude a positive role of ERK1/2 MAPK in adipogenesis (Benito, Porras et al. 1991), whilst later studies demonstrate PPAR γ as a substrate of ERK1/2 MAPK and phosphorylation by active ERK1/2 MAPK decreases its transcriptional activity and therefore, inhibits adipocyte differentiation (Camp, Tafuri 1997, Hu, Kim et al. 1996, Ristow, Muller-Wieland et al. 1998). A hypothesis concluded from these results states that function of ERK1/2 MAPK in adipogenesis is regulated in a timely manner; ERK1/2 MAPK is activate for the proliferative step (PPAR γ activity not required), whilst, it has to be deactivated in the later stages to avoid phosphorylation of PPAR γ (required for terminal differentiation) (Bost et al. 2005). Mice lacking ERK1 MAPK have decreased adiposity and less adipocytes and when challenged with high fat diet (HFD) are resistant to obesity and are protected from insulin resistance (Bost, Aouadi, Caron, Even et al. 2005). Studies in 3T3-L1 and other cell models show that activity of ERK1/2 MAPK is necessary for expression of crucial adipogenic regulators; C/EBP α , β , γ and PPAR γ (Prusty, Park et al. 2002, Aubert, Dessolin et al. 1999, Belmonte, Phillips et al. 2001). However, others report upregulation of these adipogeneic factors in 3T3-L1 adipocytes by attenuating ERK1/2 activity thus promoting adipogenesis (C. Liu, Huang et al. 2017).




Figure 5.33: Role of ERK1/2 MAPK in adipogenesis. ERK1/2 MAPK activation induces upregulation of adipogenic factors such as PPAR γ and C/EBP thus increasing adipogenesis, however, ERK1/2 MAPK activation phosphorylates PPAR γ , inhibiting its translocation into the nucleus thus decreasing adipogenesis. GPCR= G-protein coupled receptor, ERK1/2 MAPK= Extracellular signal-regulated kinase mitogen activated protein kinase 1/2, PPAR γ Peroxisome proliferator activated receptor γ and C/EBP= CCAAT-enhancer binding proteins




Figure 5.34: Activation of FPR2/ALX may activate ERK1/2 MAPK in mature adipocytes. Activation of FPR2/ALX by ANXA1/AC2-26 may activate ERK1/2 MAPK and increase expression of lipogenic proteins such as PPAR γ and C/EBP therefore, promoting adipogenesis. FPR2/ALX= Formyl peptide receptor 2, ANXA= Annexin A1, ERK1/2 MAPK= Extracellular signal-regulated kinase mitogen activated protein kinase 1/2, PPAR γ Peroxisome proliferator activated receptor γ and C/EBP= CCAAT-enhancer binding proteins

ERK1/2 MAPK phosphorylation was unaffected in mature adipocytes differentiated with WRW4 and/or AC2-26 compared to control adipocytes. This suggests there may be crosstalk between p38 MAPK and ERK1/2 MAPK, as presented in Figure 5.35b and may explain increased adipogenesis when partially blocking and no difference when completely blocking ANXA1/AC2-26 activity in adipogenesis (Xu et al. 2014). This suggests ERK1/2 MAPK may not be required for adipogenesis in the later stages as reported but may play a positive role if solely activated (Ning, He et al. 2016, Bost et al. 2005). To further validate the hypothesis, the mechanism by which MAPK regulates adipogenesis were investigated.




Figure 5.35: Cross talk between p38 MAPK and ERK1/2 MAPK may regulate adipogenesis (hypothesis based on results). a) Activation of FPR1 decreased p38 MAPK phosphorylation, activation of FPR2/ALX may induce ERK1/2 MAPK activation, however, downregulation of p38 MAPK may block the activation of ERK1/2 MAPK thus decreasing adipogenesis. b) Blocking FPR1 may force endogenous ANXA1/AC2-26 to bind to FPR2/ALX and activate ERK1/2 MAPK thus increasing adipogenesis. c) Blocking FPR1 may increase p38 MAPK activation (adipogenesis) but blocking FPR2/ALX decreases ERK1/2 MAPK phosphorylation (no adipogenesis), therefore, no difference in adipogenesis is observed.

FPR1= Formyl peptide receptor 1, FPR2/ALX= Formyl peptide receptor 2, ERK1/2 MAPK= Extracellular signal-regulated kinase mitogen activated protein kinase 1/2 and p38 MAPK= p38 mitogen activated protein kinase

5.6.3 Lipid Metabolism in Mature Adipocytes

MAPKs are thought to regulate adipogenesis through peroxisome proliferator activated receptors (PPAR's). PPAR's play important roles in adipogenesis, inflammation, atherogenesis, glucose homeostasis and cancer (Obregon 2014). PPAR's are nuclear receptors that act as transcription factors regulating the expression of proteins involved in lipid metabolism upon stimulation by thiazolidinediones and glitazones (class of insulin-sensitizing drugs). PPAR γ is the key regulator of adipose development and adipocyte insulin sensitivity (Kramer, Al-Khalili et al. 2007). Once PPAR γ expression is induced, it activates majority of the genes required for this process; aP2 (required for transport of fatty acids), perilipin (covering the surface of the lipid droplet), *de novo* lipid metabolism (including; acetyl-CoA acetyltransferase; ACC, fatty acid synthase; FAS and glycerol-3-phosphate acyltransferase; GPAT) and those involved in glucose homeostasis.

Complete inhibition of ANXA1 signalling did not affect lipid accumulation and previous data report mRNA expression of PPAR γ , ACC and FAS in *ANXA1*^{-/-} mice are not differently regulated compared to wild type mice, unless fed a HFD (Akasheh et al. 2013). In accordance to previous reports, decreased phosphorylation of p38 MAPK in mature adipocytes differentiated with WRW4 and/or AC2-26, resulted in decreased mRNA expression of PPAR γ and SREBP (Hata et al. 2003) and subsequently, may have downregulated the mRNA expression of enzymes; ACC, FASN and GPAT in the current study. Furthermore, the data in the current study indicated that mature human adipocytes differentiated with WRW4 and/or AC2-26 resulted in decreased total %ACC protein, suggesting that binding of AC2-26 to FPR1 and its activation, downregulates lipogenesis.

Previous *ANXA1*^{-/-} studies conclude *ANXA1* may play a role in regulating lipolysis (Warne et al. 2006, Akasheh et al. 2013). In line with this assertion, mature adipocytes differentiated with WRW4 and/or AC2-26 showed decreased mRNA expression of enzymes involved in lipolysis; ATGL, HSL and MGL in the current study. Furthermore, the rate of lipid peroxidation was decreased in mature adipocytes differentiated with WRW4 and/or AC2-26 (Sakamoto, Repasky et al. 1996).

5.6.4 Glucose uptake in Mature Adipocytes

Adipocytes regulate lipid and glucose homeostasis through regulating glucose uptake via adipose specific glucose transporter; GLUT4 and nonspecific glucose transporter; GLUT1. The earliest event in the onset of insulin resistance is the downregulation of GLUT4; an effect which is observed in obesity and T2DM. Clinical studies conducted in overweight diabetic patients report decrease in abundance of *ANXA1* protein in the visceral adipose tissue when compared to pre-obese nondiabetic patients (Murri, Insenser et al. 2013). However, it is reported, *ANXA1* gene is unlikely to contain variants that contribute to the risk of developing T2DM (Lindgren, Nilsson et al. 2001). Thus, the altered expression may be due to T2DM rather than causing the development of T2DM. It is proposed that *ANXA1* could be a potential physiological regulator of insulin action in target tissues as it promotes a dose dependent inhibition of insulin receptor autophosphorylation and stimulation of insulin release in MIN6N8a; an insulin secreting cell line (Lindgren et al. 2001, Won, Kang et al. 2003).

The role of *ANXA1* in insulin sensitivity is not clear. A PPAR γ agonist, bearing anti-inflammatory and insulin sensitizing properties; Rosiglitazone upregulates *ANXA1* expression in the adipose tissue whereas glucose load or exposure to TNF α upregulates

ANXA1 expression in leukocytes, inducing insulin resistance (Ahmed, Neville et al. 2010, Fujimoto, Mochizuki et al. 2010). PPAR γ stimulation improves glucose tolerance and insulin sensitivity in T2DM patients in animal models, however, the mechanisms remain unknown (Picard, Auwerx 2002). The downregulation of GLUT4 and GLUT1 mRNA expression in mature adipocytes differentiated with WRW4 or AC2-26 in the current study may be due to downregulation of PPAR γ activity in these cells. Although, activation of PPAR γ by its agonists upregulate glucose uptake, it does not increase cellular expression of GLUT4, but does increase GLUT1 (Nugent, Prins et al. 2001). Acute treatment with WRW4 and/or AC2-26 increased insulin independent glucose uptake, possibly through GLUT1, however, insulin mediated glucose uptake was reduced compared to control. The data suggests that, ANXA1 may decrease insulin mediated glucose uptake but increase insulin independent glucose uptake in mature adipocytes. However, the data in the current study demonstrates that in addition to GLUT4, GLUT1 mRNA expression was also downregulated in mature adipocytes differentiated with WRW4 and/or AC2-26. Expression of GLUT1 significantly upregulates in obesity and positively correlates with obesity-associated inflammatory cytokines such as IL6 and TNF α (Yin, Gao et al. 2009, Docanto, Ham et al. 2015).

Adipocyte size positively correlates with insulin resistance and the opposite effect has been reported previously. It is well known that inhibition of p38 MAPK improves glucose transport in L6 and 3T3-L1 cells (Furtado, Somwar et al. 2002, Michelle Furtado, Poon et al. 2003), rather than decreased glucose uptake as observed the current study. Furthermore, p38 MAPK activity was previously shown to mediate downregulation of GLUT4 and is higher in isolated adipocytes from T2DM patients (Carlson, Koterski et al. 2003), whereas, in the current study p38 MAPK activity and glucose uptake were

decreased, therefore it is thought that downregulation of GLUT4 may be due to downregulation of PPAR γ , as opposed to a direct effect of p38 MAPK.

5.6.5 Adipokines in Mature adipocytes

In addition to its role as a fat storage organ, the adipose tissue serves as an endocrine organ by synthesising and secreting soluble mediators (adipokines) that mediate interaction between the tissue, inflammation and immunity (Tilg, Moschen 2006). Furthermore, adipokines are critical regulators of lipid and glucose homeostasis and mediate crosstalk between the adipose tissue and several other metabolic organs such as, liver, muscle and the pancreas (Cao 2014).

As their synthesis and secretion increases in mature adipocytes compared to preadipocytes, their mRNA expression levels were investigated to determine the maturity of the cells differentiated with WRW4 and/or AC2-26. Leptin regulates feeding behaviours and lipid accumulation, Resistin is implicated in glucose homeostasis and is thought to have a proinflammatory role in humans and similarly, Visfatin plays a role in insulin sensitivity and secretion (Makki, Froguel et al. 2013). All three adipokines positively correlate with adipogenesis and are increased in obesity and T2DM patients (Makki et al. 2013). In the current study, the mRNA expression of Leptin, Resistin and Visfatin were downregulated in mature adipocytes differentiated with WRW4 and/or AC2-26. Furthermore, mRNA expression of Adiponectin; an adipokine decreased in obesity improves insulin resistance and glucose metabolism as it bears insulin sensitizing effects, and ANXA1 were also downregulated in mature adipocytes differentiated with WRW4 and/or AC2-26 (Makki et al. 2013). The decrease in these adipokines may have

affected insulin sensitivity and glucose uptake in these cells. These results indicate an immature adipocyte phenotype compared to control.

5.7 Conclusion

In conclusion, ANXA1 may not be involved in regulating adipogenesis in normal physiological environments, however, when its expression and signalling are dysregulated in conditions such as obesity, it may slow down the processes of adipogenesis to decrease the expression and secretion of proinflammatory adipokines and cytokines associated with a hypoxic environment, thus improving the inflammatory status. However, this effect may be overshadowed by the continued increase in systematic inflammation associated with rapid adipose tissue expansion. In addition to similar experimental limitations as Study 1 (Chapter 4), technical approach used for imaging ORO staining may have produced experimental bias, therefore, it would be of interest to employ an alternative technique or seek technical assistance.

Chapter 6: Plasma inflammatory biomarkers, insulin sensitivity and gut hormones in response to rapid weight loss through Bariatric Surgery

Plasma Annexin A1 inversely correlate with increasing adiposity and with pro-inflammatory cytokines. However, the effect of weight loss on plasma Annexin A1 levels is unknown. Bariatric surgery is associated with reversal of plasma biomarkers including, adipokines, cytokines and gut hormones. Therefore, the objective of this chapter was to investigate the plasma Annexin A1 levels and correlate it with anthropometric measures and plasma inflammatory biomarkers in bariatric surgery patients. Furthermore, the results of Chapter 4 showed AC2-26 upregulated the gene expression of gut hormone; Dipeptidyl Peptidase 4, therefore, plasma gut hormones were correlated with plasma Annexin A1 in the cohort.

6.1 Abstract

Introduction: Bariatric surgery improves metabolic abnormalities independent of weight loss. Changes to the anatomy of the gut induces physiological changes resulting in altered plasma levels of lipids, adipokines, cytokines and gut hormones. The altered circulatory profile leads to improved local and systemic inflammation, however, the exact mechanisms underlying these changes are unclear. Plasma Annexin A1 (ANXA1) levels are reduced in obesity and inversely correlate with markers of adiposity, however, the effect of weight loss plasma ANXA1 levels is unknown.

Aim: To investigate the plasma ANXA1 levels and correlate with plasma lipids, inflammatory biomarkers, markers of insulin sensitivity and gut hormone in response to rapid weight loss induced by bariatric surgery.

Methodology: Patients due to undergo bariatric surgery who matched the inclusion/exclusion criteria were identified and 27 were recruited into the study. Anthropometric measures and blood samples were taken pre-operation (pre-opt), 6 weeks and 3 months post-operation (post-opt) from patients who consented to participate. The participant cohort was further subcategorised based on type of surgery (SG vs RYGB) or type 2 diabetes mellitus (T2DM) status (non-T2DM vs T2DM). Plasma samples were analysed, and results were statistically analysed using GraphPad prism 8.0. Statistical significance at 95% was determined using One-tailed Pearson's correlation coefficient One-way and Two-way ANOVA's followed by Tukey's Post-hoc test.

Results: The average body mass index (BMI) significantly reduced at 6 weeks and 3 months post-opt compared to pre-opt of the complete, SG, RYGB, non-T2DM and T2DM cohorts. Plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt and correlated with BMI, plasma high density lipoprotein (HDL), tumour necrosis

factor α (TNF α) and adiponectin in the complete cohort. Plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt in the SG and RYGB cohorts and no significant difference was observed between the cohorts at any time point. In the SG cohort, plasma ANXA1 correlated with BMI, plasma adiponectin and PYY levels. In the RYGB cohort, plasma ANXA1 correlated with plasma triglycerides, HDL, C-reactive protein (CRP), interleukin 1 β (IL-1 β), TNF α and glucagon-like peptide 1 (GLP-1). Plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt in the non-T2DM and T2DM cohorts and no significant difference was observed between the cohorts at any time point. In the non-T2DM cohort, plasma ANXA1 correlated with plasma HDL, LDL and homeostatic model assessment of pancreatic β cell function. In the T2DM cohort, plasma ANXA1 correlated with BMI, plasma TNF α and homeostatic model assessment of insulin sensitivity.

Conclusion: Plasma ANXA1 indicate increasing trends with decreasing BMI, however, a firm conclusion regarding its associations with plasma biomarkers is difficult to form due to low number of participants.

6.2 Introduction

Bariatric surgery is considered as the most effective long-term treatment option for morbid obese individuals as it has extensive and sustained effects on weight and significantly reduces the risk of developing obesity associated co-morbidities (Colquitt, Picot et al. 2009). It is beneficial for obese individuals with body mass index (BMI) $\geq 40\text{kg/m}^2$ or $\geq 35\text{kg/m}^2$ who have at least 1 co-morbidity present and if non-surgical treatments have failed to achieve or sustain substantial weight loss as outlined by the guidelines published by the National Institute for Health and Clinical Excellence. The primary aim of bariatric surgery is to reduce the patient's total volume of ingestion and/or absorption of contents of food intake, therefore, the procedures are grouped into 3 broad categories, 1) pure gastric restriction, 2) gastric restriction with some malabsorption and 3) gastric restriction with significant intestinal malabsorption (Jaunoo, Southall 2010).




Figure 6.1: Bariatric surgery procedures. The two common forms of bariatric surgery procedures include, Roux-en-Y Gastric Bypass (RYGB) and Vertical Sleeve Gastrectomy (SG). RYGB combines gastric restriction and malabsorption and involves the division of the stomach into two pouches. The smaller of the pouch is connected to the proximal jejunum bypassing the duodenum. SG is based on gastric restriction and involves the vertical division of the stomach.

(Frikke-Schmidt, O'Rourke et al. 2016)

The most common procedures include, gastric banding, sleeve gastrectomy (SG), Roux-en-Y gastric bypass (RYGB) and biliopancreatic diversion.

SG is based on gastric restriction and is a less invasive and a less technically challenging procedure compared to RYGB (Frikke-Schmidt, O'Rourke et al. 2016). It is relatively quick to carry out with reduced risk of complications, therefore, was initially offered to high risk bariatric surgery patients (Hayoz, Hermann et al. 2018). It involves vertical division of the stomach reducing its size to about 25% (Figure 6.1). It does not affect gastric function or digestion as it does not involve reconstruction of the pylorus or the intestines. Complications of SG include, leakage from the stomach and vomiting from over-eating. RYGB combines gastric restriction with malabsorption techniques where the stomach is divided into 2 parts; a small proximal pouch (15-20mL) and a large distal pouch. The small proximal pouch is attached to the proximal jejunum, bypassing the large distal gastric pouch and duodenum (Figure 6.1). A small stomach pouch and bypassing the duodenum decreases the digestion and absorption of food, thereby, restricting energy intake (Jaunoo, Southall 2010, Arceo-Olaiz, Espana-Gomez et al. 2008, Yan, Ko et al. 2008). Complications of RYGB include, anastomotic leak, delayed gastric emptying, vomiting and dumping syndrome. These procedures were believed to be primarily effective due to their restrictive and malabsorptive properties, however, growing evidence supports the hypotheses that alterations in the gut anatomy have profound effects on the gut physiology, including alterations in gut hormones and their effects on regulating feeding and metabolism (Frikke-Schmidt, O'Rourke et al. 2016). Hormones involved in regulating appetite, satiety and insulin production, such as, gastrointestinal insulinotropic polypeptide (GIP), glucagon-like peptide 1 (GLP-1) and peptide YY (PYY) are

upregulated following surgery, however, the exact mechanism by which bariatric surgery has this effect is unclear (Goktas, Moustaid-Moussa et al. 2013).

Furthermore, bariatric surgery induces significant changes in the anatomical localisation and physiology of the adipose tissue. Up to 50% of the adipose tissue is lost in the first year following bariatric surgery with significant changes in lipid deposition and improvements in metabolic abnormalities (Galanakis, Daskalakis et al. 2015). Hypertrophic adipocytes have altered intracellular functions, induce inflammation and insulin resistance in obesity (Wood, de Heredia et al. 2009). Adipose tissue mass that is lost in response to bariatric surgery is associated with reduced adipocyte hypertrophy, rather than adipocyte hyperplasia, thereby, returning the intracellular functions to baseline and reducing associated abnormalities (Frikke-Schmidt, O'Rourke et al. 2016). Specific changes in adipose tissue depots are also considered to induce metabolic improvements compared to total loss of adipose tissue (Frikke-Schmidt, O'Rourke et al. 2016). For example, visceral adipose tissue (VAT) has increased lipolytic capacity, inflammation, vascularization, increased secretion of adipokines compared to subcutaneous adipose tissue (SAT) and is an independent risk factor for type 2 diabetes mellitus (T2DM) (regardless of fasting insulin levels, insulin secretion, total adiposity and family history) and cardiovascular disease (CVD) (Boyko, Fujimoto et al. 2000, Liu, Fox et al. 2010, Gallagher, Kelley et al. 2009). Reduction in the percentage of VAT induced by bariatric surgery is significantly associated with improvements in insulin sensitivity and remission of T2DM, reduction in flux of free fatty acids and improvements in the inflammatory status (Karki, Farb et al. 2015, Faria, Pestana et al. 2014). These improvements are achieved through multiple pathways associated with systemic and adipose tissue inflammation (Goktas, Moustaid-Moussa et al. 2013).

Adipokines and cytokines are synthesised and secreted by adipocytes and immune cells are active factors that modulate local and systemic metabolism and are detectable in the adipose tissue and the systemic circulation. Dysregulated synthesis and secretion of these adipokines by hypertrophic adipocytes in obesity is hypothesized to contribute to the development of chronic low-grade inflammation, impairing glucose metabolism and insulin sensitivity leading to the onset of insulin resistance and other metabolic co-morbidities (Lindegaard, Jorgensen et al. 2015). In obesity, both circulating and tissue levels of pro-inflammatory proteins are increased, whilst anti-inflammatory proteins are decreased, a situation that is reversed following bariatric surgery, aiding in improving the metabolic profile (Pardina, Ferrer et al. 2012, Viana, Araujo-Dasilio et al. 2013, Sams, Blackledge et al. 2016). Plasma levels and adipocyte expression levels of Annexin A1 (ANXA1) are altered in obesity, however, whether this alteration is reversed with weight loss has not been investigated (Kosicka, Cunliffe et al. 2013). Furthermore, results from Chapter 4 of this thesis suggest, ANXA1 regulates the gene expression of dipeptidyl peptidase 4 (DPP4), thereby possibly modulating the synthesis of gut hormone, GLP-1, therefore, it is of significant interest to investigate the plasma levels of ANXA1 and its effects in response to bariatric surgery.

6.3 Aim, objectives and Hypothesis

The primary aim of this pilot study was to investigate the plasma levels of ANXA1 in response to rapid weight loss through bariatric surgery and correlate it with plasma lipids, plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormone. The objectives of this pilot study were:

- To analyse anthropometric and plasma lipid changes in bariatric surgery patients pre-operatively (pre-opt), 6 weeks and 3 months post-operatively (post-opt)
- To analyse and correlate plasma ANXA1 levels with plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormone in bariatric surgery patients pre-opt, 6 weeks and 3 months post-opt
- To analyse and correlate plasma ANXA1 levels with plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormone by types of bariatric procedures in bariatric surgery patients pre-opt, 6 weeks and 3 months post-opt
- To analyse and correlate plasma ANXA1 levels with plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormone in non-T2DM and T2DM bariatric surgery patients pre-opt, 6 weeks and 3 months post-opt


It was hypothesised that weight loss would significantly increase plasma ANXA1 levels, would inversely correlate with pro-inflammatory biomarkers and directly correlate with anti-inflammatory biomarkers, markers of insulin sensitivity and plasma gut hormone, therefore, effectively aiding the resolution of inflammation and decreasing the risk of developing metabolic co-morbidities.

6.4 Materials and Methods

6.4.1 Participants and Experimental design

Following Coventry University (P50481) and NHS (IRAS:220666) ethical approvals, participants were recruited from the general surgery pre-operative assessment clinic at Heartlands Hospital, Birmingham. 52 patients who were scheduled to undergo bariatric surgery (Sleeve Gastrectomy or Roux-en-Y gastric bypass) were screened and identified using the general surgery pre-operative assessment booking database. 34 patients meeting the inclusion criteria (Male/Female adults, 18-75 years old, BMI>40kg/m² if do not have T2DM or BMI>35kg/m² if known to have T2DM) were approached at their pre-operative assessment appointment and the study design and, the aims were discussed with them (Flow chart 6.1). They were provided with study information sheets and any questions and apprehensions were discussed. 27 patients were happy to take part in the study and provided written consent. Following consent, the baseline clinical assessment (weight and height) was measured by experienced Health Care professionals at the pre-operative assessment clinics and the relevant data (age, gender, past medical history, type of bariatric surgery, co-morbidities, weight, height and BMI) was extracted from patient notes. A number of patients withdrew from the study due to reasons listed below:

Study Recruits		
Total Recruits	27	
No of Pre-opt samples	20/27	5 withdrew before pre-opt bloods <ul style="list-style-type: none"> - 1 transferred to a different site for opt - 1 opted out on their own will - 1 did not meet inclusion criteria (BMI <40kg/m²) + no longer wanted opt. - 2 surgeries cancelled due to other medical reasons 1 withdrew before pre-opt bloods, but decided to continue post-opt 1; surgery kept rearranging thus missed pre-opt bloods
No of 6 Weeks Post-opt samples	16/20	4 withdrew before 6 Weeks bloods <ul style="list-style-type: none"> - 1 did not attend post opt appoint (did call to arrange but patient was unable to attend) - 1 withdrew after operation - 1 cancelled operation (no longer wanting)
No of 3 Months Post-opt samples	9/16	7 withdrew before 3 Months bloods <ul style="list-style-type: none"> - 2 I could not meet - 1 left without seeing me - 2 rearranged - 1 had side effects from OGGT - 1 no show



Flow Chart 6.1: Flow chart depicting the process of patient recruitment including the inclusion and exclusion criteria.

Following consent, 10mL of whole blood was collected in purple top anticoagulant-treated tubes (EDTA-treated) via venepuncture for biomarker analysis following their routine bloods by a trained professional (baseline measures). Patients were contacted after their surgery to confirm their willingness to continue their participation. Patients who were happy to continue were met at their 6 weeks post bariatric surgery follow up appointment at the Diabetes and Endocrinology centre at Heartlands Hospital, Birmingham. The clinical assessments (weight, height) and 10mL of whole blood was collected in EDTA-treated tubes via venepuncture for biomarker analysis following routine bloods by a trained professional. Similarly, patients were contacted prior to their 3 months post bariatric surgery follow up to confirm their willingness to continue their participation. Patients who were happy to continue were met at their 3 months post bariatric surgery follow up appointment at the Outpatients Department at Heartlands Hospital, Birmingham. The clinical assessments (weight and height) and 10mL of whole blood was collected in EDTA-treated tubes via venepuncture for biomarker analysis following routine bloods by a trained professional.

Routine blood samples were sent to the local routine pathology lab within the hospital for analysis of the following biochemical measurements; total cholesterol, low density lipoprotein-cholesterol (LDL-C), high density lipoprotein-cholesterol (HDL-C) and triglycerides.

The blood samples collected in EDTA-treated tubes were centrifuged at 1200 rpm for 15 minutes, at 4°C. The plasma was aliquoted (0.5-1.0mL) into fresh tubes and stored at -80°C.

6.4.2 Metabolic measurements

6.4.2.1 Enzyme linked immunosorbent Assays

Plasma inflammatory biomarkers, fasting plasma insulin levels and fasting plasma gut hormones were quantified using enzyme linked immunosorbent assay (ELISA). The assays were based on the sandwich technique, as described in Chapter 3 (section 3.3).

- The R&D systems Human C-Reactive Protein (CRP) Quantikine ELISA kit (#DCRP00) was used to quantify CRP in plasma samples, following the manufacturers protocol. The mean intra-assay precision was determined using coefficient of variation (CV). The mean CV of the CRP assay was 2.76%.
- The Cusabio Human Annexin I (ANXAI) ELISA kit (#CSB-E12155h) was used to quantify Annexin A1 levels in plasma samples, following the manufacturers protocol. The mean %CV of the ANXA1 assay was 6.38%.
- The R&D systems Human Interleukin 1 β (IL-1 β) Quantikine ELISA kit (#DLB50) was used to quantify IL-1 β in plasma samples, following the manufacturers protocol. The mean %CV of the IL-1 β assay was 4.75%.
- The R&D systems Human Interleukin 10 (IL-10) Quantikine ELISA kit (#D1000B) was used to quantify IL-10 in plasma samples, following the manufacturers protocol. The mean %CV of the IL-10 assay was 6.14%.
- The R&D systems Human Tumour Necrosis Factor α (TNF α) Quantikine ELISA kit (#DTA00D) was used to quantify TNF α in plasma samples, following the manufacturers protocol. The mean %CV of the IL-10 assay was 1.78%.
- The R&D systems Human Leptin Quantikine ELISA kit (#DLP00) was used to quantify Leptin in plasma samples. The mean %CV of the Leptin assay was 3.22%.

- The R&D systems Human Adiponectin Quantikine ELISA kit (#DRP300) was used to quantify Adiponectin in plasma samples. The mean %CV of the Adiponectin assay was 8.89%.
- The DRG systems Human Insulin ELISA kit (#DX-EIA-2935) was used to quantify Insulin in fasted plasma samples. The mean %CV of the insulin assay was 5.05%.
- The Diagenics Human GLP-1 ELISA kit (#EH1053) was used to quantify Insulin in fasted plasma samples, following the manufacturers protocol. The mean %CV of the GLP-1 assay was 9.1%.

Plasma glucose levels were measured using the Biosen C.Line EKF Diagnostic machine following the manufacturers protocol. Briefly, following calibration, 20µL of fasting plasma sample was measured twice. A mean was calculated to determine the concentration of glucose in the patient samples.

6.4.2.2 Calculation of insulin resistance, sensitivity and pancreatic β cell function

The homeostasis model assessment (HOMA) was used to estimate insulin resistance (HOMA-IR), insulin sensitivity (HOMA-%S) and pancreatic β -cell function (HOMA-% β) using fasted glucose and insulin levels. The following equations were used to calculate HOMA-IR, HOMA-%S and HOMA-% β (Glucose is mmol/L and Insulin μ IU/mL) (Crofts, C., Wheldon, M., Zinn, C., Lan-Pidhainy, X., Wolever, T., Schofield, G. 2017):

$$\text{HOMA-IR} = (\text{Glucose} \times \text{Insulin}) / 22.5$$

$$\text{HOMA-\%S} = 1/\text{HOMA-\%}\beta$$

$$\text{HOMA-\%}\beta = (20 \times \text{Insulin}) / (\text{Glucose} - 3.5)$$

6.4.3 Statistical analysis

Data presented in this study is expressed as Mean \pm Standard Error of the Mean (SEM). GraphPad Prism 8 was used to statistically analyse the data and produce the graphs. One-way and Two-way ANOVA, Tukey's post hoc test and one-tailed Pearson's correlation coefficient were used to statistically analyse the data and to assess the differences in the data sets, a p value of $p < 0.05$ was considered statistically significant.

6.5 Results

Anthropometric and plasma samples of consented bariatric surgery patients were analysed. Plasma samples were analysed for plasma lipids, inflammatory biomarkers, markers of insulin sensitivity and gut hormones pre-opt, 6 weeks and 3 months post-operatively post-opt and correlated with plasma ANXA1 levels. The participant cohort was further subdivided into types of bariatric surgery (SG vs RYGB) and T2DM status (non-T2DM vs T2DM) for some of the results presented in this chapter.

6.5.1 Pre and post bariatric surgery: Full cohort

The participant characteristics of the complete cohort are presented in table 6.1. Of the 26 participants at pre-opt, 4 were male and 22 were female, with an average age of 48 ± 1.9 . At 6 weeks, 16 females (average age 46 ± 2.6) and at 3 months, 11 females (average age 51 ± 2.7) continued their participation.

Participant Characteristics				
		Pre-Opt	6 Weeks Post-Opt	3 Months Post-Opt
Number of participants		26	16	11
Gender				
	Males	4	0	0
	Females	22	16	11
Age, years				
		48±1.9 (n=26)	46±2.6 (n=16)	51±2.7 (n=11)
Type of Surgery				
	Sleeve Gastrectomy	13	7	5
	Roux-en-Y Gastric Bypass	13	9	6
Type 2 diabetes mellitus				
	Diabetics	10	7	4
	Non-Diabetics	16	9	7
Anthropometric Measures				
	Weight (kg)	131.5±6.4 (n=26)	121.9±5.5 (n=15)	105.9±5.3 [#] (n=9)
	BMI (kg/m ²)	49.7±1.2 (n=26)	44.6±1.6* (n=15)	38.8±1.5*** (n=9)
Lipid Profiles				
	Triglycerides (mmol/L)	1.3±0.1 (n=25)	1.5±0.1 (n=15)	1.4±0.3 (n=9)
	Total Cholesterol (mmol/L)	4.5±0.1 (n=25)	4.7±0.2 (n=15)	4.5±0.4 (n=9)
	HDL (mmol/L)	1.2±0.1 (n=25)	1.1±0.1 (n=15)	1.4±0.1 [#] (n=8)
	LDL (mmol/L)	2.7±0.1 (n=25)	2.9±0.2 (n=15)	2.5±0.2 (n=9)

Table 6.1: Participant characteristics at pre and post- Bariatric surgery. The anthropometric measures and lipid profiles were analysed in participants undergoing bariatric surgery at pre-, 6 weeks and 3 months post-operatively. One-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *vs pre opt, #vs 6 weeks
Pre-opt; Pre-operatively, Post-opt; Post-operatively, BMI; Body mass index, HDL; High density lipoprotein and LDL; Low density lipoprotein

6.5.1.1 Anthropometric and plasma lipid levels

On average, the average weight was significantly reduced at 3 months post-opt compared to pre-opt (131.5 ± 6.4 kg, $n=26$ vs 105.9 ± 5.3 kg, $n=9$, $p=0.049$, respectively) (Table 6.1). Similarly, the average BMI was significantly reduced at 6 weeks and 3 months post-opt compared to pre-opt (44.6 ± 1.6 kg/m², $n=15$, $p=0.025$ and 38.8 ± 1.5 kg/m², $n=9$, $p<0.0001$ vs 49.7 ± 1.2 kg/m², $n=26$, respectively), as shown in Figure 6.2.




Figure 6.2: Average BMI (kg/m²) of participant cohort pre- and post- Bariatric surgery. The average Body mass index was significantly reduced at 6 weeks and 3 months post-opt compared to pre-opt ($p=0.025$, $n=15$ and $p<0.0001$, $n=9$, respectively). One-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$. BMI= Body mass index, Pre-opt= Pre-operatively and Post-opt= post-operatively

The changes in the average plasma lipid levels of triglycerides, total cholesterol and LDL post-opt compared to pre-opt did not significantly differ (Table 6.1). However, the average plasma HDL levels significantly improved at 3 months post-opt compared to 6

weeks post-opt (1.4 ± 0.1 mmol/L, n=8 vs 1.1 ± 0.1 mmol/L, n=15, p=0.044, respectively) (Table 6.1).

6.5.1.2 Plasma inflammatory biomarkers, markers of insulin sensitivity and gut hormone

Table 6.2 presents the analysis of plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormones pre- and post- bariatric surgery of the complete participant cohort.

6.5.1.2.1 Plasma inflammatory Biomarkers

On average, plasma C-reactive protein (CRP) levels decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference was not statistically significant (10.1 ± 2.3 μ g/mL, n=14 and 8.0 ± 3.7 μ g/mL, n=6 vs 11.6 ± 2.4 μ g/mL, n=18, respectively).

On average, the plasma interleukin-1 β (IL-1 β) levels significantly increased at 3 months post-opt compared to pre-opt (2.2 ± 0.3 pg/mL, n=9 vs 1.3 ± 0.2 pg/mL, n=17, p=0.019, respectively) and 3 months post-opt compared to 6 weeks post-opt (2.2 ± 0.3 pg/mL, n=9 vs 1.3 ± 0.2 pg/mL, n=14, p=0.027, respectively). On average, the plasma interleukin-10 (IL-10) levels significantly increased at 6 weeks post-opt compared to pre-opt (9.3 ± 2.2 pg/mL, n=13 vs 6.9 ± 0.7 pg/mL, n=18, p=0.048, respectively) and remained at a similar level at 3 months post-opt compared to 6 weeks post-opt (9.1 ± 0.8 pg/mL, n=8 vs 9.3 ± 0.6 pg/mL, n=13, respectively). On average, the plasma tumour necrosis factor α (TNF α) levels remained similar at 6 weeks and 3 months post-opt compared to pre-opt (21.2 ± 0.8 pg/mL, n=15 and 21.7 ± 1.4 pg/mL, n=9 vs 21.7 ± 0.4 pg/mL, n=19, respectively). On average, the plasma leptin levels significantly decreased at 6 weeks and 3 months post-

opt compared to pre-opt (34.6±5.0 ng/mL, n=15, p<0.0001 and 35.7±4.8 ng/mL, n=9, p=0.0001 vs 80.9±6.8 ng/mL, n=20, respectively). On average, the plasma adiponectin

Plasma Biomarkers: Full Cohort				
		Pre-Opt	6 Weeks Post-Opt	3 Months Post-Opt
Inflammatory Biomarkers				
	CRP (µg/mL)	11.6±2.4 (n=18)	10.1±2.3 (n=14)	8.0.6±3.7 (n=6)
	Annexin A1 (ng/mL)	1.1±0.1 (n=20)	1.4±0.2 (n=16)	1.6±0.1 (n=9)
	IL-1β (pg/mL)	1.3±0.2 (n=17)	1.3±0.2 (n=14)	2.2±0.3* [#] (n=9)
	IL-10 (pg/mL)	6.9±0.7 (n=18)	9.3±2.2* (n=13)	9.1±0.8 (n=8)
	TNFα (pg/mL)	21.7±0.4 (n=19)	21.2±0.8 (n=15)	21.7±1.4 (n=9)
	Leptin (ng/mL)	80.9±6.8 (n=20)	34.6±5.0*** (n=15)	35.7±4.8*** (n=9)
	Adiponectin (µg/mL)	5.3±0.6 (n=19)	8.1±0.8* (n=13)	8.9±0.8** (n=8)
Insulin Sensitivity				
	Fasting Glucose (mmol/L)	4.2±0.3 (n=18)	3.9±0.1 (n=9)	4.2±0.8 (n=7)
	Fasting Insulin (µIU/mL)	25.7±1.9 (n=19)	18.1±1.9* (n=9)	14.7±2.4** (n=7)
	HOMA-IR	3.2±0.3 (n=15)	2.2±0.2 (n=9)	1.8±0.3 (n=7)
	HOMA-%S (%)	34.0±2.5 (n=15)	44.6±3.1 (n=8)	57.5±2.47.6*** (n=6)
	HOMA-%β (%)	321.0±27.0 (n=14)	260.4±10.6 (n=9)	252.2±24.0 (n=6)
Gut Hormone				
	Fasting GLP-1 (ng/mL)	264.2±12.0 (n=19)	311.3±16.1 (n=9)	354.5±17.6*** (n=7)

Table 6.2: Plasma biomarkers analysis of participant cohort at pre- and post- Bariatric surgery. Plasma inflammatory biomarkers, fasting plasma glucose and insulin levels and plasma gut hormone were analysed and, HOMA-IR, HOMA-%S and HOMA-%β were calculated using the equations presented in section 6.4.2.2. One-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean±SEM. P value was set at p <0.05 and is denoted by *vs pre opt, [#]vs 6 weeks of the corresponding group

Pre-opt= Pre-operatively, Post-opt= post-operatively, CRP= C-reactive protein, IL= Interleukin, TNFα= Tumour necrosis factor α, HOMA-IR= Homeostatic model assessment of insulin resistance, HOMA-%S= Homeostatic model assessment of insulin sensitivity, HOMA-%β= Homeostatic model assessment of pancreatic β cell function and GLP-1= Glucagon like peptide 1

levels significantly increased at 6 weeks and 3 months post-opt compared to pre-opt ($8.1 \pm 0.8 \mu\text{g/mL}$, $n=13$, $p=0.018$ and $8.9 \pm 0.8 \mu\text{g/mL}$, $n=8$, $p=0.009$ vs $5.3 \pm 0.6 \mu\text{g/mL}$, $n=19$, respectively).

6.4.1.2.2 Plasma markers of insulin sensitivity

On average, the fasting plasma glucose levels remained similar at 6 weeks and 3 months post-opt compared to pre-opt ($3.9 \pm 0.1 \text{ mmol/L}$, $n=9$ and $4.2 \pm 0.8 \text{ mmol/L}$, $n=7$ vs $4.2 \pm 0.3 \text{ mmol/L}$, $n=18$, respectively). However, the average fasting plasma insulin levels significantly decreased at 6 weeks and 3 months post-opt compared to pre-opt ($18.1 \pm 1.9 \mu\text{IU/mL}$, $n=9$, $p=0.047$ and $14.7 \pm 2.4 \mu\text{IU/mL}$, $n=7$, $p=0.006$ vs $25.7 \pm 1.9 \mu\text{IU/mL}$, $n=19$, respectively). On average, HOMA-IR significantly reduced at 6 weeks and 3 months post-opt compared to pre-opt (2.2 ± 0.2 , $n=9$, $p=0.021$ and 1.8 ± 0.3 , $n=7$, $p=0.003$ vs 3.2 ± 0.3 , $n=15$, respectively). On average, HOMA-%S significantly improved at 3 months post-opt compared to pre-opt ($57.5 \pm 7.6 \%$, $n=6$ vs $34 \pm 2.5 \%$, $n=15$, $p=0.0009$, respectively). On average, HOMA-% β reduced at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference was not statistically significant ($260.4 \pm 10.6 \%$, $n=9$ and $252.2 \pm 24.0 \%$, $n=6$ vs $321 \pm 27.0 \%$, $n=14$, respectively).

6.5.1.2.3 Plasma gut hormone

On average, the fasting plasma GLP-1 levels significantly increased at 3 months post-opt compared to pre-opt ($354.5 \pm 17.6 \text{ ng/mL}$, $n=7$ vs $264.2 \pm 12.0 \text{ ng/mL}$, $n=19$, $p=0.0003$ respectively).

6.5.1.3 Annexin A1

On average, the plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference was not statistically significant in the full cohort (1.4 ± 0.2 ng/mL, $n=16$, $p=0.080$ and 1.6 ± 0.1 ng/mL, $n=9$, $p=0.170$, vs 1.1 ± 0.1 ng/mL, $n=20$, respectively), as shown in Figure 6.3.




Figure 6.3: Average plasma Annexin A1 (ng/mL) levels of participant cohort pre- and post- Bariatric surgery. The average plasma Annexin A1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt ($p=0.080$, $n=16$ and $p=0.170$, $n=9$, respectively). One-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$.

Plasma ANXA1 levels inversely correlated with BMI ($r=0.219$, $p=0.077$, $n=44$) (Figure 6.4), directly correlated with plasma HDL levels ($r=0.242$, $p=0.064$, $n=41$) (Figure 6.5), significantly directly correlated with plasma TNF α levels ($r=0.321$, $p=0.018$, $n=43$) (Figure 6.6) and directly correlated with plasma adiponectin levels ($r=0.226$, $p=0.083$, $n=39$) (Figure 6.7), in the complete cohort.




Figure 6.4: Correlation between plasma Annexin A1 levels and BMI in participant cohort. Plasma Annexin A1 levels inversely correlated with BMI. One tailed Person correlation coefficient was used to statistically analyse the data.
BMI= Body mass index




Figure 6.5: Correlation between plasma Annexin A1 levels and plasma HDL levels in participant cohort.
Plasma Annexin A1 levels directly correlated with plasma HDL levels. One tailed Person correlation coefficient was used to statistically analyse the data.
HDL= High density lipoprotein




Figure 6.6: Correlation between plasma Annexin A1 levels and plasma TNFα levels in participant cohort.
Plasma Annexin A1 levels significantly directly correlated with plasma TNFα levels. One tailed Person correlation coefficient was used to statistically analyse the data.
TNFα= Tumour necrosis factor α




Figure 6.7: Correlation between plasma Annexin A1 levels and plasma Adiponectin levels in participant cohort. Plasma Annexin A1 levels directly correlated with plasma Adiponectin levels. One tailed Person correlation coefficient was used to statistically analyse the data.

6.5.2 Pre and post bariatric surgery: Surgery Type

Participant Characteristics: Surgery Type							
		Pre-Opt		6 Weeks Post-Opt		3 Months Post-Opt	
		SG	RYGB	SG	RYGB	SG	RYGB
Number of participants		13	13	7	9	5	6
Gender							
	Males	2	2	0	0	0	0
	Females	11	11	7	9	5	6
Age, years							
		51±2.3 (n=13)	45±2.9 (n=13)	49±4.1 (n=7)	44±3.3 (n=9)	54±2.4 (n=5)	48±4.2 (n=6)
Type 2 diabetes mellitus							
	Diabetics	5	5	3	4	2	2
	Non-Diabetics	8	8	4	5	3	4
Anthropometric Measures							
	Weight (kg)	136.1±5.3 (n=13)	138±5.4 (n=13)	117.5±7.0 (n=7)	125.8±8.0 (n=9)	104.6±7.6 [#] (n=5)	107.5±8.3* (n=4)
	BMI (kg/m ²)	49.9±1.8 (n=13)	49.5±1.5 (n=13)	43.3±2.4 (n=7)	45.8±2.1 (n=9)	38.8±2.1* (n=5)	38.8±2.5* (n=4)
Lipid Profiles							
	Triglycerides (mmol/L)	1.3±0.1 (n=12)	1.3±0.1 (n=13)	1.5±0.1 (n=7)	1.5±0.1 (n=8)	2.0±0.5 (n=4)	1.0±0.1 [#] (n=5)
	Total Cholesterol (mmol/L)	4.4±0.1 (n=12)	4.5±0.2 (n=13)	5.1±0.4 (n=7)	4.5±0.2 (n=8)	5.2±0.6 (n=4)	4.0±0.3 (n=5)
	HDL (mmol/L)	1.3±0.1 (n=12)	1.1±0.1 (n=13)	1.2±0.1 (n=7)	1.0±0.04 (n=8)	1.5±0.1 (n=4)	1.2±0.1 (n=5)
	LDL (mmol/L)	2.5±0.1 (n=12)	2.9±0.2 (n=13)	3.1±0.4 (n=7)	2.8±0.2 (n=8)	2.8±0.3 (n=4)	2.3±0.2 (n=5)

Table 6.3: Participant characteristics at pre- and post- Sleeve Gastrectomy and Roux-en-Y Gastric Bypass. The anthropometric measures and lipid profiles were analysed in participants undergoing bariatric surgery at pre-, 6 weeks and 3 months post-operatively and compared between surgery types (SG vs RYGB). Two-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *vs pre opt, #vs 6 weeks of the corresponding group. Pre-opt= Pre-operatively, Post-opt= Post-operatively, SG= Sleeve gastrectomy, RYGB= Roux-en-Y Gastric Bypass, BMI= Body mass index, HDL= High density lipoprotein and LDL= Low density lipoprotein

The characteristics of the SG and RYGB cohorts are presented in table 6.3. Of the 13 SG participants at pre-opt, 2 were male and 11 were female, with an average age of 51 ± 2.3 . At 6 weeks, 7 females (average age 49 ± 4.1) and at 3 months, 5 females (average age 54 ± 2.4) continued their participation. Similarly, of the 13 RYGB participants at pre-opt, 2 were males and 11 were females, with an average age of 45 ± 2.9 . At 6 weeks, 9 females (average age 44 ± 3.3) and at 3 months, 6 females (average age 48 ± 4.2) continued their participation.

6.5.2.1 Anthropometric and plasma lipid levels

The effect of time ($p=0.001$) but not surgery type ($p=0.840$) nor it's interaction with time ($p=0.773$) was statistically significant on the average weight loss. The effect of time ($p=0.003$) (but not surgery type ($p=0.724$) nor it's interaction with time ($p=0.954$)) was statistically significant on the average change in BMI. Furthermore, post-hoc analysis did not detect specific significant differences in the average weight and BMI between the cohorts at any time point.

However, the average weight decreased in both cohorts at post-opt compared to pre-opt. On average, in the SG cohort the average weight was significantly reduced at 3 months post-opt compared to 6 weeks post opt (104.6 ± 7.6 kg, $n=5$ vs 117.5 ± 7.0 kg, $n=7$, $p=0.025$ respectively) (Table 6.3). Similarly, in the RYGB the average weight was significantly reduced at 3 months post-opt compared to pre-opt (107 ± 8.3 kg, $n=4$ vs 138 ± 5.4 kg, $n=13$, $p=0.040$, respectively) (Table 6.3). Furthermore, the average BMI was reduced in both cohorts at post-opt compared to pre-opt. On average, in the SG cohort the average BMI was significantly reduced at 3 months post-opt compared to 6 weeks post-opt (39.0 ± 2.1 kg/m², $n=5$ vs 43.3 ± 2.4 kg/m², $n=7$, $p=0.0247$, respectively), as shown in Figure 6.8.

Similarly, in the RYGB cohort the average BMI was significantly reduced at 3 months post-opt compared to pre-opt ($38.6 \pm 2.5 \text{ kg/m}^2$, $n=4$ vs $49.5 \pm 1.5 \text{ kg/m}^2$, $n=13$, $p=0.039$, respectively), as shown in Figure 6.8.




Figure 6.8: Average BMI (kg/m^2) of participant cohort pre- and post- Sleeve gastrectomy and Roux-en-Y Gastric Bypass. The average BMI was significantly reduced at 3 months post-opt compared to 6 weeks post-opt in the SG cohort ($p=0.037$, $n=13$). Similarly, the average BMI was significantly reduced at 3 months post-opt compared to pre-opt in the RYGB cohort ($p=0.038$, $n=13$). However, no statistically significant differences were observed between the cohorts at any time point. Two-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *vs SG pre opt, #vs RYGB pre-opt of the corresponding group BMI= Body mass index, SG= Sleeve gastrectomy, RYGB= Roux-en-Y Gastric Bypass, Pre-opt= Pre-operatively, Post-opt= Post-operatively

The effect of surgery type was statistically significant on the average plasma triglyceride levels ($p=0.013$), plasma total cholesterol levels ($p=0.026$) and plasma HDL levels ($p=0.015$), however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. However, no significant difference was observed in the average plasma LDL levels in either cohorts at any timepoint.

6.5.2.2 Plasma inflammatory biomarkers, markers of insulin sensitivity and gut hormone

Table 6.4 presents the analysis of plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormones pre- and post- SG and RYGB.

6.5.2.2.1 Plasma inflammatory biomarkers

The effect of time ($p=0.184$), surgery type ($p=0.232$) nor their interaction ($p=0.429$) was statistically significant on the average plasma CRP levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average plasma CRP levels decreased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma CRP levels decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (10.6 ± 3.7 $\mu\text{g/mL}$, $n=7$ and 9.5 ± 4.2 $\mu\text{g/mL}$, $n=5$ vs 12.9 ± 3.5 $\mu\text{g/mL}$, $n=11$, respectively). Similarly, in the RYGB cohort, the average plasma CRP levels decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (9.6 ± 3.0 $\mu\text{g/mL}$, $n=7$ and 3.0 ± 2.5 $\mu\text{g/mL}$, $n=2$ vs 9.6 ± 2.8 $\mu\text{g/mL}$, $n=7$, respectively).

The effect of time ($p=0.016$) and surgery type ($p=0.016$) but not their interaction ($p=0.864$) was statistically significant on the average plasma IL-1 β levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any timepoint. However, the average plasma IL-1 β levels increased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma IL-1 β levels increased at 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (2.0 ± 0.2 pg/mL , $n=5$ vs 1.0 ± 0.2 pg/mL , $n=10$,

respectively). Similarly, in the RYGB cohort, the average plasma IL-1 β levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (1.6 ± 0.2 pg/mL, n=9 and 2.5 ± 0.6 pg/mL, n=4 vs 1.8 ± 0.3 pg/mL, n=7, respectively).

The effect of time (p=0.056), surgery type (p=0.661) nor their interaction (p=0.940) was statistically significant on the average plasma IL-10 levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average plasma IL-10 levels increased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma IL-10 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (9.5 ± 1.0 pg/mL, n=6 and 9.0 ± 0.6 pg/mL, n=5 vs 6.9 ± 1.1 pg/mL, n=11, respectively). Similarly, in the RYGB cohort, the average plasma IL-10 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (10.1 ± 1.2 pg/mL, n=8 and 9.1 ± 2.0 pg/mL, n=3 vs 6.9 ± 0.7 pg/mL, n=7, respectively).

The effect of time (p=0.914), surgery type (p=0.653) nor their interaction (p=0.748) was statistically significant on the average plasma TNF α levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average plasma TNF α levels remained similar in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma TNF α levels remained similar at 6 weeks and 3 months post-opt (22.0 ± 1.3 pg/mL, n=6 and 21.8 ± 2.1 pg/mL, n=5 vs 21.9 ± 0.6 pg/mL, n=11, respectively). Similarly, in the RYGB cohort, the average plasma TNF α levels remained similar at 6 weeks and 3 months post-opt compared to pre-opt (20.6 ± 1.1 pg/mL, n=9 and 21.6 ± 1.8 pg/mL, n=4 vs 21.4 ± 0.5 pg/mL, n=8, respectively).

The effect of time ($p=0.001$) but not the surgery type ($p=0.994$) nor their interaction ($p=0.551$) was statistically significant on the average plasma leptin levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any timepoint. However, the average plasma leptin levels decreased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma leptin levels significantly decreased at 6 weeks and 3 months post-opt, however, the difference observed was not statistically significant (35.0 ± 6.3 ng/mL, $n=7$ and 32.8 ± 8.4 ng/mL, $n=5$ vs 85.3 ± 9.1 ng/mL, $n=12$, respectively). A similar pattern was observed in the RYGB cohort, the average plasma leptin levels significantly decreased at 6 weeks post-opt compared to pre-opt, however, the difference observed was not statistically significant (36.1 ± 7.4 ng/mL, $n=9$ vs 74.3 ± 10.5 ng/mL, $n=8$, $p=0.004$, respectively) and at 3 months post-opt compared to pre-opt (39.2 ± 4.0 ng/mL, $n=4$ and 74.3 ± 10.5 ng/mL, $n=8$, respectively).

The effect of time ($p=0.015$) but not the surgery type ($p=0.397$) nor their interaction ($p=0.257$) was statistically significant on the average plasma adiponectin levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any timepoint. However, the average plasma adiponectin levels increased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma adiponectin levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (8.9 ± 1.6 μ g/mL, $n=6$ and 8.0 ± 1.1 μ g/mL, $n=5$ vs 5.7 ± 0.8 μ g/mL, $n=11$, respectively). Similarly, in the RYGB cohort, the average plasma adiponectin levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (7.6 ± 1.2 μ g/mL, $n=9$ and 10.5 ± 0.8 μ g/mL, $n=3$ vs 6.0 ± 0.7 μ g/mL, $n=6$, respectively).

6.5.2.2.2 Plasma biomarkers of insulin sensitivity

The effect of time ($p=0.933$), surgery type ($p=0.878$) nor their interaction ($p=0.765$) was statistically significant on the average fasting plasma glucose levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average fasting plasma glucose levels remained similar in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average fasting plasma glucose levels remained similar at 6 weeks and 3 months post-opt compared to pre-opt (4.1 ± 0.4 mmol/L, $n=5$ and 3.8 ± 0.1 mmol/L, $n=3$ vs 4.1 ± 0.2 mmol/L, $n=11$, respectively). Similarly, in the RYGB cohort, the average fasting plasma glucose levels remained similar at 6 weeks and 3 months post-opt compared to pre-opt (4.0 ± 0.2 mmol/L, $n=4$ and 4.4 ± 0.5 mmol/L, $n=4$ vs 3.9 ± 0.8 mmol/L, $n=8$, respectively).

The effect of time ($p=0.063$), surgery type ($p=0.126$) nor their interaction ($p=0.563$) was statistically significant on the average fasting plasma insulin levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average fasting plasma insulin levels decreased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average fasting plasma insulin levels significantly decreased at 6 weeks compared to pre-opt, however, the difference observed was not statistically significant (18.2 ± 3.5 μ U/mL, $n=5$ vs 29.2 ± 2.9 μ U/mL, $n=11$, $p=0.032$, respectively). Similarly, in the RYGB cohort, the average fasting plasma insulin levels decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (18.0 ± 0.9 μ U/mL, $n=4$ and 13.8 ± 2.8 μ U/mL, $n=4$ vs 21.0 ± 1.4 μ U/mL, $n=8$, respectively).

The effect of surgery type ($p=0.007$) but not time ($p=0.553$) nor their interaction ($p=0.723$) was statistically significant on the average HOMA-IR, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any timepoint. However, the average HOMA-IR remained similar in both at post-opt compared to pre-opt. In the SG cohort, the average HOMA-IR remained similar at 6 weeks and 3 months post-opt compared to pre-opt (3.6 ± 0.4 , $n=6$ and 3.3 ± 0.7 , $n=3$ vs 3.6 ± 0.4 , $n=9$, respectively). Similarly, in the RYGB cohort, the average HOMA-IR remained similar at 6 weeks and 3 months post-opt compared to pre-opt (2.8 ± 0.6 , $n=2$ and 2.7 ± 0.3 , $n=4$ vs 2.6 ± 0.2 , $n=6$, respectively).

The effect of time ($p=0.492$), surgery type ($p=0.054$) nor their interaction ($p=0.969$) was statistically significant on the average HOMA-%S, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average HOMA-%S remained similar in both cohorts post-opt compared to pre-opt. In the SG cohort, the average HOMA-%S remained similar at 6 weeks and 3 months post-opt compared to pre-opt (30.3 ± 4.4 %, $n=6$ and 33.7 ± 8.8 %, $n=3$ vs 30.8 ± 3.5 %, $n=9$, respectively). Similarly, in the RYGB cohort, the average HOMA-%S remained similar at 6 weeks and 3 months post-opt compared to pre-opt (2.8 ± 0.6 %, $n=2$ and 2.7 ± 0.3 %, $n=4$ vs 2.6 ± 0.2 %, $n=6$, respectively).

The effect of time ($p=0.285$), surgery type ($p=0.289$) nor their interaction ($p=0.879$) was statistically significant on the average HOMA-% β , therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average HOMA-% β reduced in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average HOMA-% β reduced at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (267.2 ± 16.2 %, $n=5$ and 252.2 ± 42.2

%, n=3 vs 326.4 ± 42.4 %, n=9, respectively). Similarly, in the RYGB cohort, the average HOMA-% β reduced at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (251.9 ± 13.6 %, n=4 and 204.2 ± 53.2 %, n=4 vs 269.7 ± 42.2 %, n=6, respectively).

6.5.2.2.3 Plasma gut hormone

The effect of time (p=0.001) but not the surgery type (p=0.998) nor their interaction (p=0.496) was statistically significant on the average plasma GLP-1 levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any timepoint. However, the average plasma GLP-1 levels increased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma GLP-1 levels increased at 6 weeks (308.2 ± 24.1 ng/mL, n=5 vs 259.7 ± 17.4 ng/mL, n=11, respectively) and significantly 3 months post-opt compared to pre-opt (376.8 ± 17.7 ng/mL, n=3 vs 259.7 ± 17.4 ng/mL, n=11, p<0.01) respectively). Similarly, in the RYGB cohort, the average plasma GLP-1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (315.0 ± 23.9 ng/mL, n=4 and 337.7 ± 26.6 ng/mL, n=4 vs 270.3 ± 16.8 ng/mL, n=8, respectively).

Plasma Biomarkers: Surgery Type							
		Pre-opt		6 Weeks Post-opt		3 Months Post-opt	
Surgery Type		SG	RYGB	SG	RYGB	SG	RYGB
Inflammatory Biomarkers							
	CRP ($\mu\text{g/mL}$)	12.9 \pm 3.5 (n=11)	9.6 \pm 2.8 (n=7)	10.6 \pm 3.7 (n=7)	9.6 \pm 3.0 (n=7)	9.5 \pm 4.2 (n=5)	3.0 \pm 2.5 (n=2)
	Annexin A1 (ng/mL)	1.1 \pm 0.1 (n=12)	1.1 \pm 0.1 (n=8)	1.7 \pm 0.3 (n=7)	1.2 \pm 0.1 (n=9)	1.6 \pm 0.2 (n=5)	1.6 \pm 0.3 (n=4)
	IL-1 β (pg/mL)	1.0 \pm 0.2 (n=10)	1.8 \pm 0.3 (n=7)	1.1 \pm 0.2 (n=6)	1.6 \pm 0.2 (n=9)	2.0 \pm 0.2 (n=5)	2.5 \pm 0.6 (n=4)
	IL-10 (pg/mL)	6.9 \pm 1.1 (n=11)	6.9 \pm 0.7 (n=7)	9.5 \pm 1.0 (n=6)	10.1 \pm 1.2 (n=8)	9.0 \pm 0.6 (n=5)	9.1 \pm 2.0 (n=3)
	TNF α (pg/mL)	21.9 \pm 0.6 (n=11)	21.4 \pm 0.5 (n=8)	22.0 \pm 1.3 (n=6)	20.6 \pm 1.1 (n=9)	21.8 \pm 2.1 (n=5)	21.6 \pm 1.8 (n=4)
	Leptin (ng/mL)	85.3 \pm 9.1 (n=12)	74.3 \pm 10.5 (n=8)	35.0 \pm 6.3 (n=7)	36.1 \pm 7.4** (n=9)	32.8 \pm 8.4 (n=5)	39.2 \pm 4.0 (n=4)
	Adiponectin ($\mu\text{g/mL}$)	5.7 \pm 0.8 (n=11)	6.0 \pm 0.7 (n=6)	8.9 \pm 1.6 (n=6)	7.6 \pm 1.2 (n=9)	8.0 \pm 1.1 (n=5)	10.5 \pm 0.8 (n=3)
Insulin Sensitivity							
	Fasting Glucose (mmol/L)	4.1 \pm 0.2 (n=11)	3.9 \pm 0.8 (n=8)	4.1 \pm 0.4 (n=5)	4.0 \pm 0.2 (n=4)	3.8 \pm 0.1 (n=3)	4.4 \pm 0.5 (n=4)
	Fasting Insulin ($\mu\text{IU/mL}$)	29.2 \pm 2.9 (n=11)	21.0 \pm 1.4 (n=8)	18.2 \pm 3.5* (n=5)	18.0 \pm 0.9 (n=4)	15.8 \pm 5.0 (n=3)	13.8 \pm 2.8 (n=4)
	HOMA-IR	3.6 \pm 0.4 (n=9)	2.6 \pm 0.2 (n=6)	3.6 \pm 0.4 (n=6)	2.8 \pm 0.6 (n=2)	3.3 \pm 0.7 (n=3)	2.7 \pm 0.3 (n=4)
	HOMA-%S (%)	30.8 \pm 3.5 (n=9)		30.3 \pm 4.4 (n=6)		33.7 \pm 8.8 (n=3)	
	HOMA-% β (%)	326.4 \pm 42.2 (n=9)	269.7 \pm 42.2 (n=6)	267.2 \pm 16.2 (n=2)	251.9 \pm 13.6 (n=4)	252.2 \pm 42.2 (n=3)	204.2 \pm 53.2 (n=4)
Gut Hormone							
	GLP-1 (ng/mL)	259.7 \pm 17.4 (n=11)	270.3 \pm 16.8 (n=8)	308.2 \pm 24.1 (n=5)	315.0 \pm 23.9 (n=4)	376.8 \pm 17.7** (n=3)	337.7 \pm 26.6 (n=4)

Table 6.4: Plasma biomarkers analysis of participant cohort at pre- and post- Sleeve Gastrectomy and Roux-en-Y Gastric Bypass. Plasma inflammatory biomarkers, fasting plasma glucose and insulin levels and plasma gut hormone were analysed and, HOMA-IR, HOMA-%S and HOMA-% β were calculated using the equations presented in section 6.4.2.2. One-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *vs pre opt of the corresponding group
Pre-opt= Pre-operatively, Post-opt= Post-operatively, SG= Sleeve gastrectomy, RYGB= Roux-en-Y Gastric Bypass, CRP= C-reactive protein, IL= Interleukin, TNF α = Tumour necrosis factor α , HOMA-IR= Homeostatic model assessment of insulin resistance, HOMA-%S= Homeostatic model assessment of insulin sensitivity, HOMA-% β = Homeostatic model assessment of pancreatic β cell function and GLP-1= Glucagon like peptide 1

6.5.2.3 Annexin A1

The effect of time ($p=0.058$), surgery type ($p=0.321$) nor their interaction ($p=0.283$) was statistically significant on the average plasma ANXA1 levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, plasma ANXA1 levels increased in both cohorts at post-opt compared to pre-opt. In the SG cohort, the average plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (1.7 ± 0.3 ng/mL, $n=7$ and 1.6 ± 0.2 ng/mL, $n=5$ vs 1.1 ± 0.1 ng/mL, $n=12$, respectively), as shown in Figure 6.9. Similarly, in the RYGB cohort, the average plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (1.2 ± 0.1 ng/mL, $n=9$ and 1.6 ± 0.3 ng/mL, $n=4$ vs 1.1 ± 0.1 ng/mL, $n=8$, respectively), as shown in Figure 6.9.




Figure 6.9: Average plasma Annexin A1 (ng/mL) levels of participant cohort pre- and post- Sleeve Gastrectomy and Roux-en-Y Gastric Bypass. The average plasma Annexin A1 levels increased at 3 months post-opt compared to 6 weeks post-opt and pre-opt in both cohorts, however, no statistically significant difference was observed at any time point within and between cohorts. Two-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$. SG= Sleeve gastrectomy, RYGB= Roux-en-Y Gastric Bypass, Pre-opt= Pre-operatively, Post-opt= Post-operatively

6.5.2.3.1 Annexin A1: SG

Plasma ANXA1 levels significantly inversely correlated with BMI ($r=-0.389$, $p=0.030$, $n=24$) (Figure 6.10) and significantly directly correlated with plasma adiponectin levels ($r=0.440$, $p=0.020$, $n=22$) (Figure 6.11) in the SG cohort.




Figure 6.10: Correlation between plasma Annexin A1 levels and BMI in Sleeve Gastrectomy participant cohort. Plasma Annexin A1 levels significantly inversely correlated with BMI. One tailed Person correlation coefficient was used to statistically analyse the data.
BMI= Body mass index




Figure 6.11: Correlation between plasma Annexin A1 levels and plasma Adiponectin levels in Sleeve Gastrectomy participant cohort. Plasma Adiponectin levels significantly directly correlated with BMI. One tailed Person correlation coefficient was used to statistically analyse the data.

6.5.2.3.2 Annexin A1: RYGB

Plasma ANXA1 levels significantly inversely correlated with plasma triglyceride levels ($r=-0.421$, $p=0.032$, $n=20$) (Figure 6.12), directly correlated with plasma HDL levels ($r=0.364$, $p=0.058$, $n=20$) (Figure 6.13), significantly directly correlated with plasma CRP levels ($r=0.518$, $p=0.020$, $n=16$) (Figure 6.14), significantly directly with plasma IL-1 β levels ($r=0.464$, $p=0.026$, $n=18$) (Figure 6.15), directly with plasma TNF α levels ($r=0.363$, $p=0.053$, $n=21$) (Figure 6.16) and significantly directly with plasma GLP-1 levels ($r=0.689$, $p=0.002$, $n=16$) (Figure 6.17) in the RYGB cohort.




Figure 6.12: Correlation between plasma Annexin A1 levels and plasma Triglyceride levels in Roux-en-Y Gastric Bypass participant cohort. Plasma Annexin A1 levels significantly inversely correlated with plasma triglyceride levels. One tailed Person correlation coefficient was used to statistically analyse the




Figure 6.13: Correlation between plasma Annexin A1 levels and plasma HDL levels in Roux-en-Y Gastric Bypass participant cohort. Plasma Annexin A1 levels directly correlated with plasma HDL levels. One tailed Person correlation coefficient was used to statistically analyse the data.
HDL= High density lipoprotein




Figure 6.14: Correlation between plasma Annexin A1 levels and plasma CRP levels in Roux-en-Y Gastric Bypass participant cohort. Plasma Annexin A1 levels significantly directly correlated with plasma CRP levels. One tailed Person correlation coefficient was used to statistically analyse the data.
CRP= C-reactive peptide




Figure 6.15: Correlation between plasma Annexin A1 levels and plasma IL-1 β levels in Roux-en-Y Gastric Bypass participant cohort. Plasma Annexin A1 levels significantly directly correlated with plasma IL-1 β levels. One tailed Person correlation coefficient was used to statistically analyse the data. IL-1 β = interleukin-1 β




Figure 6.16: Correlation between plasma Annexin A1 levels and plasma TNF α levels in Roux-en-Y Gastric Bypass participant cohort. Plasma Annexin A1 levels directly correlated with plasma TNFα levels. One tailed Person correlation coefficient was used to statistically analyse the data.
TNFα= Tumour necrosis factor α




Figure 6.17: Correlation between plasma Annexin A1 levels and plasma GLP-1 levels in Roux-en-Y Gastric Bypass participant cohort. Plasma Annexin A1 levels significantly directly correlated with plasma GLP-1 levels. One tailed Person correlation coefficient was used to statistically analyse the data. GLP-1= Glucagon like peptide- 1

6.5.3 Pre and post bariatric surgery: Type 2 diabetes mellitus Status

Participant Characteristics: Type 2 diabetes mellitus status							
		Pre-Opt		6 Weeks Post-Opt		3 Months Post-Opt	
		Non-T2DM	T2DM	Non-T2DM	T2DM	Non-T2DM	T2DM
Number of participants		16	10	9	7	7	4
Gender							
	Males	3	1	0	0	0	0
	Females	13	9	9	7	7	4
Age, years							
		50±2.4 (n=16)	45±2.9 (n=10)	48±3.7 (n=9)	43±3.5 (n=7)	53±2.8 (n=7)	47±5.6 (n=4)
Type of Surgery							
	Sleeve Gastrectomy	8	5	4	3	3	2
	Roux-en-Y Gastric Bypass	8	5	5	4	4	2
Anthropometric Measures							
	Weight (kg)	125.1±9.4 (n=16)	141.7±6.3 (n=10)	115.4±6.2 (n=9)	131.7±9.3 (n=6)	110.9±6.0 (n=5)	99.65±9.1** (n=4)
	BMI (kg/m ²)	48.5±1.4 (n=16)	51.6±2.0 (n=10)	42.1±1.6* (n=9)	48.5±2.5 (n=6)	39.8±1.9** (n=5)	37.6±2.6***/## (n=4)
Lipid Profiles							
	Triglycerides (mmol/L)	1.3±0.1 (n=15)	1.3±0.1 (n=10)	1.4±0.1 (n=9)	1.6±0.1 (n=7)	1.3±0.2 (n=5)	2.6±0.4 (n=4)
	Total Cholesterol (mmol/L)	4.6±1.2 (n=15)	4.3±0.2 (n=10)	4.6±0.4 (n=9)	4.9±0.3 (n=7)	4.3±0.3 (n=5)	4.8±0.7 (n=4)
	HDL (mmol/L)	1.1±0.1 (n=15)	1.2±0.1 (n=10)	1.1±0.1 (n=9)	1.1±0.1 (n=7)	1.3±0.1 (n=5)	1.4±0.1 (n=4)
	LDL (mmol/L)	2.8±0.1 (n=15)	2.5±0.2 (n=10)	2.8±0.3 (n=9)	3.1±0.2 (n=7)	2.4±0.2 (n=5)	1.6±0.6 (n=4)

Table 6.5: Participant characteristics of non-T2DM and T2DM cohort pre- and post- Bariatric Surgery. The anthropometric measures and lipid profiles were analysed in participants undergoing bariatric surgery at pre-, 6 weeks and 3 months post-operatively and compared between surgery types. Two-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *vs pre opt of the corresponding group. Pre-opt= Pre-operatively, Post-opt= Post-operatively, T2DM= Type 2 diabetes mellitus, BMI= Body mass index, HDL= High density lipoprotein and LDL= Low density lipoprotein

The characteristics of the non-T2DM and T2DM cohorts are presented in table 6.5. Of the 16 non-T2DM participants at pre-opt, 3 were male and 13 were female, with an average age of 50 ± 2.4 . At 6 weeks, 9 females (average age 48 ± 3.7) and at 3 months, 7 females (average age 53 ± 2.8) continued their participation. Similarly, of the 10 T2DM participants at pre-opt, 1 was male and 9 were females, with an average age of 45 ± 2.9 . At 6 weeks, 7 females (average age 43 ± 3.5) and at 3 months, 4 females (average age 47 ± 5.6) continued their participation.

6.5 .3.1 Anthropometric and plasma lipid levels

The effect of time ($p=0.060$), T2DM status ($p=0.400$) nor their interaction ($p=0.391$) was statistically significant on the average weight. The effect of time ($p=0.0001$) (but not T2DM status ($p=0.167$) nor it's interaction with time ($p=0.196$)) was statistically significant on the average change in BMI. Therefore, no significant differences in the average weight and BMI were observed between the two cohorts at any timepoint.

However, the average weight decreased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average weight was reduced at 6 weeks and 3 months post-opt compared to pre-op, however, the difference observed was not statistically significant (115.4 ± 6.2 kg, $n=9$ and 110.9 ± 6.0 kg, $n=5$ vs 125.1 ± 9.4 kg, $n=16$, respectively) (Table 6.5). Similarly, in the T2DM cohort, the average weight was reduced at 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (99.7 ± 9.1 kg, $n=4$ vs 141.7 ± 6.3 kg, $n=10$, $p=0.008$, respectively) (Table 6.5). Furthermore, the average BMI was reduced in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average BMI was significantly reduced at 6 weeks and 3 months post-opt compared to pre-opt, (42.1 ± 1.6 kg/m², $n=9$, $p=0.029$ and 39.8 ± 1.9

kg/m², n=5, p=0.009 vs 48.5±1.3 kg/m², n=16, respectively), as shown in Figure 6.18. Similarly in the T2DM cohort, the average BMI was reduced at 3 months post-opt compared to pre-opt (37.6±2.6 kg/m², n=4 vs 51.6±2.0 kg/m², n=10, p=0.0003, respectively) and to 6 weeks post-opt (48.5±2.5 kg/m², n=6 vs 37.6±2.6 kg/m², n=4, p=0.009, respectively), as shown in Figure 6.18.

The effect of time, T2DM status nor their interaction was significantly significant on the average plasma triglyceride levels (p=0.323, p=0.235 and p=0.688, respectively), plasma total cholesterol levels (p=0.447, p=0.464 and p=0.339, respectively), plasma HDL levels (p=0.181, p=0.537 and p=0.701, respectively) and plasma LDL levels (p=0.248, p=0.857 and p=0.242, respectively), therefore, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point.




Figure 6.18: Average BMI (kg/m²) of non-T2DM and T2DM cohort pre- and post- Bariatric surgery. The average BMI was significantly reduced at 6 weeks and 3 months post-opt compared to pre-opt in the non-T2DM cohort ($p=0.029$, $n=9$ and $p=0.009$, $n=5$, respectively). Similarly, the average BMI was reduced at 3 months post-opt compared to pre-opt ($p=0.0003$, $n=10$) and to 6 weeks post opt ($p=0.009$, $n=4$) in T2DM cohort. However, no statistically significant differences were observed between the cohorts at any time point. Two-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM.

P value was set at $p < 0.05$ and is denoted by *vs pre opt and #vs 6 weeks post-opt of the corresponding group BMI= Body mass index, T2DM= Type 2 diabetes mellitus, Pre-opt= Pre-operatively, Post-opt= Post-operatively

6.5.3.2 Plasma inflammatory biomarkers, markers of insulin sensitivity and gut hormones

Table 6.6 presents the analysis of plasma inflammatory biomarkers, plasma markers of insulin sensitivity and plasma gut hormones in non-T2DM and T2DM cohorts pre and post- bariatric surgery.

6.5.3.2.1 Plasma inflammatory biomarkers

The effect of time ($p=0.657$), T2DM status ($p=0.540$) nor their interaction ($p=0.706$) was statistically significant on the average plasma CRP levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average plasma CRP levels decreased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma CRP levels decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant ($8.8 \pm 2.8 \mu\text{g/mL}$, $n=7$ and $5.4 \pm 1.9 \mu\text{g/mL}$, $n=4$ vs $12.1 \pm 2.0 \mu\text{g/mL}$, $n=12$, respectively). Similarly in the T2DM cohort, the average plasma CRP levels decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant ($11.3 \pm 3.8 \mu\text{g/mL}$, $n=7$ and $10.8 \pm 7.3 \mu\text{g/mL}$, $n=3$ vs $10.6 \pm 6.3 \mu\text{g/mL}$, $n=6$, respectively).

The effect of time ($p=0.044$) but not T2DM status ($p=0.806$) nor its interaction with time ($p=0.732$) was statistically significant on the average plasma IL-1 β levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. However, the average plasma IL-1 β levels increased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma IL-1 β levels increased at 3 months post-opt compared to pre-opt, however, the difference observed

was not statistically significant (2.3 ± 0.4 pg/mL, $n=5$ vs 1.2 ± 0.2 pg/mL, $n=12$, respectively). Similarly in the T2DM cohort, the average plasma IL-1 β levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (1.5 ± 0.3 pg/mL, $n=4$ and 2.1 ± 0.4 pg/mL, $n=4$ vs 1.5 ± 0.5 pg/mL, $n=5$, respectively).

The effect of time ($p=0.001$) but not T2DM status ($p=0.459$) nor its interaction with time ($p=0.544$) was statistically significant on the average plasma IL-10 levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. However, the average plasma IL-10 levels increased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma IL-10 levels significantly increased at 6 weeks and 3 months post-opt compared to pre-opt (10.0 ± 1.2 pg/mL, $n=8$, $p=0.021$ and 9.3 ± 0.9 pg/mL, $n=4$ vs 7.2 ± 1.0 pg/mL, $n=12$, respectively). Similarly in the T2DM cohort, the average plasma IL-10 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (9.6 ± 1.0 pg/mL, $n=6$ and 8.8 ± 1.4 pg/mL, $n=4$ vs 6.4 ± 0.8 pg/mL, $n=6$, respectively).

The effect of time ($p=0.789$), T2DM status ($p=0.915$) nor their interaction ($p=0.699$) was statistically significant on the average plasma TNF α levels, therefore, no significant differences were observed between the two cohorts at any time point. The average plasma TNF α levels remained similar in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma TNF α levels remained similar at 6 weeks and 3 months post-opt (21.5 ± 1.1 pg/mL, $n=9$ and 21.2 ± 1.5 pg/mL, $n=5$ vs 21.7 ± 0.4 pg/mL, $n=14$, respectively). Similarly, in the T2DM cohort, the average plasma TNF α levels remained

similar at 6 weeks and 3 months post-opt compared to pre-opt (20.6 ± 1.4 pg/mL, n=6 and 22.4 ± 2.6 pg/mL, n=4 vs 21.6 ± 1.1 pg/mL, n=5, respectively).

The effect of time ($p=0.0002$) but not T2DM status ($p=0.374$) nor its interaction with time ($p=0.699$) was statistically significant on the average plasma leptin levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. However, the average plasma leptin levels decreased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma leptin levels significantly decreased at 6 weeks and 3 months post-opt (35.1 ± 7.5 ng/mL, n=9, $p=0.02$ and 42.4 ± 5.9 ng/mL, n=5, vs 83.5 ± 9.3 ng/mL, n=14, respectively). Similarly in the T2DM cohort, the average plasma leptin levels significantly decreased at 6 weeks and 3 months post-opt compared to pre-opt (36.9 ± 6.1 ng/mL, n=7, $p<0.05$ and 27.3 ± 6.4 ng/mL, n=4 vs 75.0 ± 7.1 ng/mL, n=6, respectively).

The effect of time ($p=0.015$) but not T2DM status ($p=0.919$) nor its interaction with time ($p=0.531$) was statistically significant on the average plasma adiponectin levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. However, the average plasma adiponectin levels increased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma adiponectin levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (8.6 ± 1.2 μ g/mL, n=9 and 8.1 ± 1.3 μ g/mL, n=4 vs 6.1 ± 0.7 μ g/mL, n=12, respectively). Similarly in the T2DM cohort, the average plasma adiponectin levels increased at 6 weeks compared to pre-opt, however, the difference observed was not statistically significant (8.4 ± 1.3 μ g/mL, n=5 and 9.7 ± 1.0 μ g/mL, n=4 vs 4.7 ± 1.0 μ g/mL, n=6, respectively).

6.5.3.2.2 Plasma biomarkers of insulin sensitivity

The effect of time ($p=0.783$), T2DM status ($p=0.097$) nor their interaction ($p=0.622$) was significantly significant on the average fasting plasma glucose levels, therefore, no significant differences were observed between the two cohorts at any timepoint. The average fasting plasma glucose levels remained similar in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average fasting plasma glucose levels were not significantly altered at 6 weeks and 3 months post-opt compared to pre-opt (3.8 ± 0.2 mmol/L, $n=6$ and 3.9 ± 0.2 mmol/L, $n=4$ vs 3.6 ± 0.3 mmol/L, $n=13$, respectively). Similarly in the T2DM cohort, the average fasting plasma glucose levels were not significantly altered at 6 weeks and 3 months post-opt compared to pre-opt (4.2 ± 0.7 mmol/L, $n=3$ and 4.5 ± 0.7 mmol/L, $n=3$ vs 4.9 ± 0.8 mmol/L, $n=6$, respectively).

The effect of time ($p=0.011$) (but not T2DM status ($p=0.859$)) and its interaction with T2DM status ($p=0.002$) were statistically significant on the average fasting plasma insulin levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. The average fasting plasma insulin levels decreased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average fasting plasma insulin significantly levels decreased at 6 weeks and 3 months post-opt compared to pre-opt (17.0 ± 2.5 μ IU/mL, $n=6$, $p=0.038$ and 18.0 ± 3.2 μ IU/mL, $n=4$ vs 23.9 ± 1.9 μ IU/L, $n=13$, respectively). Similarly in the T2DM cohort, the average fasting plasma insulin levels decreased 6 weeks and at 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (18.7 ± 2.5 μ IU/mL, $n=4$ and 10.3 ± 2.2 μ IU/mL, $n=3$ vs 29.7 ± 4.5 μ IU/mL, $n=6$, $p=0.022$, respectively).

The effect of time ($p=0.027$) (but not T2DM status ($p=0.925$)) and its interaction with T2DM status ($p=0.004$) were statistically significant on the average HOMA-IR, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. The average HOMA-IR was reduced in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average HOMA-IR reduced at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (2.0 ± 0.3 , $n=6$ and 2.1 ± 0.4 , $n=4$ vs 2.9 ± 0.3 , $n=9$, respectively). Similarly in the T2DM cohort, the average HOMA-IR decreased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (2.5 ± 0.3 , $n=4$ and 1.3 ± 0.2 , $n=3$ vs 3.6 ± 0.5 , $n=6$, respectively).

The effect of time ($p=0.013$) (but not T2DM status ($p=0.399$)) and its interaction with T2DM status ($p=0.026$) were statistically significant on the average HOMA-%S, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. The average HOMA-%S was improved in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average HOMA-%S improved at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (57.9 ± 12.0 %, $n=6$ and 52.0 ± 10.5 %, $n=4$ vs 36.2 ± 3.1 , $n=9$, respectively). Similarly in the T2DM cohort, the average HOMA-%S was improved at 3 months post opt compared to pre-opt and 6 weeks post opt, however, the difference observed was not statistically significant (41.8 ± 5.6 %, $n=3$ and 93.4 ± 25.2 %, 30.7 ± 4.1 %, $n=6$, respectively).

The effect of time ($p=0.198$), T2DM status ($p=0.420$) nor their interaction ($p=0.305$) was statistically significant on the average HOMA-% β , therefore, no significant differences were observed between the two cohorts at any timepoint. The average HOMA-% β

remained similar in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average HOMA-% β remained similar at 6 weeks and 3 months post-opt compared to pre-opt (264.3 ± 12.4 %, n=6 and 268.1 ± 33.0 %, n=4 vs 296.4 ± 24.6 %, n=9, respectively). Whereas, in the T2DM cohort, the average HOMA-% β reduced at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (252.6 ± 23.0 %, n=3 and 167.4 ± 55.0 %, n=3 vs 314.7 ± 70.6 %, n=6, respectively).

6.5.3.2.3 Plasma gut hormone

The effect of time (p=0.001) (but not T2DM status (p=0.569) nor its interaction with time (p=0.834)) was statistically significant on the average plasma GLP-1 levels, however, the post-hoc analysis did not detect specific significant differences between the cohorts at any time point. The average plasma GLP-1 levels increased in both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma GLP-1 levels increased at 6 weeks post-opt (313.7 ± 15.3 ng/mL vs 257.7 ± 14.8 ng/mL, n=13, respectively) and significantly at 3 months post-opt compared to pre-opt (339.2 ± 25.7 ng/mL, n=4 vs 257.7 ± 14.8 ng/mL, n=13, p=0.029, respectively). Similarly, in the T2DM cohort, the average plasma GLP-1 levels increased at 6 weeks (306.3 ± 43.7 ng/mL, n=3 vs 278.1 ± 21.1 ng/mL, n=6, respectively) and significantly at 3 months post-opt compared to pre-opt (374.8 ± 21.9 ng/mL, n=3 and vs 278.1 ± 21.1 ng/mL, n=6, p=0.038, respectively).

Plasma Biomarkers: Type 2 diabetes mellitus Status							
		Pre-opt		6 Weeks Post-opt		3 Months Post-opt	
Type 2 diabetes mellitus Status		Non-T2DM	T2DM	Non-T2DM	T2DM	Non-T2DM	T2DM
Inflammatory Biomarkers							
	CRP ($\mu\text{g/mL}$)	12.1 \pm 2.0 (n=12)	10.6 \pm 6.3 (n=6)	8.8 \pm 2.8 (n=7)	11.3 \pm 3.8 (n=7)	5.4 \pm 1.9 (n=4)	10.8 \pm 7.3 (n=3)
	Annexin A1 (ng/mL)	1.2 \pm 0.1 (n=14)	1.1 \pm 0.1 (n=6)	1.3 \pm 0.1 (n=9)	1.6 \pm 0.3 (n=7)	1.6 \pm 0.2 (n=5)	1.6 \pm 0.2 (n=4)
	IL-1 β (pg/mL)	1.2 \pm 0.2 (n=12)	1.5 \pm 0.5 (n=5)	1.3 \pm 0.2 (n=9)	1.5 \pm 0.3 (n=4)	2.3 \pm 0.4 (n=5)	2.1 \pm 0.4 (n=4)
	IL-10 (pg/mL)	7.2 \pm 1.0 (n=12)	6.4 \pm 0.8 (n=6)	10.0 \pm 1.2* (n=8)	9.6 \pm 1.0 (n=6)	9.3 \pm 0.9 (n=4)	8.8 \pm 1.4 (n=4)
	TNF α (pg/mL)	21.7 \pm 0.4 (n=14)	21.6 \pm 1.1 (n=5)	21.5 \pm 1.1 (n=9)	20.6 \pm 1.4 (n=6)	21.1 \pm 1.5 (n=5)	22.4 \pm 2.6 (n=4)
	Leptin (ng/mL)	83.5 \pm 9.3 (n=14)	75.0 \pm 7.1 (n=6)	35.1 \pm 7.5* (n=9)	36.9 \pm 6.1* (n=7)	42.4 \pm 5.9 (n=5)	27.3 \pm 6.4 (n=4)
	Adiponectin ($\mu\text{g/mL}$)	6.1 \pm 0.7 (n=12)	4.7 \pm 1.0 (n=6)	8.6 \pm 1.2 (n=9)	8.4 \pm 1.3 (n=5)	8.1 \pm 1.3 (n=4)	9.7 \pm 1.0 (n=4)
Insulin Sensitivity							
	Fasting Glucose (mmol/L)	3.6 \pm 0.3 (n=13)	4.9 \pm 0.8 (n=6)	3.8 \pm 0.2 (n=6)	4.2 \pm 0.7 (n=3)	3.9 \pm 0.2 (n=4)	4.5 \pm 0.7 (n=3)
	Fasting Insulin ($\mu\text{IU/mL}$)	23.9 \pm 1.9 (n=13)	29.7 \pm 4.5 (n=6)	17.0 \pm 2.5* (n=6)	18.7 \pm 2.5 (n=4)	18.0 \pm 3.2 (n=4)	10.3 \pm 2.2 (n=3)
	HOMA-IR	2.9 \pm 0.3 (n=9)	3.6 \pm 0.5 (n=6)	2.0 \pm 0.3 (n=6)	2.5 \pm 0.3 (n=4)	2.1 \pm 0.4 (n=4)	1.3 \pm 0.2 (n=3)
	HOMA-%S (%)	36.2 \pm 3.1 (n=9)	41.8 \pm 5.6 (n=3)	57.9 \pm 12.0 (n=6)	93.4 \pm 25.2 (n=3)	52.0 \pm 10.5 (n=4)	30.7 \pm 4.1 (n=6)
	HOMA-% β (%)	296.4 \pm 24.6 (n=9)	314.7 \pm 70.6 (n=6)	264.3 \pm 12.4 (n=6)	252.6 \pm 23.0 (n=3)	296.4 \pm 24.6 (n=9)	167.4 \pm 55.0 (n=3)
Gut Hormone							
	GLP-1 (ng/mL)	257.7 \pm 14.8 (n=13)	278.1 \pm 21.1 (n=6)	313.7 \pm 15.3 (n=6)	306.3 \pm 43.7 (n=3)	339.2 \pm 25.7* (n=4)	374.8 \pm 21.9* (n=3)

Table 6.6: Plasma biomarkers analysis of non-T2DM and T2DM cohorts at pre- and post- Bariatric surgery. Plasma inflammatory biomarkers, fasting glucose and plasma insulin levels and plasma gut hormone were analysed and HOMA-IR, HOMA-%S and HOMA-% β were calculated using the equations presented in section 6.4.2.2. One-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM.

P value was set at $p < 0.05$ and is denoted by *vs pre opt

Pre-opt= Pre-operatively, Post-opt= Post-operatively, T2DM= Type 2 diabetes mellitus, CRP= C-reactive protein, IL= Interleukin, TNF α = Tumour necrosis factor α , HOMA-IR= Homeostatic model assessment of insulin resistance, HOMA-%S= Homeostatic model assessment of insulin sensitivity, HOMA-% β = Homeostatic model assessment of pancreatic β cell function and GLP-1= Glucagon like peptide 1

6.5.3.3 Annexin A1

The effect of time ($p=0.080$), T2DM status ($p=0.679$) nor their interaction ($p=0.505$) was statistically significant on the average plasma ANXA1 levels, therefore, no significant differences were observed between the two cohorts at any timepoint. However, the average plasma ANXA1 increased both cohorts at post-opt compared to pre-opt. In the non-T2DM cohort, the average plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (1.3 ± 0.1 ng/mL, $n=9$ and 1.6 ± 0.2 ng/mL, $n=5$ vs 1.2 ± 0.1 ng/mL, $n=14$, respectively), as shown in Figure 6.19. Similarly in the T2DM cohort, the average plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt, however, the difference observed was not statistically significant (1.6 ± 0.3 ng/mL, $n=7$ and 1.6 ± 0.2 ng/mL, $n=4$ vs 1.1 ± 0.1 ng/mL, $n=6$, respectively), as shown in Figure 6.19.




Figure 6.19: Average plasma Annexin A1 (ng/mL) levels of non-T2DM and T2DM cohort pre- and post-Bariatric surgery. The average plasma Annexin A1 levels were increased at 6 weeks and 3 months post-opt compared to pre-opt in the non-T2DM and T2DM cohorts. However, no statistically significant differences were observed between the cohorts at any time point. Two-way ANOVA followed by Tukey's Post-hoc multiple comparison test was used to statistically analyse the data. Data presented as Mean \pm SEM. P value was set at $p < 0.05$.

T2DM= Type 2 diabetes mellitus, Pre-opt= Pre-operatively, Post-opt= Post-operatively

6.5.3.3.1 Annexin A1: Non-Type 2 diabetes mellitus

Plasma ANXA1 levels directly correlated with plasma HDL levels ($r=0.323$, $p=0.058$, $n=25$) (Figure 6.20), significantly inversely correlated with plasma LDL levels ($r=-0.340$, $p=0.048$, $n=25$) (Figure 6.21) and directly correlated with HOMA- β ($r=0.343$, $p=0.074$, $n=19$) (Figure 6.22) in the non-T2DM cohort.




Figure 6.20: Correlation between plasma Annexin A1 levels and plasma HDL in non-T2DM participant cohort. Plasma Annexin A1 levels directly correlated with plasma HDL levels. One tailed Person correlation coefficient was used to statistically analyse the data.
HDL= High density lipoprotein




Figure 6.21: Correlation between plasma Annexin A1 levels and plasma LDL in non-T2DM participant cohort. Plasma Annexin A1 levels significantly inversely correlated with plasma LDL levels. One tailed Person correlation coefficient was used to statistically analyse the data.
LDL= Low density lipoprotein




Figure 6.22: Correlation between plasma Annexin A1 levels and HOMA-%β in non-T2DM participant cohort. Plasma Annexin A1 levels directly correlated with HOMA-%β. One tailed Person correlation coefficient was used to statistically analyse the data.
HOMA-%β= Homeostatic model assessment of pancreatic β cell function

6.5.3.3.2 Annexin A1: Type 2 diabetes mellitus

Plasma ANXA1 levels inversely correlated with BMI ($r=-0.377$, $p=0.075$, $n=16$) (Figure 6.23), directly correlated with plasma TNF α levels ($r=0.419$, $p=0.053$, $n=16$) (Figure 6.24) and directly correlated with HOMA-%S ($r=0.429$, $p=0.082$, $n=12$) (Figure 6.25) in the T2DM cohort.




Figure 6.23: Correlation between plasma Annexin A1 levels and BMI in T2DM participant cohort.
Plasma Annexin A1 levels significantly inversely correlated with BMI. One tailed Person correlation coefficient was used to statistically analyse the data.
BMI= Body mass index




Figure 6.24: Correlation between plasma Annexin A1 levels and plasma TNF α in T2DM participant cohort. Plasma Annexin A1 levels directly correlated with plasma TNF α levels. One tailed Person correlation coefficient was used to statistically analyse the data.
TNF α = Tumour necrosis factor α




Figure 6.25: Correlation between plasma Annexin A1 levels and HOMA-%S in T2DM participant cohort.
Plasma Annexin A1 levels directly correlated with HOMA-%S. One tailed Person correlation coefficient was used to statistically analyse the data.
HOMA-%S= Homeostatic model assessment of insulin sensitivity

6.6 Discussion

In the current study, the plasma ANXA1 levels were analysed in bariatric surgery patients and correlated with plasma inflammatory biomarkers, markers of insulin sensitivity and gut hormones. Interestingly, plasma ANXA1 levels increased at 6 weeks and 3 months post-opt compared to pre-opt in all studied cohorts, however, a statistical power could not be reached, possibly owing to small participant number as the retention rate throughout the study was relatively poor. Interestingly, no significant difference was observed between the average BMI of SG and RYGB cohort, yet the average plasma ANXA1 levels were higher in the SG cohort at 6 weeks and significantly inversely correlated with BMI in the SG cohort only, perhaps reflecting the less invasive nature of SG and inducing a quicker anti-inflammatory/pro-resolving response compared to RYGB (Frikke-Schmidt, O'Rourke et al. 2016). Interestingly, Pietrani 2018 report higher circulating levels of ANXA1 in obese T2DM patients compared to lean or overweight T2DM patients and a direct correlation with BMI. Whereas, in the current study, the average plasma ANXA1 levels increased in the T2DM cohort at 6 weeks post- compared to pre-opt and significantly inversely correlated with BMI. Furthermore, plasma ANXA1 levels exhibited strong correlations with BMI, plasma lipid levels, plasma CRP, IL-10, IL-1 β and TNF α levels, HOMA-%S, HOMA-% β and plasma GLP-1 levels. The correlations between plasma ANXA1 and anthropometric measures, plasma lipids and plasma biomarkers analysed differ between surgery type and T2DM status, suggesting ANXA1 may have different effects in different populations, i.e. non-T2DM vs T2DM.

Approximately, 60-70% of obese individuals have abnormal lipid profiles (dyslipidaemia), including increased circulating LDL levels, increased circulating triglycerides and decreased circulating HDL, thereby, significantly increasing the risk of

cardiovascular diseases in these individuals (Bays, Toth et al. 2013). Bariatric surgery induced weight loss is allied with reduced circulating LDL levels, reduced triglycerides levels and increased circulating HDL levels, thereby significantly reversing dyslipidaemia (Gloy, Briel et al. 2013, Dash, Xiao et al. 2016, Heffron, Parikh et al. 2016).

In the current study, the remission of dyslipidaemia was greater in RYGB cohort compared to the SG cohort and this observation is supported by previous literature (Heffron, Parikh et al. 2016). Furthermore, the results of the current study are the first to show a direct correlation between plasma ANXA1 and plasma HDL levels and inverse correlation between plasma ANXA1 and LDL levels in bariatric surgery patients. Both ANXA1 and HDL mediate cardioprotective effects due to their anti-oxidative, anti-inflammatory and anti-atherogenic properties (Pan, Kong et al. 2016). Increased expression levels of ANXA1 were observed in atherosclerotic plaques in carotid stenosis patients without clinical symptoms compared to those with significant cerebrovascular symptoms, suggesting ANXA1 may be involved in resolving and/or attenuating the progression of plaque formation (Cheuk, Cheng 2011). Furthermore, deletion of FPR2/ALX or ANXA1 in Apolipoprotein E (*APO*^{-/-}) high fat diet (HFD) fed mice enhances atherosclerotic lesion formation, adhesion, recruitment and accumulation of arterial myeloid cells, whereas, early atherogenesis and plaque formation is reduced upon treatment with AC2-26 (Drechsler, de Jong et al. 2015). However, FPR2/ALX expression levels are upregulated in human carotid atherosclerotic lesions compared to healthy vessels and is suggested to promote disease progression and increase plaque stability (Petri, Laguna-Fernandez et al. 2015). A recent study reported a novel anti-inflammatory mechanism of HDL through modulating the expression of ANXA1 in vascular

endothelial cells. HDL induced the expression of ANXA1 in endothelial cells in a dose dependent manner and this increase was associated with inhibition of cell surface adhesion molecules and secretion of chemotactic factors (vascular cell adhesion molecule 1, intracellular adhesion molecule 1, E-selection and monocyte chemoattractant protein 1), and inhibiting monocyte adhesion, thereby attenuating formation of atherosclerotic lesions (Pan, Kong et al. 2016). Interestingly, low HDL levels are associated with increased risk of developing T2DM, however, in the current study no significant differences in the plasma HDL levels between T2DM and non-T2DM were observed (Chahil, Ginsberg 2006). Interestingly, the anti-oxidative, anti-inflammatory and anti-atherogenic properties of HDL are suggested to be compromised in diabetes mellitus patients, due to oxidative modification and glycation of HDL proteins (Femlak, Gluba-Brzãzka et al. 2017). In the current study, plasma HDL levels directly correlated with plasma ANXA1 in the non-T2DM cohort but not in the T2DM cohort, possibly reflecting its compromised function in T2DM patients.

Obesity associated inflammation is initiated within the adipose tissue and liver, with elevated macrophage infiltration and expression of proinflammatory cytokines and adipokines. These pro-inflammatory cytokines and adipokines enter the circulation and cause systemic inflammation and contribute to the onset of obesity associated co-morbidities such as, CVD and T2DM (Wood, de Heredia et al. 2009). Changes in cytokine and adipokine synthesis and secretion have been suggested to improve metabolic risk factors in obesity in response to weight loss induced by lifestyle changes and surgical procedures, however, the results reported are inconsistent. In the current study, plasma levels of pro-inflammatory cytokines showed an inconsistent trend suggesting that the circulatory levels all pro-inflammatory cytokines are not regulated in a similar manner

and the invasive nature of surgery may contribute to the discrepancies (Rakotoarivelo, Lacraz et al. 2018).

In the current study, plasma CRP levels were reduced, whereas, plasma IL-1 β were increased post-surgery in all studied cohorts. CRP is an acute phase protein involved in regulating the innate immune system and is used as a marker to identify inflammatory diseases as it exacerbates the inflammatory status and induces systemic inflammation (Wang, Nakayama 2010). Circulating levels of CRP are higher in overweight and obese individuals compared to healthy individuals and are associated with distribution of body fat independent of BMI (Visser, Bouter et al. 1999). Elevated CRP levels are an independent risk factor for CVD and insulin resistance and have been suggested to reflect the amount and activity of other pro-inflammatory, pro-atherogenic cytokines such as, TNF α , interleukin 1 (IL-1) and interleukin-6 (Lagrand, Visser et al. 1999, Yudkin, Stehouwer et al. 1999, Tchernof, Nolan et al. 2002). Therefore, the decrease in plasma CRP following bariatric surgery reflects an improved inflammatory status and reduced risk of developing co-morbidities (Rojano-Rodriguez, Valenzuela-Salazar et al. 2014, Netto, Bettini et al. 2015). IL-1 β is a pleiotropic cytokine and is involved in regulating pro-inflammatory processes that facilitate monocyte infiltration and migration to the site of inflammation, insulin secretion and appetite regulation (Sica, Wang et al. 1990, Kupper, Groves 1995, Mandrup-Poulsen 1996, Plata-Salaman 1994). Though IL-1 β is considered as a pro-inflammatory cytokine, its involvement in regulating gastric acid secretion and emptying, adipocyte development and adipose biology may induce an increase post bariatric surgery (Delikat, Harris et al. 1993, El-Omar 2001). It stimulates gastrin release from enteroendocrine G cells in the gastric antrum and duodenum (Weigert, Schaffer et al. 1996). Gastrin is released in response to food and gastric

distension, acts to increase the secretion of hydrochloric acid, pepsinogen and pancreatic juices and reduces appetite (Meek, Lewis et al. 2016). Circulating levels of gastrin are reduced 12 months post RYGB, whereas, SG has been associated with increased plasma gastrin levels in both human and animal studies, but the exact mechanism that induces this change is unclear (Sundbom, Holdstock et al. 2007, Sillakivi, Suumann et al. 2013, Grong, Arbo et al. 2015). In the current study, plasma IL-1 β levels increase in both RYGB and SG cohorts, therefore, the increase in IL-1 β may act to increase gastrin levels, reduce appetite and aid in weight loss. The previous literature reports conflicting results regarding the effect of bariatric surgery induced weight loss on circulating TNF α levels (Hafida, Mirshahi et al. 2016). Some authors report no change in circulating TNF α levels following RYGB, whereas, others report reduced plasma TNF α following SG and RYGB (Catalan, Gomez-Ambrosi et al. 2007, Sams, Blackledge et al. 2016, Viana, Araujo-Dasilio et al. 2013, Netto, Bettini et al. 2015). In the current study, no significant changes in plasma TNF α levels were observed post- compared to pre-opt. Furthermore, in the current study, direct correlations between plasma ANXA1 levels and CRP, TNF α and IL-1 β were observed, whereas, an inverse correlation between plasma ANXA1 and CRP in obesity has been reported (Kosicka, Cunliffe et al. 2013), suggesting the direct correlation may act as a compensatory mechanism underlying systemic inflammation and possibly masking the effect of ANXA1. Furthermore, the direct correlation with plasma IL-1 β may suggest a role for ANXA1 in regulating appetite, adipose tissue development and body weight.

IL-10 is a pleiotropic anti-inflammatory cytokine involved in immunoregulation and inflammation. It is synthesised and secreted by macrophages, dendritic cells, B cells, CD4⁺, CD8⁺ and Th2 T cells and inhibits the major histocompatibility complex class II

and costimulatory molecules on monocytes and macrophages, and limits the synthesis and secretion of pro-inflammatory cytokines and chemokines in T-helper 1 T cells and natural killer cells by inhibiting nuclear factor kappa β to aid in resolving inflammation (Couper, Blount et al. 2008). IL-10 is considered to play a protective role in human metabolism, as it is shown to provide protection against endothelial dysfunction (by inhibiting superoxide thereby, suppressing impaired endothelium-dependent relaxation) during diabetes mellitus (Gunneth, Heistad et al. 2002). Plasma IL-10 levels are significantly increased after weight loss induced by lifestyle changes and bariatric surgery, and significantly correlates with plasma adiponectin levels (Jung, Park et al. 2008, Netto, Bettini et al. 2015, Wolf, Wolf et al. 2004). Adiponectin induces monocytes and macrophages towards an anti-inflammatory phenotype by inducing synthesis and secretion of IL-10 in human obesity, thus, the increase in plasma adiponectin levels after bariatric surgery may act as a stimulant for increasing plasma IL-10 levels in these individuals (Wolf, Wolf et al. 2004, Jung, Park et al. 2008). Furthermore, recombinant ANXA1 stimulate IL-10 secretion by macrophages primed with liposaccharide in a dose dependent manner, suggesting ANXA1 may partly mediate its anti-inflammatory actions by modulating the secretion of IL-10 (Ferlazzo, D'Agostino et al. 2003). However, plasma IL-10 and plasma ANXA1 levels increased post- compared to pre-opt in the current study, no correlation was observed, suggesting the increase in plasma IL-10 may be independent of plasma ANXA1. Plasma adiponectin and leptin levels are significantly altered following bariatric surgery and reflect the decrease in adipose tissue mass, increased insulin sensitivity and reduced inflammation. In the current study, plasma adiponectin levels directly correlated with plasma ANXA1 levels reflecting the insulin sensitising role of ANXA1. Furthermore, direct correlation with adiponectin suggests, plasma ANXA1

levels may be regulated by total adiposity. The endocrine function of ANXA1 may be compromised in obesity and may primarily act in an autocrine, paracrine manner in the adipose tissue to resolve inflammation, however, this effect may be reversed in response to weight loss, therefore, aiding in reducing systemic inflammation and the risk of developing obesity associated co-morbidities.

Insulin sensitivity and T2DM status are improved following bariatric surgery independent of weight loss (Batterham, Cummings 2016). This improvement is associated with changes in the hormonal secretion in the gut that have a positive influence on insulin secretion, sensitivity and glucose tolerance (Fried 2013). RYGB is considered to produce a greater reduction in insulin resistance soon after the surgery, possibly due to increased secretion of incretins that enhance insulin sensitivity, however, SG, despite being based on a different technique also improves insulin sensitivity to a similar magnitude (Benaiges, Flores Le-Roux et al. 2013). In the current study, HOMA-IR, HOMA-%S and HOMA-% β in the fasted state were used to assess insulin resistance, insulin sensitivity and insulin secretion from pancreatic β cells, respectively. As expected, in the current study, HOMA-IR decreased and HOMA-%S increased, suggestive of decreased insulin resistance and improved insulin sensitivity post bariatric surgery in all study cohorts (Benaiges, Flores Le-Roux et al. 2013, Ferrannini, Mingrone 2009). Furthermore, a HOMA-% β value close to 100% reflects healthy pancreatic β cell function and a decrease in HOMA-% β index indicates improved pancreatic β function post-opt. Previous literature reports increased circulating levels of ANXA1 in obese T2DM patients compared to healthy weight and overweight T2DM patients as well as non-T2DM controls and no correlation with BMI (Pietrani, Ferreira et al. 2018, Purvis, Collino et al. 2019). However, in the current study, no difference in plasma ANXA1 between obese

T2DM and non-T2DM cohort was observed pre-operatively. Furthermore, plasma ANXA1 levels increased following surgery in T2DM and non-T2DM participants in a similar manner and presented an inverse correlation with BMI in the T2DM cohort only. Evidence suggests ANXA1 may be important for regulating blood glucose levels in a diseased state (Purvis, Solito et al. 2019). ANXA1^{-/-} mice HFD fed develop a more severe diabetic phenotype, characterised by increased blood glucose levels, elevated insulin levels and augmentation in oral glucose tolerance test (Purvis, Collino et al. 2019). Furthermore, ANXA1 is suggested to be a key regulator of RhoA activity which restores insulin receptor substrate 1 signalling and recombinant human ANXA1 is proposed as a novel candidate for treatment of T2DM (Purvis, Collino et al. 2019). Therefore, the direct correlation between plasma ANXA1 and HOMA-%S in the T2DM observed in the current study may indicate a role of ANXA1 in improving the glycaemic index and T2DM status of obese T2DM patients following bariatric surgery. Furthermore, ANXA1 is reported to stimulate insulin secretion from rat pancreatic and MIN6N8s cells (Ohnishi, Tokuda et al. 1995) and in the current study a direct correlation between plasma ANXA1 levels and HOMA-% β was observed in the non-T2DM cohort. However, HOMA-% β is not considered a suitable indicator for evaluating improvements in insulin secretion due to the changes in hormones with incretin effect post-surgery as these hormones will only influence insulin secretion postprandially, therefore, HOMA-% β may not truly reflect the efficiency of insulin secretion (Benaiges, Flores Le-Roux et al. 2013). Thus, in addition to fasted state, it would be useful to correlate plasma ANXA1 and markers of insulin resistance, sensitivity and secretion postprandially to identify discrepancies.

The rapid metabolic effects of bariatric surgery have been linked to the changes in the synthesis and secretion of pancreatic and gut hormones. Several gut hormones are

involved in regulating appetite, satiety and gastric emptying and motility (Meek, Lewis et al. 2016). PYY and GLP-1 are secreted by the L cells in the gut and reduce gastric emptying and motility thereby, promoting satiety. Reduced gastric emptying is associated with improved glucose tolerance as it results in a more gradual supply of nutrients to the gut (Schirra, Katschinski et al. 1996). The increase in postprandial PYY and GLP-1 levels post bariatric surgery are unclear, however, the ‘hindgut’ hypothesis suggests this rise is due to anatomical changes or increased intestinal transit (Nausheen, Shah et al. 2013, Strader, Vahl et al. 2005, Yousseif, Emmanuel et al. 2014). The ‘foregut’ hypothesis suggests the exclusion of the upper small intestine is responsible for the changes, possibly due to the decreased secretion of the anti-incretin factor (Meek, Lewis et al. 2016). In the current study, fasting plasma GLP-1 levels significantly increase post-opt regardless of surgery type and T2DM status and directly correlated with plasma ANXA1 levels. The direct correlation further reflects the insulin sensitising effects of ANXA1, which could be partly mediated by modulating the expression of DPP4.

6.7 Conclusion

Due to the low participant numbers (owing to several factors including, no incentives, pre-opt stress and travelling from afar) and poor retention rate (owing to several factors including, delayed surgeries and participants unable to attend nor reschedule their post-opt appointments in sufficient time to meet the timelines of this study) throughout the study, a firm conclusion cannot be formed, however, plasma ANXA1 levels show increasing trends in relation to decreasing BMI in all studied cohorts. The limitations of this study include, small cohort, only women during the follow up, limited follow up and non-randomisation.

However, the remission of dyslipidaemia and increase in plasma HDL levels (in particular) after bariatric surgery induced weight loss may partly modulate the expression of ANXA1 to reduce the risk of developing obesity associated co-morbidities, thus, it would be of significant interest to study correlations between ANXA1 and markers of cardiovascular risk is well in obese individuals and bariatric surgery patients.

Chapter 7: Evidence of aberrant inflammation in patients with Lipodystrophy

Plasma Annexin A1 levels inversely correlate with adiposity and markers of inflammation in obesity, however, it is unclear, whether this decrease is due to increasing fat mass or presence of low-grade chronic inflammation. Lipodystrophy patients have little to no fat mass yet display similar metabolic abnormalities as obese individuals. Therefore, the objective of this study was to investigate the plasma Annexin A1 levels in lipodystrophy patients and correlate it with anthropometric measures and inflammatory biomarkers to elucidate the relationship between plasma Annexin A1 levels, adiposity and inflammation.

7.1 Abstract

Introduction: Lipodystrophy and obesity represent extreme and opposite ends of the adiposity spectrum. Lipodystrophy diseases are characterised by abnormal accumulation, distribution and/or total loss of the adipose tissue. They are characterised by increased metabolic abnormalities and elevated risk of developing co-morbidities such as, type 2 diabetes mellitus, like obese individuals.

Aim: The aim of this study was to analyse plasma Annexin A1 (ANXA1) levels in lipodystrophy patients and correlate with inflammation to understand the relationship between ANXA1, adiposity and inflammation.

Methodology: Plasma samples of Lipodystrophy patients were gifted by Professor David Savage, Metabolic Research Laboratories, Cambridge University and were compared against body mass index (BMI) matched controls. Plasma samples were analysed, and results were statistically analysed using GraphPad Prism 5.0. Statistical significance at 95% was determined using One-tailed unpaired students T-test and One-tailed Pearson's correlation coefficient.

Results: Plasma ANXA1 levels were significantly reduced in lipodystrophy patients of all genotypes in comparison to control and inversely correlated with BMI and plasma C-reactive protein levels.

Conclusion: Therefore, it is proposed that attenuated levels of ANXA1 may play a role in inducing insulin resistance and development of associated co-morbidities by failing to resolve inflammation in obesity and by failing to regulate insulin sensitivity in lipodystrophy.

7.2 Introduction

Lipodystrophies are a heterogeneous group of diseases characterised by generalised or partial loss of adipose tissue. If the loss is localised, it is often associated with pathological accumulation of lipids in non-adipose tissue depots (Mantzoros 2012). Lipodystrophies have been reported in the literature for more than 100 years, but only recently new subsets of diseases have been recognised. The loss of adipose tissue resulting from genetic mutations (autosomal dominant and recessive subtypes) is extremely rare and loss resulting from autoimmune disorders or drugs such as, protease inhibitors or insulin injections is more common (Herranz, de Lucas et al. 2008). The adipose tissue of lipodystrophic patients is characterised by smaller adipocytes possibly due to abnormally low expression of crucial adipogenic transcription factors such as, peroxisome proliferator activated receptor γ (PPAR γ) and adipocyte differentiation related genes, greater cell size variation, disruption of cell membranes and signs of apoptosis (Bastard, Caron et al. 2002, Jan, Cervera et al. 2004).

Lipodystrophy diseases are associated with similar metabolic abnormalities as obese individuals, such as, hypertriglyceridemia, dyslipidaemia, hyperinsulinemia and insulin resistance and the onset of co-morbidities such as, type 2 diabetes mellitus (T2DM), hepatic steatosis, cardiovascular disease and cancer (Brown, Araujo-Vilar et al. 2016, Phan, Reue 2005). The severity of the metabolic abnormalities positively correlates with the degree of adipose tissue lost (Haque, Shimomura et al. 2002). For example, complete loss of adipose tissue in congenital generalised lipodystrophies (CGL) and acquired generalised lipodystrophies (AGL) induces extreme insulin resistance, whereas, partial loss of adipose tissue in familial partial lipodystrophies (FPL) and acquired partial

lipodystrophies (APL) induces mild to moderate insulin resistance with little to no metabolic complications (Haque et al. 2002).

CGL are autosomal recessive diseases caused by mutations of specific genes involved in lipid metabolism and adipocyte function, including *AGPAT2* (CGL1), *BSCL2* (CGL2), *CAVI* (CGL3) and *PTRF* (CGL4) (Table 7.1) (Brown et al. 2016). They are characterised by near to total loss of adipose tissue and prominent muscles at birth or shortly after (Agarwal, Simha et al. 2003). Due to the absence of fully functioning adipocytes, dietary and endogenously synthesised lipids are stored aberrantly in metabolically important tissues such as, muscle and liver inducing lipotoxicity and development of metabolic abnormalities (Van Maldergem, Magre et al. 2002). Similarly, AGL typically associated with autoimmune disorders (Table 7.2), are characterised by gradual loss of adipose tissue in individuals born with normal adipose tissue distribution. The loss typically begins in childhood or adolescence and is more frequent in females than in males (Misra, Garg 2003). CGL and AGL patients develop hyperphagia in early childhood, have accelerated growth, advanced bone age, features suggestive of acromegaly (enlarged hands and feet), acanthosis nigricans and early onset of T2DM (Herranz et al. 2008). Female patients have additional clinical features including hirsutism, clitoromegaly, irregular menstrual periods, polycystic ovaries and/or infertility (Garg 2004).

Subtype	Gene	Molecular Basis	Unique clinical features
CGL1	<i>AGPAT2</i>	<ul style="list-style-type: none"> - 1-Acylglycerol-3-Phosphate O-Acyltransferase 2 catalyses the formation of phospholipids and triglycerides in various organs - AGPAT isoform 2 is highly expressed in the adipose tissue 	<ul style="list-style-type: none"> - Metabolic active adipose tissue is lost - Mechanical adipose tissue is preserved - Lytic bone lesions in long bones are frequent
CGL2	<i>BSCL2</i>	<ul style="list-style-type: none"> - Seipin regulates lipid homeostasis and adipocyte differentiation - Lipid lipogenesis and lipid droplet accumulation in non-adipocytes - Promotes adipogenesis in periods of excess energy 	<ul style="list-style-type: none"> - Both metabolic active and mechanical adipose tissue are lost - Cardiomyopathy - Lytic bone lesions in long bones are occasional - Abnormal morphology of sperm that affects fertility
CGL3	<i>CAVI</i>	<ul style="list-style-type: none"> - Galectin-3 binds and transports fatty acids to lipid droplets 	<ul style="list-style-type: none"> - Metabolically active adipose tissue is lost - Mechanical and bone marrow adipose tissue is preserved - Functional enlargement of the oesophagus - Short stature, hypocalcaemia, vitamin D resistance
CGL4	<i>PTRF</i>	<ul style="list-style-type: none"> - Cavin-1 regulates the expression of caveolins 1 and 3 	<ul style="list-style-type: none"> - Metabolically active adipose tissue is lost - Mechanical and bone marrow adipose tissue is preserved - Cardiomyopathy - Sudden death - Congenital pyloric stenosis - Developmental delay - Muscle weakness

Table 7.1: Subsets of congenital generalised lipodystrophies. Congenital generalised lipodystrophies (CGL) diseases are characterised by general loss of adipose tissue caused by mutations in genes involved in adipocyte function and lipid metabolism. CGL are subdivided into 4 subtypes based on the gene involved and the severity of clinical features and metabolic abnormalities vary between the subsets. AGPAT2= 1-Acylglycerol-3-phosphate O-Acyltransferase 2, BSCL2= Seipin, CAVI= Galectin-3 and PTRF= Cavin-1.

(Herranz et al. 2008)

Subtype	Prevalence	Clinical features
Panniculitis associated AGL	~25%	<ul style="list-style-type: none"> - Initial development of panniculitis, followed by localised fat loss when lesions heal - Ongoing panniculitis later results in generalised loss of subcutaneous adipose tissue
Auto-immune AGL	~25%	<ul style="list-style-type: none"> - Gradual generalised fat loss associated with auto-immune disease; juvenile dermatomyositis
Idiopathic AGL	~50%	<ul style="list-style-type: none"> - Gradual loss of subcutaneous adipose tissue with unclear aetiology

Table 7.2: Subtypes of acquired generalised lipodystrophies. *Acquired generalised lipodystrophy (AGL) syndromes are characterised by general loss of adipose tissue caused by inflammatory and autoimmune diseases. (Herranz et al. 2008, Misra, Garg 2003)*

FPL are rare autosomal dominant diseases caused by mutations of specific genes involved in lipid metabolism and adipocyte function including; LMNA (FPL2), PPAR γ (FPL3), PLIN1 (FPL4) and AKT2 (AKT2-lined) (Table 7.3). Accumulation and distribution of adipose tissue is typically normal in FPL patients during early childhood, but selective and progressive loss occurs from various parts of the body around late childhood or puberty (Brown et al. 2016). Patients have reduced adipose tissue in the arms, legs, head and the trunk regions, with excessive accumulation in other parts of the body including the neck, face and intra-abdominal regions, giving them a ‘Cushingoid appearance’ (Herranz et al. 2008). Metabolic abnormalities are more common and severe in these patients and they are at a higher risk of developing co-morbidities during adulthood (Subramanyam, Simha et al. 2010, Vantyghem, Vincent-Desplanques et al. 2008). Women with FPL experience menstrual irregularities have a higher occurrence of developing polycystic ovarian syndrome and experience severe metabolic abnormalities compared to men (Garg 2000).

Subtype	Gene	Molecular Basis	Unique clinical features
FPL 1	<i>Unknown</i>		<ul style="list-style-type: none"> - Loss of adipose tissue is confined to arms and legs - May develop central obesity - Only reported in women
FPL 2	<i>LMNA</i>	<ul style="list-style-type: none"> - Lamin A is associated with adipocyte differentiation, insulin signalling and PPARγ - Mutations in Lamin A and C result in premature cell death of adipocytes 	<ul style="list-style-type: none"> - Increased amount of adipose tissue in the cervico-facial area - Premature aging - Metabolic laminopathies resembling the metabolic syndrome
FPL 3	<i>PPARG</i>	<ul style="list-style-type: none"> - Peroxisome proliferator activated receptor γ is essential for adipocyte differentiation 	<ul style="list-style-type: none"> - Loss of adipose tissue is more prominent in the calves and forearms - Metabolic abnormalities are more prominent
FPL 4	<i>PLIN1</i>	<ul style="list-style-type: none"> - Perilipin is the most abundant protein coating the surface of lipid droplets - Essential for storage triglycerides and release of fatty acids 	<ul style="list-style-type: none"> - Loss of adipose tissue is more prominent in the lower limbs and buttocks - Muscle hypertrophy may be more prominent in the calves
FPL 5	<i>CIDEA</i> (autosomal recessive)	<ul style="list-style-type: none"> - Cell death inducing DFFA like effector C is expressed and plays a role in storage of lipid within the lipid droplets 	<ul style="list-style-type: none"> - Loss of adipose tissue in the arms and legs - Hypertension and severe insulin resistance
FPL 6	<i>LIPE</i> (autosomal recessive)	<ul style="list-style-type: none"> - Hormone sensitive lipase is involved in triacylglycerol degradation and glycerolipid metabolism 	<ul style="list-style-type: none"> - Partial lipodystrophy - Severe insulin resistance - Fatty liver - Acanthosis nigricans
AKT2-linked	<i>AKT2</i>	<ul style="list-style-type: none"> - Protein Kinase B β is thought to play a role in post receptor insulin signalling and may be involved in regulating PPARγ 	<ul style="list-style-type: none"> - Severe lipodystrophy - Impaired insulin activity

Table 7.3: Subsets of familial partial lipodystrophies. Familial partial lipodystrophy (FPL) diseases are characterised by partial loss of adipose tissue caused by mutations in genes involved in adipocyte function and lipid metabolism. FPL are subdivided into different subtypes based on the gene involved and the severity of clinical features and metabolic abnormalities vary.

LMNA= Lamin A, *PPARG*= Peroxisome proliferator activated receptor γ , *PLIN1*= Perilipin, *CIDEA*= Cell death inducing DFFA like effector C, *LIPE*= Hormone sensitive lipase and *AKT2*= Protein kinase B β (Herranz et al. 2008, Brown et al. 2016)

Lipodystrophies and obesity represent extreme and opposite ends of the adiposity spectrum; however, they induce similar metabolic abnormalities and increase the risk of developing co-morbidities (Herrero, Shapiro et al. 2010). Lipodystrophic mice display increased systemic inflammation with increased expression of proinflammatory cytokines such as, tumour necrosis factor α (TNF α), interleukin 1 β and interleukin 6 (IL6) and decreased expression of anti-inflammatory cytokines such as, interleukin 10 (Herrero et al. 2010). Furthermore, mRNA expression of essential adipokines such as, leptin, adiponectin and resistin are downregulated in the lipodystrophic adipose tissue and are consistent with the degree of lipoatrophy and metabolic abnormalities (Herrero et al. 2010). Human immunodeficiency virus (HIV)-associated lipodystrophy patients undergoing highly active antiretroviral therapy display increased systemic proinflammatory cytokines such as, TNF α and IL6 (Johnson 2004). However, data regarding the plasma profiles of inflammatory biomarkers in other forms of human lipodystrophies is limited. Therefore, comparing the underlying mechanisms of obesity and lipodystrophies may further our understanding of the relationship between adiposity, inflammation and the onset of the metabolic syndrome.

7.2.2 Aim, objectives and Hypothesis

The aim of this study was to analyse plasma ANXA1 levels in lipodystrophy patients and correlate with an inflammatory biomarker. It was hypothesised that, plasma ANXA1 levels are reduced in lipodystrophy patients and inversely correlate with inflammation.

The objectives of this study were:

- To analyse and correlate the plasma ANXA1 levels in lipodystrophy patients and BMI matched controls
- To analyse and correlate the plasma C-reactive protein (CRP) levels in lipodystrophy patients and BMI matched controls
- To correlate plasma ANXA1 levels with plasma CRP levels of lipodystrophy patients and BMI matched control

7.4 Materials and Methods

7.4.1 Participant samples

The lipodystrophy plasma samples and associated anthropometric data were gifted by Professor David Savage, Metabolic Research Laboratories, Cambridge University and were compared against BMI matched controls.

7.4.2 Metabolic measurements

7.4.2.1 Enzyme linked immunosorbent Assays

Plasma inflammatory biomarkers, fasting plasma insulin levels and fasting plasma gut hormones were quantified using enzyme linked immunosorbent assay (ELISA). The assays were based on the sandwich technique, as described in Chapter 3 (section 3.3).

- The R&D systems Human C-Reactive Protein (CRP) Quantikine ELISA kit (#DCRP00) was used to quantify CRP in plasma samples, following the manufacturers protocol. The mean intra-assay precision was determined using coefficient of variation (CV). The mean CV of the CRP assay was 2.76%.
- The Cusabio Human Annexin I (ANXAI) ELISA kit (#CSB-E12155h) was used to quantify Annexin A1 levels in plasma samples, following the manufacturers protocol. The mean %CV of the ANXA1 assay was 6.38%.

7.4.3 Statistical analysis

Data presented in this study is expressed as Mean \pm Standard Error of the Mean (SEM). GraphPad Prism 5 was used to statistically analyse the data and produce the graphs. One-tailed, unpaired students T-test was used to compare control and disease subsets individually and one-tailed Pearson's correlation coefficient to statistically analyse the

data and to assess the differences in the data sets, a p value of $p < 0.05$ was considered statistically significant.

7.5 Results

7.5.1 Anthropometric measures

Plasma samples from 9 lipodystrophy patients and 31 BMI matched controls were analysed for ANXA1 and CRP in the current study. No statistically significant difference was observed between the average BMI of lipodystrophy patients ($23.9 \pm 1.5 \text{ kg/m}^2$, $n=9$) and matched controls ($23.2 \pm 0.6 \text{ kg/m}^2$, $n=29$), $p=0.324$ (Figure 7.1).




Figure 7.1: Average BMI of lipodystrophy patients and matched controls. The average BMI of 9 lipodystrophy patients and 29 controls were compared. No statistically significant difference was observed between the two study groups ($p=0.324$). One-tailed, unpaired student's T-test was used to statistically analyse the data. Data presented as Mean \pm SEM. BMI= Body Mass Index

Similarly, no statistically significant difference was observed between the average BMI of lipodystrophy patients and matched controls when characterised per BMI group, underweight (BMI <19.9); 16.4 kg/m², n=1 vs 19.1±0.1 kg/m², n=1 healthy weight (BMI 20-24.9); 22.7±0.7 kg/m², n=5 vs 22.4±0.4 kg/m², n=5, p=0.339, overweight (BMI 25-29.9); 26.7±1.4 kg/m², n=2, 27.1±0.6 kg/m², n=7, p=0.400 and obese (BMI 30+); 31.6 kg/m², n=1 vs 31.1±0.0 kg/m², n=1, respectively (Figure 7.2).




Figure 7.2: Average BMI of control and lipodystrophy patients per BMI categories. The average BMI of 9 lipodystrophy patients and 29 controls were compared per BMI category. One-tailed, unpaired student's T-test was used to statistically analyse the data. No statistically significant difference was observed between the two study groups who were underweight, healthy (p=0.284), overweight (p=0.400) or obese. Data presented as Mean±SEM. BMI= Body Mass Index

Furthermore, no statistically significant difference was observed between the average BMI of lipodystrophy patients when characterised per disease subtype; CGL vs FPL (22.0±0.6 kg/m², n=2 vs 24.7±1.5 kg/m², n=4, p=0.147), CGL vs AGL (20.2±3.8 kg/m², n=2, p=0.346) and FPL vs AGL (p=0.118) (Figure 7.3).




Figure 7.3: Average BMI of lipodystrophy patients based on subtypes. Lipodystrophy diseases are categorised into different subtypes based on their aetiology and degree of adipose tissue loss. The average BMI of CGL, FPL and AGL were compared. One-tailed, unpaired student's T-test was used to statistically analyse the data. No statistically significant difference was observed between the subtypes; CGL vs FPL ($n=2$ and $n=4$ respectively, $p=0.147$), CGL vs AGL ($n=2$, $p=0.346$) and FPL vs AGL ($n=4$ and $n=2$ respectively, $p=0.118$). Data presented as Mean \pm SEM. BMI= Body Mass Index, CGL= Congenital generalised lipodystrophy, FPL= Familial partial lipodystrophy and AGL= Acquired generalised lipodystrophy

7.5.2 Inflammatory Biomarkers

7.5.2.1 Plasma CRP

The average plasma CRP levels were significantly higher in lipodystrophy patients compared to control (3.3 ± 1.0 $\mu\text{g/mL}$, $n=9$ vs 1.4 ± 0.3 $\mu\text{g/mL}$, $n=29$, $p=0.021$), as shown in Figure 7.4. However, no significant correlation was observed between plasma CRP levels and BMI, as shown in Figure 7.5. The average plasma CRP levels were higher in healthy weight compared to underweight controls (1.7 ± 0.4 $\mu\text{g/mL}$, $n=5$ vs 0.9 ± 0.2 $\mu\text{g/mL}$, $n=17$), but was lower in overweight compared to healthy weight controls (1.1 ± 0.5 $\mu\text{g/mL}$, $n=7$) and in obese compared to overweight controls (0.47 ± 0.0 $\mu\text{g/mL}$, $n=1$), as shown in Figure 7.6. The average CRP levels in lipodystrophy patients followed a similar

pattern when categorised based on BMI. The average CRP levels were higher in healthy weight compared to underweight patients (3.1 ± 1.1 $\mu\text{g/mL}$, $n=5$ vs 0.4 ± 0.0 $\mu\text{g/mL}$, $n=1$), in overweight compared to healthy weight patients (4.6 ± 4.1 $\mu\text{g/mL}$, $n=2$) and lower in obese compared to overweight patients (1.0 ± 0.0 $\mu\text{g/mL}$, $n=1$), as shown in Figure 6.6. The average plasma CRP levels were significantly higher in healthy weight lipodystrophy patients compared to control ($p=0.034$). The average plasma CRP levels were significantly increased in FPL and AGL patients compared to control (3.9 ± 1.8 $\mu\text{g/mL}$, $n=4$, $p=0.0079$ and 4.0 ± 3.6 $\mu\text{g/mL}$, $n=2$, $p=0.0254$ vs 1.4 ± 0.3 $\mu\text{g/mL}$, $n=29$, respectively). However, no significant difference was observed between the average plasma CRP levels between the different lipodystrophy disease subtypes; CGL vs FPL (2.2 ± 1.8 $\mu\text{g/mL}$, $n=2$, $p=0.283$) CGL vs AGL ($p=0.336$) and FPL vs AGL ($p=0.491$), as shown in Figure 7.7.




Figure 7.4: Plasma CRP (µg/mL) levels in lipodystrophy patients vs control. Plasma CRP levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls. One-tailed, unpaired student's T-test was used to statistically analyse the data. The average plasma CRP levels were significantly higher in lipodystrophy patients compared to control ($p=0.021$). Data presented as Mean \pm SEM. P value was set at $p < 0.05$ as denoted by *. BMI= Body mass index and CRP= C-reactive protein




Figure 7.5: Correlation between plasma CRP levels and BMI. Plasma CRP levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls and correlated with BMI. One tailed Person correlation coefficient was used to statistically analyse the data.
BMI= Body Mass Index, CRP= C-reactive protein




Figure 7.6: Plasma CRP ($\mu\text{g/mL}$) levels in lipodystrophy patients vs BMI matched control per BMI category. Plasma CRP levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls. One-tailed, unpaired student's T-test was used to statistically analyse the data. The average plasma CRP levels were higher in healthy weight compared to underweight controls, lower in overweight compared to healthy weight controls and in obese compared to overweight controls, however, no statistically significant difference was observed between each group. Similarly, plasma CRP levels were higher in healthy weight compared to underweight lipodystrophy patients, in overweight compared to healthy weight lipodystrophy patients and lower in obese compared to overweight lipodystrophy patients, however, no statistically significant difference was observed between each group. Plasma CRP levels were significantly higher in healthy weight lipodystrophy patients compared to healthy weight controls ($n=5$, $n=16$, $p=0.034$). Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *. CRP= C-reactive protein, BMI= Body Mass Index





Figure 7.7: Plasma CRP (µg/mL) levels in lipodystrophy patients categorised per disease subtype. Plasma CRP levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls. One-tailed, unpaired student's T-test was used to statistically analyse the data. Plasma CRP levels significantly higher in FPL and AGL patients compared to controls (n=4, p=0.0079 and n=2, p=0.0283, respectively). No statistically significant difference was observed between the disease subtypes. Data presented as Mean±SEM. P value was set at p <0.05.

BMI= Body mass index, CGL= Congenital generalised lipodystrophy, FPL= Familial partial lipodystrophy, AGL= Acquired generalised lipodystrophy, CRP= C-reactive protein

7.5.2.2 Plasma Annexin A1

The average plasma ANXA1 levels were significantly lower in lipodystrophy patients compared to controls (0.2 ± 0.1 ng/mL, n=9 vs 0.94 ± 0.1 ng/mL, n=29, p=0.001), as shown in Figure 7.8. Although, an inverse correlation was observed between plasma ANXA1 levels and BMI, it was not statistically significant, as shown in Figure 7.9. The average plasma ANXA1 levels were higher in healthy weight compared to underweight controls (1.3 ± 0.2 ng/mL, n=17 vs 0.7 ± 0.2 ng/mL, n=5), but was lower in overweight compared to healthy weight controls (0.7 ± 0.2 ng/mL, n=7) and in higher in obese compared to

overweight controls (0.019 ± 0.0 ng/mL, $n=1$), as shown in Figure 7.10. The average ANXA1 levels in lipodystrophy patients followed a similar pattern when categorised based on BMI. The average ANXA1 levels were higher in healthy weight compared to underweight patients (0.2 ± 0.1 ng/mL, $n=5$ vs 0.01 ± 0.0 ng/mL, $n=1$), overweight compared to healthy weight patients (0.5 ± 0.03 ng/mL, $n=2$) and lower in obese compared to overweight patients (0.08 ± 0.0 ng/mL, $n=1$), as shown in Figure 7.10. The average plasma ANXA1 levels were significantly lower in healthy weight lipodystrophy patients compared to control ($p=0.002$). The average plasma ANXA1 levels were significantly lower in FPL patients compared to control (0.3 ± 0.1 ng/mL, $n=4$ vs 0.94 ± 0.1 ng/, $n=29$, $p=0.021$). However, no significant difference was observed between the average plasma ANXA1 concentrations between the different lipodystrophy disease subtypes; CGL vs FPL (0.08 ± 0.0 ng/mL, $n=2$) CGL vs AGL (0.4 ± 0.3 ng/mL, $n=4$) and FPL vs AGL ($p=0.312$), as shown in Figure 7.11. Although an inverse correlation between plasma ANXA1 and plasma CRP was observed, it was not statistically significant, as shown in Figure 7.12.






Figure 7.9: Correlation between Plasma Annexin A1 levels and BMI. Plasma ANXA1 level were analysed in 9 lipodystrophy patients and 29 BMI matched controls and correlated with BMI. One tailed Person correlation coefficient was used to statistically analyse the data. An inverse correlation between the two biomarkers was observed, however it was not of statistical significance. BMI= Body Mass Index and ANXA1= Annexin A1




Figure 7.10: Plasma Annexin A1 (ng/mL) levels in lipodystrophy patients vs control per BMI category. Plasma ANXA1 levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls. One-tailed, unpaired student's T-test was used to statistically analyse the data. The average plasma ANXA1 levels were higher in healthy weight compared to underweight controls, lower in overweight compared to healthy weight controls and in obese compared to overweight controls, however, no statistically significant difference was observed between each group. Similarly, plasma ANXA1 levels were higher in healthy weight compared to underweight lipodystrophy patients and lower in overweight compared to healthy weight lipodystrophy patients and in obese compared to overweight lipodystrophy patients, however, no statistically significant difference was observed between each group. Plasma ANXA1 levels were significantly lower in healthy weight lipodystrophy patients compared to healthy weight controls ($n=5$ and $n=17$, respectively, $p=0.007$). Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *. BMI= Body Mass Index and ANXA1= Annexin A1




Figure 7.11: Plasma Annexin A1 (ng/mL) levels in lipodystrophy patients characterised per disease subtype. Plasma ANXA1 levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls. One-tailed, unpaired student's T-test was used to statistically analyse the data. Plasma ANXA1 levels were lower in disease subtypes compared to control. Plasma ANXA1 levels were significantly lower in FPL patients compared to controls ($n=4$, $p=0.021$, respectively). No statistically significant difference was observed between the disease subtypes. Data presented as Mean \pm SEM. P value was set at $p < 0.05$ and is denoted by *. BMI= Body mass index, ANXA1= Annexin A1, CGL= Congenital generalised lipodystrophy, FPL= Familial partial lipodystrophy and AGL= Acquired generalised lipodystrophy.




Figure 7.12: Correlation between plasma Annexin A1 and plasma CRP levels. Plasma Annexin A1 levels and plasma CRP levels were analysed in 9 lipodystrophy patients and 29 BMI matched controls and correlated. An inverse correlation was observed between the two biomarkers however, it was not of statistical significance. One tailed Person correlation coefficient was used to statistically analyse the data. BMI= Body mass index and CRP= C-reactive protein

7.6 Discussion

In the current study, plasma ANXA1 levels in lipodystrophy patients were analysed and correlated with the pro-inflammatory biomarker, CRP. Interestingly, plasma ANXA1 levels were significantly decreased in lipodystrophy patients compared to control and inversely correlated with BMI, in a similar manner that is observed in obese individuals, suggesting the degree of adiposity may not play a significant role in attenuating plasma ANXA1 levels (Kosicka, Cunliffe et al. 2013). Furthermore, plasma CRP levels were significantly increased in lipodystrophy patients in comparison to controls, suggestive of systemic inflammation and were inversely correlated with plasma ANXA1 levels. The results of the current study indicate plasma ANXA1 levels are attenuated possibly due to an increased inflammatory environment, however, to further understand the relationship between plasma ANXA1 and adiposity, correlation with % body fat (in non-lipodystrophy population) or adipokines may serve as a better indicator than BMI. Furthermore, adiposity and the deposition of fat depots differs amongst lipodystrophy subtypes, however, due to low patient numbers (as low as n=2 for CGL and AGL) the results are inconclusive, therefore, it is of interest to increase subject numbers to investigate relationship between plasma ANXA1 and adiposity.

The origin of inflammation in lipodystrophy varies amongst the different subtypes and may depend on the aetiology. For example, inflammation in congenital lipodystrophies is thought to originate from dystrophic adipose tissue, ectopic lipid accumulation and lipotoxicity (Mantzoros 2012). Whereas in, acquired lipodystrophies induced by autoimmune diseases such as HIV, the increased inflammatory environments created by increased secretion of proinflammatory cytokines and chemokines, and macrophage expression of markers in response to the virus (Ryden, Arner 2007). Anti-retroviral

therapy drugs prescribed to treat HIV further impair adipocyte differentiation by downregulating crucial transcription factors such as PPAR γ , induce the generation of reactive oxygen species, increase recruitment of macrophages, inhibit glucose transporter 4 mediated glucose uptake, impair insulin signalling, alter the secretion of adipokines; leptin and adiponectin and enhance production and secretion of proinflammatory cytokines such as IL6 and TNF α (Fiorenza, Chou et al. 2011, Blumer, van Vonderen et al. 2008, Divi, Haverkos et al. 2007, Pacenti, Barzon et al. 2006). Furthermore, they increase lipolysis leading to elevated levels of free fatty acids and ectopic accumulation in non-adipose tissue depots, inducing lipotoxicity and worsening the metabolic profile in these patients (Villarroya, Domingo et al. 2010).

Obesity induced inflammation is hypothesised to be a contributing factor in the development of associated metabolic abnormalities and early on-set of metabolic diseases such as, T2DM however, it is considered to play a differential role in lipodystrophy (Herrero et al. 2010). For example, increased infiltration of classically activated macrophages (M1) into the adipose tissue and increased production of TNF α and IL6 are hypothesized to interfere with the normal activity of insulin receptors and contribute to the development of insulin resistance in obesity (Strissel, Stancheva et al. 2007). Although, macrophage infiltration is also increased lipodystrophic adipose tissue, their phenotype and behaviour (mixture of M1 and M2) is very different. They are believed to be involved in tissue remodelling and repair due to increased apoptosis and not in the pathogenesis of insulin resistance (Herrero et al. 2010). Furthermore, anti-inflammatory drugs known to decrease inflammation and improve hyperglycaemia by lowering glucose and insulin levels in human obese and diabetic individuals do not have similar effects in lipodystrophy mice (Herrero et al. 2010). Instead, the altered synthesis and secretion of

adipokines is hypothesised to induce lipodystrophy associated metabolic abnormalities. Plasma leptin levels positively correlate with adiposity, therefore, it is not surprising that the levels are significantly reduced in CGL and AGL patients, but not as much in FPL patients (Pardini, Victoria et al. 1998, Hegele, Cao et al. 2000). Leptin replacement therapy markedly improves dyslipidaemia, insulin sensitivity, reduces haemoglobin A1c levels and intrahepatic lipid content in lipodystrophy patients (Mantzoros 2012). Similarly, plasma adiponectin levels are significantly reduced in CGL and AGL patients, whereas not so much in FPL and APL patients (Haque et al. 2002). The low levels of adiponectin are proposed to induce insulin resistance and the metabolic syndrome in lipodystrophy (Trujillo, Scherer 2005).

7.7 Conclusion

In addition to its pro-resolving properties, ANXA1 is proposed to mediate insulin sensitising effects, therefore, attenuated levels may contribute to the onset of insulin resistance by failing to resolve inflammation in obesity and by failing to regulate insulin sensitivity in lipodystrophy (Lindgren, Nilsson et al. 2001). Therefore, it would be of interest to correlate plasma ANXA1 levels with glycaemic biomarkers using a larger study cohort (and greater anthropometric measures and biomarkers) to further understand the relationship between attenuated plasma ANXA1 levels and insulin resistance in patients with varying degrees of adiposity.

Chapter 8: General Discussion

The primary aim of this project was to investigate the role of ANXA1 in obesity and associated inflammation. The secondary aims and results are summarised in table 8.1.

Study	Aim	Results
1 (Chapter 4)	To investigate the potential roles of ANXA1 in adipocytes using a model mimicking the inflammatory status that is observed during obesity	Acute AC2-26 treatment significantly downregulated the mRNA expression of pro-inflammatory, insulin resistant genes and upregulated the mRNA expression of anti-inflammatory, insulin sensitive genes.
2 (Chapter 5)	To investigate the role of ANXA1 in the differentiation of preadipocytes into mature adipocytes	ANXA1 may regulate the rate of adipogenesis through differentional regulation of MAPK intracellular pathways.
3 (Chapter 6)	To investigate the plasma ANXA1 levels in response to rapid weight loss induced by Bariatric Surgery	Plasma ANXA1 levels show increasing trends with decreasing BMI in the full cohort, SG participants, RYGB participants, non-T2DM participants and T2DM participants
4 (Chapter 7)	To investigate the plasma ANXA1 levels in Lipodystrophy patients	Plasma ANXA1 levels are reduced in lipodystrophy patients and inversely correlate with plasma CRP levels

Table 8.1: Summary of Aims and Results of the project.

ANXA1= Annexin A1, MAPK= Mitogen activated protein kinase, BMI= Body mass index, CRP= C-reactive protein, SG= Sleeve gastrectomy, RYGB= Roux-en-Y gastric bypass, T2DM= Type 2 diabetes mellitus

The *in vitro* results from study 1 (Chapter 4) of this thesis indicate several roles of ANXA1 in adipocyte physiology in obesity. The differential regulation of genes unknown to be dysregulated in obesity in response to AC2-26 treatment indicate ANXA1 may be involved in regulating cellular survival and apoptosis, cellular proliferation and differentiation, lipid and glucose metabolism and, intracellular stress and inflammation. It is interesting to note that acute AC2-26 treatment downregulated the expression of genes such as, *CD36*, *DPP4*, *LEP* and *TNF* that are involved in inducing insulin resistance and upregulated the expression of genes such as, *ADIPOQ* that are involved in inducing insulin sensitivity in mature obesogenic adipocytes, reflecting an insulin sensitising effect

of ANXA1. Insulin mediates its biological responses by inducing tyrosine phosphorylation of insulin receptor substrate (IRS) proteins which in turn activate downstream signalling pathways (Figure 8.1). Agents such as, inflammatory cytokines, chemokines and adipokines, free fatty acids and cellular stress attenuate insulin activity and induce insulin resistance by modulating IRS phosphorylation (tyrosine phosphorylation activates and serine phosphorylation inhibits the activity) (Gual, Le Marchand-Brustel et al. 2005, Godsland 2009). In addition to changes in phosphorylation of IRS proteins, downregulation of *IRS1* gene expression is considered to play a central role in altered IRS1 tyrosine phosphorylation in adipocytes of obese and T2DM humans and animals (White 2002, Thirone, Huang et al. 2006). Therefore, it was surprising to observe that acute AC2-26 treatment significantly downregulated *IRS1* gene expression in mature obesogenic adipocytes. Though the exact mechanisms and reasoning of this observation are unclear, ANXA1 has shown to promote a dose-dependent inhibition of insulin receptor autophosphorylation, inhibiting its tyrosine kinase activity (Melki, Hullin et al. 1994). These conflicting results warrant further investigation.

Furthermore, acute AC2-26 treatment differentially regulated the expression of genes associated with lipid accumulation and adipogenesis in mature obesogenic adipocytes. It upregulated the expression of genes associated with inhibiting adipogenesis or adipocyte hypertrophy such as, *PPARA* and *PER3*, and downregulated the expression of genes associated with enhancing adipogenesis such as, *PPP3CC*, perhaps indicating a protective role of ANXA1 against adipocyte hypertrophy. Therefore, the role of ANXA1 in adipogenesis was investigated. The *in vitro* results from study 2 (Chapter 5) of this thesis indicate that ANXA1 and its peptide (AC2-26) may regulate the rate of adipogenesis in the latter differential stages through differential regulation of mitogen activated protein

kinase (MAPK) intracellular pathways. Interestingly, mature adipocytes differentiated with specific ANXA1 receptor (FPR2/ALX) antagonist, WRW4 or non-specific receptor agonist, AC2-26 (can activate FPR1 and FPR2/ALX) accumulated significantly less lipids and indicated an immature adipocyte phenotype compared to control. Whereas, mature adipocytes differentiated with low doses of non-specific receptor agonist, BOC-2 (low doses block FPR1 signalling) accumulated significantly more lipids than control. However, no significant difference in lipid accumulation was observed in mature adipocytes differentiated with a high dose of BOC-2 (higher doses block FPR1 and FPR2/ALX). Therefore, the increase in ANXA1 in mature adipocytes and further increase in obese adipocyte suggests, ANXA1 may act in an autocrine and paracrine manner to regulate adipocyte hypertrophy (Kosicka, Cunliffe et al. 2013, Henegar, Tordjman et al. 2008, Alfadda, Benabdelkamel et al. 2013). Hypertrophy is considered to be a key underlying cause of adipocyte dysfunction contributing to the onset of metabolic syndrome and obesity associated co-morbidities, thereby, indicating a protective role of ANXA1, however, this effect may be overshadowed by the sustained presence of nutrients and low grade chronic inflammation (Hirsch, Fried et al. 1989, Danforth 2000, Heilbronn, Smith et al. 2004).




Figure 8.1: Pathways activated by the insulin receptor. Binding of insulin to the insulin receptor activates two intracellular signalling pathways; MAPK signalling pathway that is involved in cell growth and survival and the PI3K signalling pathway that is involved in cellular proliferation and metabolism. MAPK=Mitogen activated protein kinase, PI3K= Phosphoinositide 3 kinase, IRS= insulin receptor substrate

Extracellular ANXA1 is cleaved by proteases to generate smaller peptides including, AC2-26 which are thought to retain the biological functions of ANXA1 (Gavins, Hickey 2012). However, how much ANXA1 is cleaved and how much remains full length is unknown. Unlike ANXA1, AC2-26 is able to bind to and activate formyl peptide receptor (FPR)-1 and FPR2/ALX (Dalli, Montero-Melendez et al. 2012). The activation of different receptors and intracellular signalling pathways by AC2-26 may contribute to the conflicting results observed in Study 1. Therefore, it would be of interest to investigate the roles of FPR1 and FPR2/ALX individually in mature obesogenic adipocytes and adipogenesis using receptor knockout models. Furthermore, adipocyte differentiation

with FPR1 and/or FPR2/ALX receptor knockouts would further emphasise the results from study 2 (Chapter 5) and confirm the results observed are due to differential activation rather than side-effects of the agonist and antagonists.

The results from the clinical pilot study 3 (Chapter 6) indicate increasing trends of plasma ANXA1 levels with decreasing body mass index (BMI). However, due to small participant numbers and poor retention it is difficult to form a solid conclusion, therefore, a full study in this area is required to determine their association. Interestingly, plasma ANXA1 directly correlated with plasma levels of an incretin hormone, glucagon-like peptide 1 (GLP-1). GLP-1 is proteolytically cleaved by dipeptidyl peptidase 4 (DPP4), thereby inhibiting its incretin effect (Bae 2016). DPP4 inhibitors are a new class of anti-hyperglycaemic agents used to treat T2DM (Ahren 2007). Furthermore, DPP4 is proposed to play a role in linking obesity and the pathogenesis of T2DM by acting as a local mediator of inflammation and insulin resistance in the adipose and hepatic tissues (Deacon 2019). Interestingly, the clinical results from study 1 (Chapter 4) show acute treatment of AC2-26 significantly downregulated the gene expression of *DPP4* in mature obesogenic adipocytes. The increase in plasma ANXA1 in response to weight loss may act in an autocrine, paracrine and endocrine manner to decrease the synthesis and secretion of DPP4, reflected by direct correlations with plasma GLP-1 levels. Therefore, it is of significant interest to further investigate the relationship between ANXA1 and DPP4 locally and systemically to dissect the mechanisms underlying the development of obesity associated co-morbidities for therapeutic purposes.

Furthermore, the results from the clinical pilot study 3 (Chapter 6) show inconsistent trends. Such as, plasma ANXA1 levels directly correlates with plasma levels of both anti- and pro-inflammatory biomarkers. These reasons behind these associations are unclear,

however, it is speculated that the invasive nature of the surgery may have skewed the findings. Most of the studies carried out in this area report pre- and >12 months post bariatric surgery data; therefore, it is of interest to continue investigating the plasma biomarkers for a longer follow up period. Furthermore, some participants within the study developed post-operative (post-opt) infections, which may have contributed to the inconsistent trends. Thus, the inclusion/exclusion criteria should be reviewed and patients with post-opt infections should be excluded in the future studies, as the biomarkers analysed form part of the inflammatory cascade and post-opt infections could potentially skew the results, giving false positive or negative results.




Figure 8.2: Proposed plasma Annexin A1 levels in relation to BMI or % body fat . Plasma Annexin A1 levels are reduced in individuals with low BMI/%BF and in individuals with high BMI/%Body fat, therefore, it is proposed that plasma Annexin A1 levels follow a normal distribution pattern where the plasma levels are highest in the healthy BMI/%Body fat range.

Finally, the results from the *in vivo* study 4 (Chapter 7) show plasma ANXA1 levels are significantly reduced in lipodystrophy patients compared to BMI matched controls. This observation is interesting because plasma ANXA1 levels are also significantly reduced in obese individuals and inversely correlate with BMI (Kosicka, Cunliffe et al. 2013) (Figure

8.2). This suggests the regulation of plasma ANXA1 levels may not be in direct relation to the amount of adipose tissue, but rather the presence of local and systemic low-grade chronic inflammation.

8.1 Conclusion

This project describes evidence for two different but overlapping functions of ANXA1. The first of which is produced locally within the adipose tissue depots and may have a role in regulating adipogenesis and reducing adipocyte hypertrophy to prevent adipocyte dysfunction, local inflammation and development of insulin resistance and secondly peripheral circulating levels (the source of which is still unknown) have been demonstrated to correlate with inflammatory biomarkers, markers of insulin sensitivity and gut hormone. Peripherally, ANXA1 may act in an endocrine manner to regulate inflammatory biomarkers to dampen inflammation, regulate insulin secretion and improve the metabolic profile to reduce the risk of developing obesity associated co-morbidities. However, this effect may be overshadowed by the continued increase in systematic inflammation associated with rapid adipose tissue expansion.

8.3 Future Studies

The results of Study 1(Chapter 4) and Study 2 (Chapter 5) indicate a protective role of ANXA1/AC2-26 in mature adipocytes incubated in an environment mimicking the inflammatory status observed in obesity by regulating the expression of genes and pathways associated with key adipocyte functions through alleviating the pro-inflammatory status, restoring metabolic mechanisms and dampening inflammation. To further validate the results of the current studies and to improve the quality and reliability of the data the following studies are suggested:

1) Analyse the expression of genes that were identified to be significantly altered in mature adipocytes that were incubated in an environment mimicking the inflammatory status observed in obesity (Study 1) using gene deletion (CRISPR) or knock down (siRNA) assays. Deleting the ANXA1 gene would further ensure the results are due to the change in ANXA1 rather than a side effect of the agonist treatment, limiting the experimental errors. Furthermore, primary adipocytes obtained from visceral adipose tissue depots could be used as models to investigate if the role or effect of ANXA1/AC2-26 differs amongst different adipose tissue depots.

2) The results of Study 2 indicate that FPR1 is involved in regulating the rate of adipocyte differentiation. Differentiation of SGBS with individual and/or grouped gene deletions or knock downs of ANXA1 and/or its receptors; FPR1 and/or FPR2/ALX into mature adipocytes would further emphasise and validate the role of ANXA1, FPR1 and FPR2/ALX in adipocyte differentiation. Similar experiments would certify the role of ANXA1 and/or its receptors in glucose uptake in mature adipocytes.

3) White adipocyte trans-differentiation to a brown adipocyte is considered a novel therapeutic target for obesity. Acute AC2-26 treatment significantly upregulated the gene expression of *PPARA*, a master regulator of brown adipocyte differentiation in mature obesogenic adipocytes. This data indicates that ANXA1/AC2-26 may induce the expression of genes associated with brown adipocytes in white adipocytes to utilise excess energy in the form of thermogenesis to reduce adipocyte hypertrophy and associated dysfunctions. Therefore, it would be of significant interest to investigate the role of ANXA1/AC2-26 in trans-differentiation for therapeutic purposes.

4) The adipose tissue is a heterogenous tissue therefore, it would be of significant interest to carry out co-culture experiments to investigate the paracrine effects of ANXA1 on immune cells to reflect the physiological state. It would be of significant interest to investigate the effects of ANXA1 synthesised and secreted by adipocytes on monocyte and macrophage migration, and macrophage polarisation.

The question whether plasma ANXA1 is regulated by adiposity or inflammation remains inconclusive. The results of Chapter 5 (Study 3) and Chapter 6 (Study 4) indicate the plasma levels of ANXA1 may be regulated by inflammation, however, it is difficult to form a firm conclusion based on small, short time framed and incomprehensive studies. Therefore, the following developments are suggested:

5) It would be of significant interest to carry out the clinical pilot study 3 on a larger scale (carry out a para-equation based on population size to determine the optimal number of participants required at each follow up) and a longer follow up (on average the studies in this field of study follow up to 12-24 months post opt). Following patients for up to 24 months with follow ups every 3 months would indicate the patterns of plasma ANXA1

proteins and other biomarkers and whether the patterns are exponential or reach a plateau. Furthermore, the difference in plasma ANXA1 in response to different bariatric surgeries and diabetes mellitus status are of interest, therefore, a full separate study would be designed to investigate the differences between these groups (wide array of anthropometric measures and plasma biomarkers). If the evidence supports it, ANXA1 agonist could be given to reduce inflammatory effects in patients with T2DM.

6) The clinical pilot study 3 aimed to investigate insulin sensitivity/resistance using oral glucose tolerance test (OGGT) pre-operatively (pre-opt) and post-operatively (post-opt) and correlate it with plasma ANXA1, however, due to small cohort and poor retention, it was not feasible. The OGGT is a more reliable test to determine insulin sensitivity/resistance than using fasting plasma glucose and plasma insulin values. Furthermore, GLP-1 and peptide YY are secreted in response to nutrients, therefore, it would be of significant interest to investigate the plasma levels of these biomarkers pre- and post-opt postprandially than fasted and correlate them with plasma ANXA1 levels to achieve physiologically accurate results.

7) To avoid the effects of surgery, it would be of significant interest to investigate the plasma levels of inflammatory biomarkers in individuals undergoing weight loss through non-surgical methods (lifestyle, exercise and/or diet).

8) In addition to investigating the plasma levels of inflammatory biomarkers and correlating them with ANXA1, it would be of significant interest to investigate the effect of weight loss on the expression and secretion of ANXA1 in adipocytes. However, the exact source of plasma ANXA1 is unknown, therefore it would be of significant interest to investigate what percentage of ANXA1 is secreted by the adipose tissue and how the

percentage changes in response to weight loss, as this would further strengthen the association between ANXA1 and adiposity.

9) It would be of significant interest to investigate the plasma levels of inflammatory biomarkers, insulin sensitivity and adipokines and correlate them with plasma ANXA1 levels in lipodystrophy patients. The results of this study would further contribute to our understanding the regulation of ANXA1 and its relationship with adiposity.

References

- ADRIAN, T.E., FERRI, G.L., BACARESE-HAMILTON, A.J., FUESSL, H.S., POLAK, J.M. and BLOOM, S.R., 1985. Human distribution and release of a putative new gut hormone, peptide YY. *Gastroenterology*, **89**(5), pp. 1070-1077.
- AGARWAL, A.K., SIMHA, V., ORAL, E.A., MORAN, S.A., GORDEN, P., O'RAHILLY, S., ZAIDI, Z., GURAKAN, F., ARSLANIAN, S.A., KLAR, A., RICKER, A., WHITE, N.H., BINDL, L., HERBST, K., KENNEL, K., PATEL, S.B., AL-GAZALI, L. and GARG, A., 2003. Phenotypic and genetic heterogeneity in congenital generalized lipodystrophy. *The Journal of clinical endocrinology and metabolism*, **88**(10), pp. 4840-4847.
- AGGARWAL, A., COSTA, M.J., RIVERO-GUTIERREZ, B., JI, L., MORGAN, S.L. and FELDMAN, B.J., 2017. The Circadian Clock Regulates Adipogenesis by a Per3 Crosstalk Pathway to Klf15. *Cell reports*, **21**(9), pp. 2367-2375.
- AHIMA, R.S., KELLY, J., ELMQUIST, J.K. and FLIER, J.S., 1999. Distinct physiologic and neuronal responses to decreased leptin and mild hyperleptinemia. *Endocrinology*, **140**(11), pp. 4923-4931.
- AHMADIAN, M., DUNCAN, R.E., JAWORSKI, K., SARKADI-NAGY, E. and SUL, H.S., 2007. Triacylglycerol metabolism in adipose tissue. *Future lipidology*, **2**(2), pp. 229-237.
- AHMED, M., NEVILLE, M.J., EDELMANN, M.J., KESSLER, B.M. and KARPE, F., 2010. Proteomic analysis of human adipose tissue after rosiglitazone treatment shows coordinated changes to promote glucose uptake. *Obesity (Silver Spring, Md.)*, **18**(1), pp. 27-34.
- AHREN, B., 2007. Dipeptidyl peptidase-4 inhibitors: clinical data and clinical implications. *Diabetes care*, **30**(6), pp. 1344-1350.
- AKASHEH, R.T., PINI, M., PANG, J. and FANTUZZI, G., 2013. Increased adiposity in annexin A1-deficient mice. *PloS one*, **8**(12), pp. e82608.
- ALBREKTSSEN, T. and FLECKNER, J., 2001. The transcription factor Fos-related antigen 1 is induced by thiazolidinediones during differentiation of 3T3-L1 cells. *Molecular pharmacology*, **59**(3), pp. 567-575.
- ALFADDA, A.A., BENABDELKAMEL, H., MASOOD, A., MOUSTAFA, A., SALLAM, R., BASSAS, A. and DUNCAN, M., 2013. Proteomic analysis of mature adipocytes from obese patients in relation to aging. *Experimental gerontology*, **48**(11), pp. 1196-1203.
- AL-GOBLAN, A.S., AL-ALFI, M.A. and KHAN, M.Z., 2014. Mechanism linking diabetes mellitus and obesity. *Diabetes, metabolic syndrome and obesity : targets and therapy*, **7**, pp. 587-591.

- ALI, A.T., HOCHFELD, W.E., MYBURGH, R. and PEPPER, M.S., 2013. Adipocyte and adipogenesis. *European journal of cell biology*, **92**(6-7), pp. 229-236.
- ALKHOURI, N., GORNICKA, A., BERK, M.P., THAPALIYA, S., DIXON, L.J., KASHYAP, S., SCHAUER, P.R. and FELDSTEIN, A.E., 2010. Adipocyte apoptosis, a link between obesity, insulin resistance, and hepatic steatosis. *The Journal of biological chemistry*, **285**(5), pp. 3428-3438.
- ALMGREN, M., ATKINSON, R., HE, J., HILDING, A., HAGMAN, E., WOLK, A., THORELL, A., MARCUS, C., NASLUND, E., OSTENSON, C.G., SCHALLING, M. and LAVEBRATT, C., 2012. Adenovirus-36 is associated with obesity in children and adults in Sweden as determined by rapid ELISA. *PloS one*, **7**(7), pp. e41652.
- AL-SUHAIMI, E.A. and SHEHZAD, A., 2013. Leptin, resistin and visfatin: the missing link between endocrine metabolic disorders and immunity. *European journal of medical research*, **18**(1), pp. 12-783X-18-12. eCollection 2013.
- ALVAREZ BARTOLOME, M., BORQUE, M., MARTINEZ-SARMIENTO, J., APARICIO, E., HERNANDEZ, C., CABRERIZO, L. and FERNANDEZ-REPRESA, J.A., 2002. Peptide YY secretion in morbidly obese patients before and after vertical banded gastroplasty. *Obesity Surgery*, **12**(3), pp. 324-327.
- AMANO, S.U., COHEN, J.L., VANGALA, P., TENCEROVA, M., NICOLORO, S.M., YAWE, J.C., SHEN, Y., CZECH, M.P. and AOUDI, M., 2014. Local proliferation of macrophages contributes to obesity-associated adipose tissue inflammation. *Cell metabolism*, **19**(1), pp. 162-171.
- AOUDI, M., JAGER, J., LAURENT, K., GONZALEZ, T., CORMONT, M., BINETRUY, B., LE MARCHAND-BRUSTEL, Y., TANTI, J.F. and BOST, F., 2007. p38MAP Kinase activity is required for human primary adipocyte differentiation. *FEBS letters*, **581**(29), pp. 5591-5596.
- AOUDI, M., LAURENT, K., PROT, M., LE MARCHAND-BRUSTEL, Y., BINETRUY, B. and BOST, F., 2006. Inhibition of p38MAPK increases adipogenesis from embryonic to adult stages. *Diabetes*, **55**(2), pp. 281-289.
- ARAKI, M., NAKAGAWA, Y., OISHI, A., HAN, S.I., WANG, Y., KUMAGAI, K., OHNO, H., MIZUNOE, Y., IWASAKI, H., SEKIYA, M., MATSUZAKA, T. and SHIMANO, H., 2018. The Peroxisome Proliferator-Activated Receptor alpha (PPARalpha) Agonist Pemafibrate Protects against Diet-Induced Obesity in Mice. *International journal of molecular sciences*, **19**(7), pp. 10.3390/ijms19072148.
- ARBUTHNOTT, E., 1989. Brown adipose tissue: structure and function. *The Proceedings of the Nutrition Society*, **48**(2), pp. 177-182.
- ARCEO-OLAIZ, R., ESPANA-GOMEZ, M.N., MONTALVO-HERNANDEZ, J., VELAZQUEZ-FERNANDEZ, D., PANTOJA, J.P. and HERRERA, M.F., 2008. Maximal weight loss after banded and unbanded laparoscopic Roux-en-Y gastric bypass:

a randomized controlled trial. *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*, **4**(4), pp. 507-511.

ASHBY, D. and BLOOM, S.R., 2007. Recent progress in PYY research--an update report for 8th NPY meeting. *Peptides*, **28**(2), pp. 198-202.

ASTRUP, A., GOTZSCHE, P.C., VAN DE WERKEN, K., RANNERIES, C., TOUBRO, S., RABEN, A. and BUEMANN, B., 1999. Meta-analysis of resting metabolic rate in formerly obese subjects. *The American Journal of Clinical Nutrition*, **69**(6), pp. 1117-1122.

ATGIE, C., D'ALLAIRE, F. and BUKOWIECKI, L.J., 1997. Role of beta1- and beta3-adrenoceptors in the regulation of lipolysis and thermogenesis in rat brown adipocytes. *The American Journal of Physiology*, **273**(4 Pt 1), pp. C1136-42.

ATIT, R., SGAIER, S.K., MOHAMED, O.A., TAKETO, M.M., DUFORT, D., JOYNER, A.L., NISWANDER, L. and CONLON, R.A., 2006. Beta-catenin activation is necessary and sufficient to specify the dorsal dermal fate in the mouse. *Developmental biology*, **296**(1), pp. 164-176.

ATKINSON, R.L., DHURANDHAR, N.V., ALLISON, D.B., BOWEN, R.L., ISRAEL, B.A., ALBU, J.B. and AUGUSTUS, A.S., 2005. Human adenovirus-36 is associated with increased body weight and paradoxical reduction of serum lipids. *International journal of obesity (2005)*, **29**(3), pp. 281-286.

ATTIE, A.D. and SCHERER, P.E., 2009. Adipocyte metabolism and obesity. *Journal of lipid research*, **50** Suppl(Suppl), pp. S395-9.

AUBERT, J., DESSOLIN, S., BELMONTE, N., LI, M., MCKENZIE, F.R., STACCINI, L., VILLAGEOIS, P., BARHANIN, B., VERNALLIS, A., SMITH, A.G., AILHAUD, G. and DANI, C., 1999. Leukemia inhibitory factor and its receptor promote adipocyte differentiation via the mitogen-activated protein kinase cascade. *The Journal of biological chemistry*, **274**(35), pp. 24965-24972.

AVRAM, M.M., AVRAM, A.S. and JAMES, W.D., 2007. Subcutaneous fat in normal and diseased states 3. Adipogenesis: from stem cell to fat cell. *Journal of the American Academy of Dermatology*, **56**(3), pp. 472-492.

BABBIN, B.A., JESAITIS, A.J., IVANOV, A.I., KELLY, D., LAUKOETTER, M., NAVA, P., PARKOS, C.A. and NUSRAT, A., 2007. Formyl peptide receptor-1 activation enhances intestinal epithelial cell restitution through phosphatidylinositol 3-kinase-dependent activation of Rac1 and Cdc42. *Journal of immunology (Baltimore, Md.: 1950)*, **179**(12), pp. 8112-8121.

BABBIN, B.A., LAUKOETTER, M.G., NAVA, P., KOCH, S., LEE, W.Y., CAPALDO, C.T., PEATMAN, E., SEVERSON, E.A., FLOWER, R.J., PERRETTI, M., PARKOS, C.A. and NUSRAT, A., 2008. Annexin A1 regulates intestinal mucosal injury,

inflammation, and repair. *Journal of immunology (Baltimore, Md.: 1950)*, **181**(7), pp. 5035-5044.

BABIYCHUK, E.B., MONASTYRSKAYA, K. and DRAEGER, A., 2008. Fluorescent annexin A1 reveals dynamics of ceramide platforms in living cells. *Traffic (Copenhagen, Denmark)*, **9**(10), pp. 1757-1775.

BACHMAN, E.S., DHILLON, H., ZHANG, C.Y., CINTI, S., BIANCO, A.C., KOBILKA, B.K. and LOWELL, B.B., 2002. betaAR signaling required for diet-induced thermogenesis and obesity resistance. *Science (New York, N.Y.)*, **297**(5582), pp. 843-845.

BAE, E.J., 2016. DPP-4 inhibitors in diabetic complications: role of DPP-4 beyond glucose control. *Archives of Pharmacal Research*, **39**(8), pp. 1114-1128.

BAE, J.S., PARK, S.J., HAM, I.R. and LEE, T.G., 2009. Dose dependent effects of intravitreal triamcinolone acetonide on diffuse diabetic macular edema. *Korean journal of ophthalmology : KJO*, **23**(2), pp. 80-85.

BAGGIO, L.L., HUANG, Q., BROWN, T.J. and DRUCKER, D.J., 2004. Oxyntomodulin and glucagon-like peptide-1 differentially regulate murine food intake and energy expenditure. *Gastroenterology*, **127**(2), pp. 546-558.

BAKKER, A.H., VAN DIELEN, F.M., GREVE, J.W., ADAM, J.A. and BUURMAN, W.A., 2004. Preadipocyte number in omental and subcutaneous adipose tissue of obese individuals. *Obesity research*, **12**(3), pp. 488-498.

BANERJEE, R.R., RANGWALA, S.M., SHAPIRO, J.S., RICH, A.S., RHOADES, B., QI, Y., WANG, J., RAJALA, M.W., POCAI, A., SCHERER, P.E., STEPPAN, C.M., AHIMA, R.S., OBICI, S., ROSSETTI, L. and LAZAR, M.A., 2004. Regulation of fasted blood glucose by resistin. *Science (New York, N.Y.)*, **303**(5661), pp. 1195-1198.

BANERJEE, S.S., FEINBERG, M.W., WATANABE, M., GRAY, S., HASPEL, R.L., DENKINGER, D.J., KAWAHARA, R., HAUNER, H. and JAIN, M.K., 2003. The Kruppel-like factor KLF2 inhibits peroxisome proliferator-activated receptor-gamma expression and adipogenesis. *The Journal of biological chemistry*, **278**(4), pp. 2581-2584.

BARBATELLI, G., MURANO, I., MADSEN, L., HAO, Q., JIMENEZ, M., KRISTIANSEN, K., GIACOBINO, J.P., DE MATTEIS, R. and CINTI, S., 2010. The emergence of cold-induced brown adipocytes in mouse white fat depots is determined predominantly by white to brown adipocyte transdifferentiation. *American journal of physiology. Endocrinology and metabolism*, **298**(6), pp. E1244-53.

BARINAGA, M., 1995. "Obese" protein slims mice. *Science (New York, N.Y.)*, **269**(5223), pp. 475-476.

BARNES, P.J., 1998. Anti-inflammatory actions of glucocorticoids: molecular mechanisms. *Clinical science (London, England : 1979)*, **94**(6), pp. 557-572.

- BARROSO, I., GURNELL, M., CROWLEY, V.E., AGOSTINI, M., SCHWABE, J.W., SOOS, M.A., MASLEN, G.L., WILLIAMS, T.D., LEWIS, H., SCHAFER, A.J., CHATTERJEE, V.K. and O'RAHILLY, S., 1999. Dominant negative mutations in human PPARgamma associated with severe insulin resistance, diabetes mellitus and hypertension. *Nature*, **402**(6764), pp. 880-883.
- BASTARD, J.P., CARON, M., VIDAL, H., JAN, V., AUCLAIR, M., VIGOUROUX, C., LUBOINSKI, J., LAVILLE, M., MAACHI, M., GIRARD, P.M., ROZENBAUM, W., LEVAN, P. and CAPEAU, J., 2002. Association between altered expression of adipogenic factor SREBP1 in lipotrophic adipose tissue from HIV-1-infected patients and abnormal adipocyte differentiation and insulin resistance. *Lancet (London, England)*, **359**(9311), pp. 1026-1031.
- BATTERHAM, R.L., COHEN, M.A., ELLIS, S.M., LE ROUX, C.W., WITHERS, D.J., FROST, G.S., GHATEI, M.A. and BLOOM, S.R., 2003. Inhibition of food intake in obese subjects by peptide YY3-36. *The New England journal of medicine*, **349**(10), pp. 941-948.
- BATTERHAM, R.L., COWLEY, M.A., SMALL, C.J., HERZOG, H., COHEN, M.A., DAKIN, C.L., WREN, A.M., BRYNES, A.E., LOW, M.J., GHATEI, M.A., CONE, R.D. and BLOOM, S.R., 2002. Gut hormone PYY(3-36) physiologically inhibits food intake. *Nature*, **418**(6898), pp. 650-654.
- BATTERHAM, R.L. and CUMMINGS, D.E., 2016. Mechanisms of Diabetes Improvement Following Bariatric/Metabolic Surgery. *Diabetes care*, **39**(6), pp. 893-901.
- BAYS, H.E., 2009. "Sick fat," metabolic disease, and atherosclerosis. *The American Journal of Medicine*, **122**(1 Suppl), pp. S26-37.
- BAYS, H.E., TOTH, P.P., KRIS-ETHERTON, P.M., ABATE, N., ARONNE, L.J., BROWN, W.V., GONZALEZ-CAMPOY, J.M., JONES, S.R., KUMAR, R., LA FORGE, R. and SAMUEL, V.T., 2013. Obesity, adiposity, and dyslipidemia: a consensus statement from the National Lipid Association. *Journal of clinical lipidology*, **7**(4), pp. 304-383.
- BECHMANN, L.P., HANNIVOORT, R.A., GERKEN, G., HOTAMISLIGIL, G.S., TRAUNER, M. and CANBAY, A., 2012. The interaction of hepatic lipid and glucose metabolism in liver diseases. *Journal of hepatology*, **56**(4), pp. 952-964.
- BELMONTE, N., PHILLIPS, B.W., MASSIERA, F., VILLAGEOIS, P., WDZIEKONSKI, B., SAINT-MARC, P., NICHOLS, J., AUBERT, J., SAEKI, K., I, A., NARUMIYA, S., AILHAUD, G. and DANI, C., 2001. Activation of extracellular signal-regulated kinases and CREB/ATF-1 mediate the expression of CCAAT/enhancer binding proteins beta and -delta in preadipocytes. *Molecular endocrinology (Baltimore, Md.)*, **15**(11), pp. 2037-2049.
- BENA, S., BRANCALEONE, V., WANG, J.M., PERRETTI, M. and FLOWER, R.J., 2012. Annexin A1 interaction with the FPR2/ALX receptor: identification of distinct

domains and downstream associated signaling. *The Journal of biological chemistry*, **287**(29), pp. 24690-24697.

BENAIGES, D., FLORES LE-ROUX, J.A., PEDRO-BOTET, J., CHILLARON, J.J., RENARD, M., PARRI, A., RAMON, J.M., PERA, M. and GODAY, A., 2013. Sleeve gastrectomy and Roux-en-Y gastric bypass are equally effective in correcting insulin resistance. *International journal of surgery (London, England)*, **11**(4), pp. 309-313.

BENCE, K.K., DELIBEGOVIC, M., XUE, B., GORGUN, C.Z., HOTAMISLIGIL, G.S., NEEL, B.G. and KAHN, B.B., 2006. Neuronal PTP1B regulates body weight, adiposity and leptin action. *Nature medicine*, **12**(8), pp. 917-924.

BENHAMOU, P.Y. and PENFORNIS, A., 2002. Natural history, prognosis, and management of transplantation-induced diabetes mellitus. *Diabetes & metabolism*, **28**(3), pp. 166-175.

BENITO, M., PORRAS, A., NEBRED, A.R. and SANTOS, E., 1991. Differentiation of 3T3-L1 fibroblasts to adipocytes induced by transfection of ras oncogenes. *Science (New York, N.Y.)*, **253**(5019), pp. 565-568.

BERG, A.H., COMBS, T.P., DU, X., BROWNLEE, M. and SCHERER, P.E., 2001. The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nature medicine*, **7**(8), pp. 947-953.

BERG, A.H. and SCHERER, P.E., 2005. Adipose tissue, inflammation, and cardiovascular disease. *Circulation research*, **96**(9), pp. 939-949.

BERISHA, S.Z., SERRE, D., SCHAUER, P., KASHYAP, S.R. and SMITH, J.D., 2011. Changes in whole blood gene expression in obese subjects with type 2 diabetes following bariatric surgery: a pilot study. *PloS one*, **6**(3), pp. e16729.

BERNDT, J., KOVACS, P., RUSCHKE, K., KLOTING, N., FASSHAUER, M., SCHON, M.R., KORNER, A., STUMVOLL, M. and BLUHER, M., 2007. Fatty acid synthase gene expression in human adipose tissue: association with obesity and type 2 diabetes. *Diabetologia*, **50**(7), pp. 1472-1480.

BERRY, D.C., STENESEN, D., ZEVE, D. and GRAFF, J.M., 2013. The developmental origins of adipose tissue. *Development (Cambridge, England)*, **140**(19), pp. 3939-3949.

BERTOLA, A., CIUCCI, T., ROUSSEAU, D., BOURLIER, V., DUFFAUT, C., BONNAFOUS, S., BLIN-WAKKACH, C., ANTY, R., IANNELLI, A., GUGENHEIM, J., TRAN, A., BOULOUMIE, A., GUAL, P. and WAKKACH, A., 2012. Identification of adipose tissue dendritic cells correlated with obesity-associated insulin-resistance and inducing Th17 responses in mice and patients. *Diabetes*, **61**(9), pp. 2238-2247.

BJORBAEK, C., ELMQUIST, J.K., MICHL, P., AHIMA, R.S., VAN BUEREN, A., MCCALL, A.L. and FLIER, J.S., 1998. Expression of leptin receptor isoforms in rat brain microvessels. *Endocrinology*, **139**(8), pp. 3485-3491.

- BLUHER, M., 2016. Adipose tissue inflammation: a cause or consequence of obesity-related insulin resistance? *Clinical science (London, England : 1979)*, **130**(18), pp. 1603-1614.
- BLUHER, M., 2012. Clinical relevance of adipokines. *Diabetes & metabolism journal*, **36**(5), pp. 317-327.
- BLUHER, M., 2009. Adipose tissue dysfunction in obesity. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association*, **117**(6), pp. 241-250.
- BLUMER, R.M., VAN VONDEREN, M.G., SUTINEN, J., HASSINK, E., ACKERMANS, M., VAN AGTMAEL, M.A., YKI-JARVINEN, H., DANNER, S.A., REISS, P. and SAUERWEIN, H.P., 2008. Zidovudine/lamivudine contributes to insulin resistance within 3 months of starting combination antiretroviral therapy. *AIDS (London, England)*, **22**(2), pp. 227-236.
- BODARY, P.F., WESTRICK, R.J., WICKENHEISER, K.J., SHEN, Y. and EITZMAN, D.T., 2002. Effect of leptin on arterial thrombosis following vascular injury in mice. *Jama*, **287**(13), pp. 1706-1709.
- BODEN, G., 1996. Fatty acids and insulin resistance. *Diabetes care*, **19**(4), pp. 394-395.
- BODEN, G., CHEN, X., MOZZOLI, M. and RYAN, I., 1996. Effect of fasting on serum leptin in normal human subjects. *The Journal of clinical endocrinology and metabolism*, **81**(9), pp. 3419-3423.
- BOEY, D., LIN, S., KARL, T., BALDOCK, P., LEE, N., ENRIQUEZ, R., COUZENS, M., SLACK, K., DALLMANN, R., SAINSBURY, A. and HERZOG, H., 2006. Peptide YY ablation in mice leads to the development of hyperinsulinaemia and obesity. *Diabetologia*, **49**(6), pp. 1360-1370.
- BONEN, A., TANDON, N.N., GLATZ, J.F., LUIKEN, J.J. and HEIGENHAUSER, G.J., 2006. The fatty acid transporter FAT/CD36 is upregulated in subcutaneous and visceral adipose tissues in human obesity and type 2 diabetes. *International journal of obesity (2005)*, **30**(6), pp. 877-883.
- BOROVIKOVA, L.V., IVANOVA, S., ZHANG, M., YANG, H., BOTCHKINA, G.I., WATKINS, L.R., WANG, H., ABUMRAD, N., EATON, J.W. and TRACEY, K.J., 2000. Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature*, **405**(6785), pp. 458-462.
- BOST, F., AOUADI, M., CARON, L. and BINETRUY, B., 2005. The role of MAPKs in adipocyte differentiation and obesity. *Biochimie*, **87**(1), pp. 51-56.
- BOST, F., AOUADI, M., CARON, L., EVEN, P., BELMONTE, N., PROT, M., DANI, C., HOFMAN, P., PAGES, G., POUYSSEGUR, J., LE MARCHAND-BRUSTEL, Y.

and BINETRUY, B., 2005. The extracellular signal-regulated kinase isoform ERK1 is specifically required for in vitro and in vivo adipogenesis. *Diabetes*, **54**(2), pp. 402-411.

BOUCHARD, C., TREMBLAY, A., DESPRES, J.P., NADEAU, A., LUPIEN, P.J., THERIAULT, G., DUSSAULT, J., MOORJANI, S., PINAULT, S. and FOURNIER, G., 1990. The response to long-term overfeeding in identical twins. *The New England journal of medicine*, **322**(21), pp. 1477-1482.

BOYKO, E.J., FUJIMOTO, W.Y., LEONETTI, D.L. and NEWELL-MORRIS, L., 2000. Visceral adiposity and risk of type 2 diabetes: a prospective study among Japanese Americans. *Diabetes care*, **23**(4), pp. 465-471.

BRANCALEONE, V., MITIDIERI, E., FLOWER, R.J., CIRINO, G. and PERRETTI, M., 2014. Annexin A1 mediates hydrogen sulfide properties in the control of inflammation. *The Journal of pharmacology and experimental therapeutics*, **351**(1), pp. 96-104.

BRAY, G.A., 2004. Medical consequences of obesity. *The Journal of clinical endocrinology and metabolism*, **89**(6), pp. 2583-2589.

BRETON, C., 2013. The hypothalamus-adipose axis is a key target of developmental programming by maternal nutritional manipulation. *The Journal of endocrinology*, **216**(2), pp. R19-31.

BROWN, R.,J., ARAUJO-VILAR,David, CHEUNG, P.,To, DUNGER,David, GARG,Abhimanyu, JACK,Michelle, MUNGAI, Lucy, ORAL, E.,A., PATNI,Nivedita, ROTHER, K.,I., VON SCHNURBEIN,Julia, SORKINA,Ekaterina, STANLEY,Takara, VIGOUROUX,Corinne, WABITSCH,Martin, WILLIAMS,Rachel and YORIFUJI,Tohru, 2016. The Diagnosis and Management of Lipodystrophy Syndromes: A Multi-Society Practice Guideline. *The Journal of Clinical Endocrinology & Metabolism*, **101**(12), pp. 4500-4511.

CADIGAN, K.M. and NUSSE, R., 1997. Wnt signaling: a common theme in animal development. *Genes & development*, **11**(24), pp. 3286-3305.

CAMP, H.S. and TAFURI, S.R., 1997. Regulation of peroxisome proliferator-activated receptor gamma activity by mitogen-activated protein kinase. *The Journal of biological chemistry*, **272**(16), pp. 10811-10816.

CAMPBELL, P.J., CARLSON, M.G. and NURJHAN, N., 1994. Fat metabolism in human obesity. *The American Journal of Physiology*, **266**(4 Pt 1), pp. E600-5.

CAMPFIELD, L.A., SMITH, F.J., GUISEZ, Y., DEVOS, R. and BURN, P., 1995. Recombinant mouse OB protein: evidence for a peripheral signal linking adiposity and central neural networks. *Science (New York, N.Y.)*, **269**(5223), pp. 546-549.

CANI, P.D., BIBILONI, R., KNAUF, C., WAGET, A., NEYRINCK, A.M., DELZENNE, N.M. and BURCELIN, R., 2008. Changes in gut microbiota control

metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes*, **57**(6), pp. 1470-1481.

CANNON, B. and NEDERGAARD, J., 2004. Brown adipose tissue: function and physiological significance. *Physiological Reviews*, **84**(1), pp. 277-359.

CAO, H., 2014. Adipocytokines in obesity and metabolic disease. *The Journal of endocrinology*, **220**(2), pp. T47-59.

CAO, Y., 2010. Adipose tissue angiogenesis as a therapeutic target for obesity and metabolic diseases. *Nature reviews. Drug discovery*, **9**(2), pp. 107-115.

CAO, Z., UMEK, R.M. and MCKNIGHT, S.L., 1991. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes & development*, **5**(9), pp. 1538-1552.

CARLSON, C.J., KOTERSKI, S., SCIOTTI, R.J., POCCARD, G.B. and RONDINONE, C.M., 2003. Enhanced basal activation of mitogen-activated protein kinases in adipocytes from type 2 diabetes: potential role of p38 in the downregulation of GLUT4 expression. *Diabetes*, **52**(3), pp. 634-641.

CATALAN, V., GOMEZ-AMBROSI, J., RAMIREZ, B., ROTELLAR, F., PASTOR, C., SILVA, C., RODRIGUEZ, A., GIL, M.J., CIENFUEGOS, J.A. and FRUHBECK, G., 2007. Proinflammatory cytokines in obesity: impact of type 2 diabetes mellitus and gastric bypass. *Obesity Surgery*, **17**(11), pp. 1464-1474.

CERNKOVICH, E.R., DENG, J., BOND, M.C., COMBS, T.P. and HARP, J.B., 2008. Adipose-specific disruption of signal transducer and activator of transcription 3 increases body weight and adiposity. *Endocrinology*, **149**(4), pp. 1581-1590.

CHAHIL, T.J. and GINSBERG, H.N., 2006. Diabetic dyslipidemia. *Endocrinology and metabolism clinics of North America*, **35**(3), pp. 491-510, vii-viii.

CHAN, E.K., MACKEY, M.A., SNOVER, D.C., SCHNEIDER, P.D., RUCKER, R.D., Jr, ALLEN, C.E. and BUCHWALD, H., 1984. Suppression of weight gain by glucagon in obese Zucker rats. *Experimental and molecular pathology*, **40**(3), pp. 320-327.

CHAU, Y.Y., BANDIERA, R., SERRELS, A., MARTINEZ-ESTRADA, O.M., QING, W., LEE, M., SLIGHT, J., THORNBURN, A., BERRY, R., MCHAFFIE, S., STIMSON, R.H., WALKER, B.R., CHAPULI, R.M., SCHEDL, A. and HASTIE, N., 2014. Visceral and subcutaneous fat have different origins and evidence supports a mesothelial source. *Nature cell biology*, **16**(4), pp. 367-375.

CHEN, B., LAM, K.S., WANG, Y., WU, D., LAM, M.C., SHEN, J., WONG, L., HOO, R.L., ZHANG, J. and XU, A., 2006. Hypoxia dysregulates the production of adiponectin and plasminogen activator inhibitor-1 independent of reactive oxygen species in adipocytes. *Biochemical and biophysical research communications*, **341**(2), pp. 549-556.

- CHEUK, B.L. and CHENG, S.W., 2011. Annexin A1 expression in atherosclerotic carotid plaques and its relationship with plaque characteristics. *European journal of vascular and endovascular surgery : the official journal of the European Society for Vascular Surgery*, **41**(3), pp. 364-371.
- CHO, K.W., MORRIS, D.L., DELPROPOSTO, J.L., GELETKA, L., ZAMARRON, B., MARTINEZ-SANTIBANEZ, G., MEYER, K.A., SINGER, K., O'ROURKE, R.W. and LUMENG, C.N., 2014. An MHC II-dependent activation loop between adipose tissue macrophages and CD4⁺ T cells controls obesity-induced inflammation. *Cell reports*, **9**(2), pp. 605-617.
- CHO, S.O., KIM, M.H. and KIM, H., 2018. beta-Carotene Inhibits Activation of NF-kappaB, Activator Protein-1, and STAT3 and Regulates Abnormal Expression of Some Adipokines in 3T3-L1 Adipocytes. *Journal of cancer prevention*, **23**(1), pp. 37-43.
- CHUNG, J.J., MARKIEWICZ, M.A., POLIC, B. and SHAW, A.S., 2014. Role of NKG2D in obesity-induced adipose tissue inflammation and insulin resistance. *PloS one*, **9**(10), pp. e110108.
- CIANFLONE, K., XIA, Z. and CHEN, L.Y., 2003. Critical review of acylation-stimulating protein physiology in humans and rodents. *Biochimica et biophysica acta*, **1609**(2), pp. 127-143.
- CILDIR, G., AKINCILAR, S.C. and TERGAONKAR, V., 2013. Chronic adipose tissue inflammation: all immune cells on the stage. *Trends in molecular medicine*, **19**(8), pp. 487-500.
- CINTI S., 2015. The Adipose Organ: Implications For Prevention And Treatment Of Obesity. In M.L. Frelut (Ed.), *The ECOG's eBook on Child and Adolescent Obesity*. Retrieved from ebook.ecogobesity.eu
- CLARKE, B.A. and CLARKE, S.D., 1982. Polymer-protomer transition of acetyl-CoA carboxylase as a regulator of lipogenesis in rat liver. *Archives of Biochemistry and Biophysics*, **218**(1), pp. 92-100.
- CLARKE, S.F., MURPHY, E.F., NILAWEERA, K., ROSS, P.R., SHANAHAN, F., O'TOOLE, P.W. and COTTER, P.D., 2012. The gut microbiota and its relationship to diet and obesity: new insights. *Gut microbes*, **3**(3), pp. 186-202.
- COBURN, C.T., KNAPP, F.F., Jr, FEBBRAIO, M., BEETS, A.L., SILVERSTEIN, R.L. and ABUMRAD, N.A., 2000. Defective uptake and utilization of long chain fatty acids in muscle and adipose tissues of CD36 knockout mice. *The Journal of biological chemistry*, **275**(42), pp. 32523-32529.
- COHEN, M.A., ELLIS, S.M., LE ROUX, C.W., BATTERHAM, R.L., PARK, A., PATTERSON, M., FROST, G.S., GHATEI, M.A. and BLOOM, S.R., 2003. Oxyntomodulin suppresses appetite and reduces food intake in humans. *The Journal of clinical endocrinology and metabolism*, **88**(10), pp. 4696-4701.

- COLQUITT, J.L., PICOT, J., LOVEMAN, E. and CLEGG, A.J., 2009. Surgery for obesity. *The Cochrane database of systematic reviews*, (2):CD003641. doi(2), pp. CD003641.
- COORAY, S.N., GOBBETTI, T., MONTERO-MELENDZ, T., MCARTHUR, S., THOMPSON, D., CLARK, A.J., FLOWER, R.J. and PERRETTI, M., 2013. Ligand-specific conformational change of the G-protein-coupled receptor ALX/FPR2 determines proresolving functional responses. *Proceedings of the National Academy of Sciences of the United States of America*, **110**(45), pp. 18232-18237.
- COSTA, M.J., SO, A.Y., KAASIK, K., KRUEGER, K.C., PILLSBURY, M.L., FU, Y.H., PTACEK, L.J., YAMAMOTO, K.R. and FELDMAN, B.J., 2011. Circadian rhythm gene period 3 is an inhibitor of the adipocyte cell fate. *The Journal of biological chemistry*, **286**(11), pp. 9063-9070.
- COUPER, K.N., BLOUNT, D.G. and RILEY, E.M., 2008. IL-10: the master regulator of immunity to infection. *Journal of immunology (Baltimore, Md.: 1950)*, **180**(9), pp. 5771-5777.
- CROFTS, C., WHELDON, M., ZINN, C., LAN-PIDHAINY, X., WOLEVER, T., SCHOFIELD, G., 2017. Assessing the test-retest repeatability of insulin resistance measures: Homeostasis model assessment 2 and oral glucose insulin sensitivity. *Journal of Insulin Resistance*, **2**(1),.
- CROXTALL, J.D., GILROY, D.W., SOLITO, E., CHOUDHURY, Q., WARD, B.J., BUCKINGHAM, J.C. and FLOWER, R.J., 2003. Attenuation of glucocorticoid functions in an Anx-A1-/- cell line. *The Biochemical journal*, **371**(Pt 3), pp. 927-935.
- D'ACQUISTO, F., MERGHANI, A., LECONA, E., ROSIGNOLI, G., RAZA, K., BUCKLEY, C.D., FLOWER, R.J. and PERRETTI, M., 2007. Annexin-1 modulates T-cell activation and differentiation. *Blood*, **109**(3), pp. 1095-1102.
- DALLI, J., MONTERO-MELENDZ, T., MCARTHUR, S. and PERRETTI, M., 2012. Annexin A1 N-terminal derived Peptide ac2-26 exerts chemokinetic effects on human neutrophils. *Frontiers in pharmacology*, **3**, pp. 28.
- DAMAZO, A.S., YONA, S., FLOWER, R.J., PERRETTI, M. and OLIANI, S.M., 2006. Spatial and temporal profiles for anti-inflammatory gene expression in leukocytes during a resolving model of peritonitis. *Journal of immunology (Baltimore, Md.: 1950)*, **176**(7), pp. 4410-4418.
- DANFORTH, E., Jr, 2000. Failure of adipocyte differentiation causes type II diabetes mellitus? *Nature genetics*, **26**(1), pp. 13.
- DANKEL, S.N., DEGERUD, E.M., BORKOWSKI, K., FJAERE, E., MIDTBO, L.K., HAUGEN, C., SOLSVIK, M.H., LAVIGNE, A.M., LIASET, B., SAGEN, J.V., KRISTIANSEN, K., MELLGREN, G. and MADSEN, L., 2014. Weight cycling

promotes fat gain and altered clock gene expression in adipose tissue in C57BL/6J mice. *American journal of physiology. Endocrinology and metabolism*, **306**(2), pp. E210-24.

DAQUINAG, A.C., ZHANG, Y. and KOLONIN, M.G., 2011. Vascular targeting of adipose tissue as an anti-obesity approach. *Trends in pharmacological sciences*, **32**(5), pp. 300-307.

DARDENNE, M., SAVINO, W., GASTINEL, L.N., NABARRA, B. and BACH, J.F., 1983. Thymic dysfunction in the mutant diabetic (db/db) mouse. *Journal of immunology (Baltimore, Md.: 1950)*, **130**(3), pp. 1195-1199.

DARNELL, J.E., Jr, 1997. STATs and gene regulation. *Science (New York, N.Y.)*, **277**(5332), pp. 1630-1635.

DAS, U.N., 2007. Acetylcholinesterase and butyrylcholinesterase as possible markers of low-grade systemic inflammation. *Medical science monitor : international medical journal of experimental and clinical research*, **13**(12), pp. RA214-21.

DASH, S., XIAO, C. and LEWIS, G.F., 2016. Effects of bariatric surgery on hepatic and intestinal lipoprotein particle metabolism. *Current opinion in lipidology*, **27**(1), pp. 14-18.

DE FERRANTI, S. and MOZAFFARIAN, D., 2008. The perfect storm: obesity, adipocyte dysfunction, and metabolic consequences. *Clinical chemistry*, **54**(6), pp. 945-955.

DE JONGE, L. and BRAY, G.A., 1997. The thermic effect of food and obesity: a critical review. *Obesity research*, **5**(6), pp. 622-631.

DE LUIS, D.A., GONZALEZ SAGRADO, M., CONDE, R., ALLER, R. and IZAOLA, O., 2010. Resistin levels and inflammatory markers in patients with morbid obesity. *Nutricion hospitalaria*, **25**(4), pp. 630-634.

DE MINICIS, S., AGOSTINELLI, L., RYCHLICKI, C., SORICE, G.P., SACCOMANNO, S., CANDELARESI, C., GIACCARI, A., TROZZI, L., PIERANTONELLI, I., MINGARELLI, E., MARZIONI, M., MUSCOGIURI, G., GAGGINI, M., BENEDETTI, A., GASTALDELLI, A., GUIDO, M. and SVEGLIATI-BARONI, G., 2014. HCC development is associated to peripheral insulin resistance in a mouse model of NASH. *PloS one*, **9**(5), pp. e97136.

DEACON, C.F., 2019. Physiology and Pharmacology of DPP-4 in Glucose Homeostasis and the Treatment of Type 2 Diabetes. *Frontiers in endocrinology*, **10**, pp. 80.

DELIKAT, S., HARRIS, R.J. and GALVANI, D.W., 1993. IL-1 beta inhibits adipocyte formation in human long-term bone marrow culture. *Experimental hematology*, **21**(1), pp. 31-37.

- DEMARIA, E.J. and JAMAL, M.K., 2005. Surgical options for obesity. *Gastroenterology clinics of North America*, **34**(1), pp. 127-142.
- DEN BROEDER, M.J., KOPYLOVA, V.A., KAMMINGA, L.M. and LEGLER, J., 2015. Zebrafish as a Model to Study the Role of Peroxisome Proliferating-Activated Receptors in Adipogenesis and Obesity. *PPAR research*, **2015**, pp. 358029.
- DENG, J., HUA, K., CAVENEY, E.J., TAKAHASHI, N. and HARP, J.B., 2006. Protein inhibitor of activated STAT3 inhibits adipogenic gene expression. *Biochemical and biophysical research communications*, **339**(3), pp. 923-931.
- DIMITRIADIS, G., MITROU, P., LAMBADIARI, V., MARATOU, E. and RAPTIS, S.A., 2011. Insulin effects in muscle and adipose tissue. *Diabetes research and clinical practice*, **93 Suppl 1**, pp. S52-9.
- DIRCKS, L. and SUL, H.S., 1999. Acyltransferases of de novo glycerophospholipid biosynthesis. *Progress in lipid research*, **38**(5-6), pp. 461-479.
- DIVELLA, R., DE LUCA, R., ABBATE, I., NAGLIERI, E. and DANIELE, A., 2016. Obesity and cancer: the role of adipose tissue and adipo-cytokines-induced chronic inflammation. *Journal of Cancer*, **7**(15), pp. 2346-2359.
- DIVI, R.L., HAVERKOS, K.J., HUMSI, J.A., SHOCKLEY, M.E., THAMIRE, C., NAGASHIMA, K., OLIVERO, O.A. and POIRIER, M.C., 2007. Morphological and molecular course of mitochondrial pathology in cultured human cells exposed long-term to Zidovudine. *Environmental and molecular mutagenesis*, **48**(3-4), pp. 179-189.
- DOCANTO, M.M., HAM, S., CORBOULD, A. and BROWN, K.A., 2015. Obesity-Associated Inflammatory Cytokines and Prostaglandin E2 Stimulate Glucose Transporter mRNA Expression and Glucose Uptake in Primary Human Adipose Stromal Cells. *Journal of interferon & cytokine research : the official journal of the International Society for Interferon and Cytokine Research*, **35**(8), pp. 600-605.
- DOMINGUEZ, H., STORGAARD, H., RASK-MADSEN, C., STEFFEN HERMANN, T., IHLEMANN, N., BAUNBJERG NIELSEN, D., SPOHR, C., KOBER, L., VAAG, A. and TORP-PEDERSEN, C., 2005. Metabolic and vascular effects of tumor necrosis factor-alpha blockade with etanercept in obese patients with type 2 diabetes. *Journal of vascular research*, **42**(6), pp. 517-525.
- DONEY, A.S., FISCHER, B., LEESE, G., MORRIS, A.D. and PALMER, C.N., 2004. Cardiovascular risk in type 2 diabetes is associated with variation at the PPARG locus: a Go-DARTS study. *Arteriosclerosis, Thrombosis, and Vascular Biology*, **24**(12), pp. 2403-2407.
- DRECHSLER, M., DE JONG, R., ROSSAINT, J., VIOLA, J.R., LEONI, G., WANG, J.M., GROMMES, J., HINKEL, R., KUPATT, C., WEBER, C., DORING, Y., ZARBOCK, A. and SOEHNLEIN, O., 2015. Annexin A1 counteracts chemokine-induced arterial myeloid cell recruitment. *Circulation research*, **116**(5), pp. 827-835.

- DRUCKER, D.J. and NAUCK, M.A., 2006. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet (London, England)*, **368**(9548), pp. 1696-1705.
- DUFTON, N. and PERRETTI, M., 2010. Therapeutic anti-inflammatory potential of formyl-peptide receptor agonists. *Pharmacology & therapeutics*, **127**(2), pp. 175-188.
- DUNCAN, R.E., AHMADIAN, M., JAWORSKI, K., SARKADI-NAGY, E. and SUL, H.S., 2007. Regulation of lipolysis in adipocytes. *Annual Review of Nutrition*, **27**, pp. 79-101.
- EDER, K., BAFFY, N., FALUS, A. and FULOP, A.K., 2009. The major inflammatory mediator interleukin-6 and obesity. *Inflammation research : official journal of the European Histamine Research Society ...[et al.]*, **58**(11), pp. 727-736.
- EFERL, R., RICCI, R., KENNER, L., ZENZ, R., DAVID, J.P., RATH, M. and WAGNER, E.F., 2003. Liver tumor development. c-Jun antagonizes the proapoptotic activity of p53. *Cell*, **112**(2), pp. 181-192.
- EGAN, J.J., GREENBERG, A.S., CHANG, M.K., WEK, S.A., MOOS, M.C., Jr and LONDOS, C., 1992. Mechanism of hormone-stimulated lipolysis in adipocytes: translocation of hormone-sensitive lipase to the lipid storage droplet. *Proceedings of the National Academy of Sciences of the United States of America*, **89**(18), pp. 8537-8541.
- EL KEBIR, D., JOZSEF, L. and FILEP, J.G., 2008. Opposing regulation of neutrophil apoptosis through the formyl peptide receptor-like 1/lipoxin A4 receptor: implications for resolution of inflammation. *Journal of leukocyte biology*, **84**(3), pp. 600-606.
- ELGAZAR-CARMON, V., RUDICH, A., HADAD, N. and LEVY, R., 2008. Neutrophils transiently infiltrate intra-abdominal fat early in the course of high-fat feeding. *Journal of lipid research*, **49**(9), pp. 1894-1903.
- ELLINGSGAARD, H., HAUSELMANN, I., SCHULER, B., HABIB, A.M., BAGGIO, L.L., MEIER, D.T., EPPLER, E., BOUZAKRI, K., WUEEST, S., MULLER, Y.D., HANSEN, A.M., REINECKE, M., KONRAD, D., GASSMANN, M., REIMANN, F., HALBAN, P.A., GROMADA, J., DRUCKER, D.J., GRIBBLE, F.M., EHSES, J.A. and DONATH, M.Y., 2011. Interleukin-6 enhances insulin secretion by increasing glucagon-like peptide-1 secretion from L cells and alpha cells. *Nature medicine*, **17**(11), pp. 1481-1489.
- EL-MESALLAMY, H.O., KASSEM, D.H., EL-DEMERDASH, E. and AMIN, A.I., 2011. Vaspin and visfatin/Nampt are interesting interrelated adipokines playing a role in the pathogenesis of type 2 diabetes mellitus. *Metabolism: clinical and experimental*, **60**(1), pp. 63-70.
- ELMORE, S., 2007. Apoptosis: a review of programmed cell death. *Toxicologic pathology*, **35**(4), pp. 495-516.

ELMQUIST, J.K., ELIAS, C.F. and SAPER, C.B., 1999. From lesions to leptin: hypothalamic control of food intake and body weight. *Neuron*, **22**(2), pp. 221-232.

EL-OMAR, E.M., 2001. The importance of interleukin 1beta in Helicobacter pylori associated disease. *Gut*, **48**(6), pp. 743-747.

ENGELMAN, J.A., BERG, A.H., LEWIS, R.Y., LIN, A., LISANTI, M.P. and SCHERER, P.E., 1999. Constitutively active mitogen-activated protein kinase kinase 6 (MKK6) or salicylate induces spontaneous 3T3-L1 adipogenesis. *The Journal of biological chemistry*, **274**(50), pp. 35630-35638.

ENGELMAN, J.A., LISANTI, M.P. and SCHERER, P.E., 1998. Specific inhibitors of p38 mitogen-activated protein kinase block 3T3-L1 adipogenesis. *The Journal of biological chemistry*, **273**(48), pp. 32111-32120.

ENGIN, A., 2017a. Adipose Tissue Hypoxia in Obesity and Its Impact on Preadipocytes and Macrophages: Hypoxia Hypothesis. *Advances in Experimental Medicine and Biology*, **960**, pp. 305-326.

ENGIN, A., 2017b. Fat Cell and Fatty Acid Turnover in Obesity. *Advances in Experimental Medicine and Biology*, **960**, pp. 135-160.

ERNST, S., LANGE, C., WILBERS, A., GOEBELER, V., GERKE, V. and RESCHER, U., 2004. An annexin 1 N-terminal peptide activates leukocytes by triggering different members of the formyl peptide receptor family. *Journal of immunology (Baltimore, Md.: 1950)*, **172**(12), pp. 7669-7676.

ESCHER, P., BRAISSANT, O., BASU-MODAK, S., MICHALIK, L., WAHLI, W. and DESVERGNE, B., 2001. Rat PPARs: quantitative analysis in adult rat tissues and regulation in fasting and refeeding. *Endocrinology*, **142**(10), pp. 4195-4202.

FARIA, G., PESTANA, D., ARAL, M., PRETO, J., NORBERTO, S., CALHAU, C., GUIMARAES, J.T. and TAVEIRA-GOMES, A., 2014. Metabolic score: insights on the development and prediction of remission of metabolic syndrome after gastric bypass. *Annals of Surgery*, **260**(2), pp. 279-286.

FAROOQI, I.S., JEBB, S.A., LANGMACK, G., LAWRENCE, E., CHEETHAM, C.H., PRENTICE, A.M., HUGHES, I.A., MCCAMISH, M.A. and O'RAHILLY, S., 1999. Effects of recombinant leptin therapy in a child with congenital leptin deficiency. *The New England journal of medicine*, **341**(12), pp. 879-884.

FAROOQI, I.S., MATARESE, G., LORD, G.M., KEOGH, J.M., LAWRENCE, E., AGWU, C., SANNA, V., JEBB, S.A., PERNA, F., FONTANA, S., LECHLER, R.I., DEPAOLI, A.M. and O'RAHILLY, S., 2002. Beneficial effects of leptin on obesity, T cell hyporesponsiveness, and neuroendocrine/metabolic dysfunction of human congenital leptin deficiency. *The Journal of clinical investigation*, **110**(8), pp. 1093-1103.

- FAROOQI, S. and O'RAHILLY, S., 2006. Genetics of obesity in humans. *Endocrine reviews*, **27**(7), pp. 710-718.
- FATIMA, L.A., CAMPELLO, R.S., SANTOS, R.S., FREITAS, H.S., FRANK, A.P., MACHADO, U.F. and CLEGG, D.J., 2017. Estrogen receptor 1 (ESR1) regulates VEGFA in adipose tissue. *Scientific reports*, **7**(1), pp. 16716-017-16686-7.
- FEMLAK, M., GLUBA-BRZĄZKA, A., CIAŁKOŃSKA-RYSZ, A. and RYSZ, J., 2017. The role and function of HDL in patients with diabetes mellitus and the related cardiovascular risk. *Lipids in health and disease*, **16**(1), pp. 207-017-0594-3. eCollection 2017.
- FENG, J., WANG, X., LI, H., WANG, L. and TANG, Z., 2018. Silencing of Annexin A1 suppressed the apoptosis and inflammatory response of preeclampsia rat trophoblasts. *International journal of molecular medicine*, .
- FERLAZZO, V., D'AGOSTINO, P., MILANO, S., CARUSO, R., FEO, S., CILLARI, E. and PARENTE, L., 2003. Anti-inflammatory effects of annexin-1: stimulation of IL-10 release and inhibition of nitric oxide synthesis. *International immunopharmacology*, **3**(10-11), pp. 1363-1369.
- FERRANNINI, E. and MINGRONE, G., 2009. Impact of different bariatric surgical procedures on insulin action and beta-cell function in type 2 diabetes. *Diabetes care*, **32**(3), pp. 514-520.
- FERRUCCI, L. and FABBRI, E., 2018. Inflammageing: chronic inflammation in ageing, cardiovascular disease, and frailty. *Nature reviews.Cardiology*, **15**(9), pp. 505-522.
- FEVE, B. and BASTARD, J.P., 2009. The role of interleukins in insulin resistance and type 2 diabetes mellitus. *Nature reviews.Endocrinology*, **5**(6), pp. 305-311.
- FIELD, J.B., 1989. Exercise and deficient carbohydrate storage and intake as causes of hypoglycemia. *Endocrinology and metabolism clinics of North America*, **18**(1), pp. 155-161.
- FILEP, J.G., 2013. Biasing the lipoxin A4/formyl peptide receptor 2 pushes inflammatory resolution. *Proceedings of the National Academy of Sciences of the United States of America*, **110**(45), pp. 18033-18034.
- FIORENZA, C.G., CHOU, S.H. and MANTZOROS, C.S., 2011. Lipodystrophy: pathophysiology and advances in treatment. *Nature reviews.Endocrinology*, **7**(3), pp. 137-150.
- FISCHER-POSOVSZKY, P., NEWELL, F.S., WABITSCH, M. and TORNQVIST, H.E., 2008. Human SGBS cells - a unique tool for studies of human fat cell biology. *Obesity facts*, **1**(4), pp. 184-189.

- FISHER, J.S., 2006. Potential Role of the AMP-activated Protein Kinase in Regulation of Insulin Action. *Cellscience*, **2**(3), pp. 68-81.
- FLIER, J.S., 1998. Clinical review 94: What's in a name? In search of leptin's physiologic role. *The Journal of clinical endocrinology and metabolism*, **83**(5), pp. 1407-1413.
- FLINT, A., RABEN, A., ASTRUP, A. and HOLST, J.J., 1998. Glucagon-like peptide 1 promotes satiety and suppresses energy intake in humans. *The Journal of clinical investigation*, **101**(3), pp. 515-520.
- FLODBY, P., BARLOW, C., KYLEFJORD, H., AHRLUND-RICHTER, L. and XANTHOPOULOS, K.G., 1996. Increased hepatic cell proliferation and lung abnormalities in mice deficient in CCAAT/enhancer binding protein alpha. *The Journal of biological chemistry*, **271**(40), pp. 24753-24760.
- FLOWER, R.J. and ROTHWELL, N.J., 1994. Lipocortin-1: cellular mechanisms and clinical relevance. *Trends in pharmacological sciences*, **15**(3), pp. 71-76.
- FOWKES, R.C., MORADI-BIDHENDI, N., BRANCALEONE, V., ZARIWALA, M.G., BRADY, D., JESSOP, D.S., PERRETTI, M. and RENSHAW, D., 2013. Annexin-A1 protein and its relationship to cortisol in human saliva. *Psychoneuroendocrinology*, **38**(5), pp. 722-727.
- FRAYN, K.N., ARNER, P. and YKI-JARVINEN, H., 2006. Fatty acid metabolism in adipose tissue, muscle and liver in health and disease. *Essays in biochemistry*, **42**, pp. 89-103.
- FRIED, M., 2013. Bariatric and metabolic surgery. *Minerva endocrinologica*, **38**(3), pp. 237-244.
- FRIED, S.K., BUNKIN, D.A. and GREENBERG, A.S., 1998. Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *The Journal of clinical endocrinology and metabolism*, **83**(3), pp. 847-850.
- FRIEDMAN, J.M. and HALAAS, J.L., 1998. Leptin and the regulation of body weight in mammals. *Nature*, **395**(6704), pp. 763-770.
- FRIKKE-SCHMIDT, H., O'ROURKE, R.W., LUMENG, C.N., SANDOVAL, D.A. and SEELEY, R.J., 2016. Does bariatric surgery improve adipose tissue function? *Obesity reviews : an official journal of the International Association for the Study of Obesity*, **17**(9), pp. 795-809.
- FROY, O., 2012. Circadian rhythms and obesity in mammals. *ISRN obesity*, **2012**, pp. 437198.

- FU, Y., LUO, N., KLEIN, R.L. and GARVEY, W.T., 2005. Adiponectin promotes adipocyte differentiation, insulin sensitivity, and lipid accumulation. *Journal of lipid research*, **46**(7), pp. 1369-1379.
- FU, Z., GILBERT, E.R. and LIU, D., 2013. Regulation of Insulin Synthesis and Secretion and Pancreatic Beta-Cell Dysfunction in Diabetes. *Current diabetes reviews*, **9**(1), pp. 25-53.
- FUJIMOTO, S., MOCHIZUKI, K., SHIMADA, M., HORI, T., MURAYAMA, Y., OHASHI, N. and GODA, T., 2010. Insulin resistance induced by a high-fat diet is associated with the induction of genes related to leukocyte activation in rat peripheral leukocytes. *Life Sciences*, **87**(23-26), pp. 679-685.
- FUKUHARA, A., MATSUDA, M., NISHIZAWA, M., SEGAWA, K., TANAKA, M., KISHIMOTO, K., MATSUKI, Y., MURAKAMI, M., ICHISAKA, T., MURAKAMI, H., WATANABE, E., TAKAGI, T., AKIYOSHI, M., OHTSUBO, T., KIHARA, S., YAMASHITA, S., MAKISHIMA, M., FUNAHASHI, T., YAMANAKA, S., HIRAMATSU, R., MATSUZAWA, Y. and SHIMOMURA, I., 2005. Visfatin: a protein secreted by visceral fat that mimics the effects of insulin. *Science (New York, N.Y.)*, **307**(5708), pp. 426-430.
- FURUKAWA, S., FUJITA, T., SHIMABUKURO, M., IWAKI, M., YAMADA, Y., NAKAJIMA, Y., NAKAYAMA, O., MAKISHIMA, M., MATSUDA, M. and SHIMOMURA, I., 2004. Increased oxidative stress in obesity and its impact on metabolic syndrome. *The Journal of clinical investigation*, **114**(12), pp. 1752-1761.
- GALANAKIS, C.G., DASKALAKIS, M., MANIOS, A., XYDA, A., KARANTANAS, A.H. and MELISSAS, J., 2015. Computed tomography-based assessment of abdominal adiposity changes and their impact on metabolic alterations following bariatric surgery. *World journal of surgery*, **39**(2), pp. 417-423.
- GALLAGHER, D., KELLEY, D.E., YIM, J.E., SPENCE, N., ALBU, J., BOXT, L., PI-SUNYER, F.X., HESHKA, S. and MRI ANCILLARY STUDY GROUP OF THE LOOK AHEAD RESEARCH GROUP, 2009. Adipose tissue distribution is different in type 2 diabetes. *The American Journal of Clinical Nutrition*, **89**(3), pp. 807-814.
- GARG, A., 2004. Acquired and inherited lipodystrophies. *The New England journal of medicine*, **350**(12), pp. 1220-1234.
- GARG, A., 2000. Gender differences in the prevalence of metabolic complications in familial partial lipodystrophy (Dunnigan variety). *The Journal of clinical endocrinology and metabolism*, **85**(5), pp. 1776-1782.
- GARRIDO-SANCHEZ, L., VENDRELL, J., FERNANDEZ-GARCIA, D., CEPERUELO-MALLAFRE, V., CHACON, M.R., OCANA-WILHELMI, L., ALCAIDE, J., TINAHONES, F.J. and GARCIA-FUENTES, E., 2012. De novo lipogenesis in adipose tissue is associated with course of morbid obesity after bariatric surgery. *PloS one*, **7**(2), pp. e31280.

- GARVEY, W.T., MAIANU, L., HUECKSTEADT, T.P., BIRNBAUM, M.J., MOLINA, J.M. and CIARALDI, T.P., 1991. Pretranslational suppression of a glucose transporter protein causes insulin resistance in adipocytes from patients with non-insulin-dependent diabetes mellitus and obesity. *The Journal of clinical investigation*, **87**(3), pp. 1072-1081.
- GASTALDELLI, A., 2011. Role of beta-cell dysfunction, ectopic fat accumulation and insulin resistance in the pathogenesis of type 2 diabetes mellitus. *Diabetes research and clinical practice*, **93 Suppl 1**, pp. S60-5.
- GAVINS, F.N. and HICKEY, M.J., 2012. Annexin A1 and the regulation of innate and adaptive immunity. *Frontiers in immunology*, **3**, pp. 354.
- GAVRILOVA, O., MARCUS-SAMUELS, B., GRAHAM, D., KIM, J.K., SHULMAN, G.I., CASTLE, A.L., VINSON, C., ECKHAUS, M. and REITMAN, M.L., 2000. Surgical implantation of adipose tissue reverses diabetes in lipoatrophic mice. *The Journal of clinical investigation*, **105**(3), pp. 271-278.
- GEARY, N., LE SAUTER, J. and NOH, U., 1993. Glucagon acts in the liver to control spontaneous meal size in rats. *The American Journal of Physiology*, **264**(1 Pt 2), pp. R116-22.
- GILL, J.A. and LA MERRILL, M.A., 2017. An emerging role for epigenetic regulation of Pgc-1 β expression in environmentally stimulated brown adipose thermogenesis. *Environmental epigenetics*, **3**(2), pp. dvx009.
- GLASS, C.K. and ROSENFELD, M.G., 2000. The coregulator exchange in transcriptional functions of nuclear receptors. *Genes & development*, **14**(2), pp. 121-141.
- GLOY, V.L., BRIEL, M., BHATT, D.L., KASHYAP, S.R., SCHAUER, P.R., MINGRONE, G., BUCHER, H.C. and NORDMANN, A.J., 2013. Bariatric surgery versus non-surgical treatment for obesity: a systematic review and meta-analysis of randomised controlled trials. *BMJ (Clinical research ed.)*, **347**, pp. f5934.
- GODSLAND, I.F., 2009. Insulin resistance and hyperinsulinaemia in the development and progression of cancer. *Clinical science (London, England : 1979)*, **118**(5), pp. 315-332.
- GOKTAS, Z., MOUSTAID-MOUSSA, N., SHEN, C.L., BOYLAN, M., MO, H. and WANG, S., 2013. Effects of bariatric surgery on adipokine-induced inflammation and insulin resistance. *Frontiers in endocrinology*, **4**, pp. 69.
- GOOSSENS, G.H., 2008. The role of adipose tissue dysfunction in the pathogenesis of obesity-related insulin resistance. *Physiology & Behavior*, **94**(2), pp. 206-218.
- GOUDRIAAN, J.R., DEN BOER, M.A., RENSEN, P.C., FEBBRAIO, M., KUIPERS, F., ROMIJN, J.A., HAVEKES, L.M. and VOSHOL, P.J., 2005. CD36 deficiency in mice impairs lipoprotein lipase-mediated triglyceride clearance. *Journal of lipid research*, **46**(10), pp. 2175-2181.

- GREENBERG, A.S. and OBIN, M.S., 2006. Obesity and the role of adipose tissue in inflammation and metabolism. *The American Journal of Clinical Nutrition*, **83**(2), pp. 461S-465S.
- GREGOR, M.F. and HOTAMISLIGIL, G.S., 2007. Thematic review series: Adipocyte Biology. Adipocyte stress: the endoplasmic reticulum and metabolic disease. *Journal of lipid research*, **48**(9), pp. 1905-1914.
- GRIESEL, B.A., WEEMS, J., RUSSELL, R.A., ABEL, E.D., HUMPHRIES, K. and OLSON, A.L., 2010. Acute inhibition of fatty acid import inhibits GLUT4 transcription in adipose tissue, but not skeletal or cardiac muscle tissue, partly through liver X receptor (LXR) signaling. *Diabetes*, **59**(4), pp. 800-807.
- GRONG, E., ARBO, I.B., THU, O.K., KUHRY, E., KULSENG, B. and MARVIK, R., 2015. The effect of duodenojejunoscopy and sleeve gastrectomy on type 2 diabetes mellitus and gastrin secretion in Goto-Kakizaki rats. *Surgical endoscopy*, **29**(3), pp. 723-733.
- GROSS, P.M., 1992. Circumventricular organ capillaries. *Progress in brain research*, **91**, pp. 219-233.
- GUAL, P., LE MARCHAND-BRUSTEL, Y. and TANTI, J.F., 2005. Positive and negative regulation of insulin signaling through IRS-1 phosphorylation. *Biochimie*, **87**(1), pp. 99-109.
- GUERRE-MILLO, M., GERVOIS, P., RASPE, E., MADSEN, L., POULAIN, P., DERUDAS, B., HERBERT, J.M., WINEGAR, D.A., WILLSON, T.M., FRUCHAR, J.C., BERGE, R.K. and STAEELS, B., 2000. Peroxisome proliferator-activated receptor alpha activators improve insulin sensitivity and reduce adiposity. *The Journal of biological chemistry*, **275**(22), pp. 16638-16642.
- GUNNETT, C.A., HEISTAD, D.D. and FARACI, F.M., 2002. Interleukin-10 protects nitric oxide-dependent relaxation during diabetes: role of superoxide. *Diabetes*, **51**(6), pp. 1931-1937.
- GUPTA, R.K., ARANY, Z., SEALE, P., MEPANI, R.J., YE, L., CONROE, H.M., ROBY, Y.A., KULAGA, H., REED, R.R. and SPIEGELMAN, B.M., 2010. Transcriptional control of preadipocyte determination by Zfp423. *Nature*, **464**(7288), pp. 619-623.
- GUPTA, R.K., MEPANI, R.J., KLEINER, S., LO, J.C., KHANDEKAR, M.J., COHEN, P., FRONTINI, A., BHOWMICK, D.C., YE, L., CINTI, S. and SPIEGELMAN, B.M., 2012. Zfp423 expression identifies committed preadipocytes and localizes to adipose endothelial and perivascular cells. *Cell metabolism*, **15**(2), pp. 230-239.
- GURSOY CALAN, O., CALAN, M., YESIL SENSES, P., UNAL KOCABAS, G., OZDEN, E., SARI, K.R., KOCAR, M., IMAMOGLU, C., SENSES, Y.M., BOZKAYA, G. and BILGIR, O., 2016. Increased adiponin is associated with carotid intima media

thickness and metabolic disturbances in polycystic ovary syndrome. *Clinical endocrinology*, **85**(6), pp. 910-917.

GUSTAFSON, B., GOGG, S., HEDJAZIFAR, S., JENND AHL, L., HAMMARSTEDT, A. and SMITH, U., 2009. Inflammation and impaired adipogenesis in hypertrophic obesity in man. *American journal of physiology. Endocrinology and metabolism*, **297**(5), pp. E999-E1003.

GUTZWILLER, J.P., DREWE, J., GOKE, B., SCHMIDT, H., ROHRER, B., LAREIDA, J. and BEGLINGER, C., 1999. Glucagon-like peptide-1 promotes satiety and reduces food intake in patients with diabetes mellitus type 2. *The American Journal of Physiology*, **276**(5 Pt 2), pp. R1541-4.

HAASE, J., WEYER, U., IMMIG, K., KLOTING, N., BLUHER, M., EILERS, J., BECHMANN, I. and GERICKE, M., 2014. Local proliferation of macrophages in adipose tissue during obesity-induced inflammation. *Diabetologia*, **57**(3), pp. 562-571.

HAEMMERLE, G., ZIMMERMANN, R., HAYN, M., THEUSSL, C., WAEG, G., WAGNER, E., SATTler, W., MAGIN, T.M., WAGNER, E.F. and ZECHNER, R., 2002. Hormone-sensitive lipase deficiency in mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *The Journal of biological chemistry*, **277**(7), pp. 4806-4815.

HAFIDA, S., MIRSHAHI, T. and NIKOLAJCZYK, B.S., 2016. The impact of bariatric surgery on inflammation: quenching the fire of obesity? *Current opinion in endocrinology, diabetes, and obesity*, **23**(5), pp. 373-378.

HALAZONETIS, T.D., GEORGOPOULOS, K., GREENBERG, M.E. and LEDER, P., 1988. c-Jun dimerizes with itself and with c-Fos, forming complexes of different DNA binding affinities. *Cell*, **55**(5), pp. 917-924.

HALBERG, N., KHAN, T., TRUJILLO, M.E., WERNSTEDT-ASTERHOLM, I., ATTIE, A.D., SHERWANI, S., WANG, Z.V., LANDSKRONER-EIGER, S., DINEEN, S., MAGALANG, U.J., BREKKEN, R.A. and SCHERER, P.E., 2009. Hypoxia-inducible factor 1alpha induces fibrosis and insulin resistance in white adipose tissue. *Molecular and cellular biology*, **29**(16), pp. 4467-4483.

HAN, J., LEE, J.E., JIN, J., LIM, J.S., OH, N., KIM, K., CHANG, S.I., SHIBUYA, M., KIM, H. and KOH, G.Y., 2011. The spatiotemporal development of adipose tissue. *Development (Cambridge, England)*, **138**(22), pp. 5027-5037.

HANNON, R., CROXTALL, J.D., GETTING, S.J., ROVIEZZO, F., YONA, S., PAUL-CLARK, M.J., GAVINS, F.N., PERRETTI, M., MORRIS, J.F., BUCKINGHAM, J.C. and FLOWER, R.J., 2003. Aberrant inflammation and resistance to glucocorticoids in annexin 1-/- mouse. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **17**(2), pp. 253-255.

- HAO, R.H., YANG, T.L., RONG, Y., YAO, S., DONG, S.S., CHEN, H. and GUO, Y., 2018. Gene expression profiles indicate tissue-specific obesity regulation changes and strong obesity relevant tissues. *International journal of obesity (2005)*, **42**(3), pp. 363-369.
- HAQUE, W.A., SHIMOMURA, I., MATSUZAWA, Y. and GARG, A., 2002. Serum adiponectin and leptin levels in patients with lipodystrophies. *The Journal of clinical endocrinology and metabolism*, **87**(5), pp. 2395.
- HARMON, D.B., SRIKAKULAPU, P., KAPLAN, J.L., OLDHAM, S.N., MCSKIMMING, C., GARMEY, J.C., PERRY, H.M., KIRBY, J.L., PROHASKA, T.A., GONEN, A., HALLOWELL, P., SCHIRMER, B., TSIMIKAS, S., TAYLOR, A.M., WITZTUM, J.L. and MCNAMARA, C.A., 2016. Protective Role for B-1b B Cells and IgM in Obesity-Associated Inflammation, Glucose Intolerance, and Insulin Resistance. *Arteriosclerosis, Thrombosis, and Vascular Biology*, **36**(4), pp. 682-691.
- HARMS, M. and SEALE, P., 2013. Brown and beige fat: development, function and therapeutic potential. *Nature medicine*, **19**(10), pp. 1252-1263.
- HARRIS, R.B., 2014. Direct and indirect effects of leptin on adipocyte metabolism. *Biochimica et biophysica acta*, **1842**(3), pp. 414-423.
- HASENFUSS, S.C., BAKIRI, L., THOMSEN, M.K., WILLIAMS, E.G., AUWERX, J. and WAGNER, E.F., 2014. Regulation of steatohepatitis and PPARgamma signaling by distinct AP-1 dimers. *Cell metabolism*, **19**(1), pp. 84-95.
- HASLAM, D.W. and JAMES, W.P., 2005. Obesity. *Lancet (London, England)*, **366**(9492), pp. 1197-1209.
- HATA, K., NISHIMURA, R., IKEDA, F., YAMASHITA, K., MATSUBARA, T., NOKUBI, T. and YONEDA, T., 2003. Differential roles of Smad1 and p38 kinase in regulation of peroxisome proliferator-activating receptor gamma during bone morphogenetic protein 2-induced adipogenesis. *Molecular biology of the cell*, **14**(2), pp. 545-555.
- HAYOZ, C., HERMANN, T., RAPTIS, D.A., BRONNIMANN, A., PETERLI, R. and ZUBER, M., 2018. Comparison of metabolic outcomes in patients undergoing laparoscopic roux-en-Y gastric bypass versus sleeve gastrectomy - a systematic review and meta-analysis of randomised controlled trials. *Swiss medical weekly*, **148**, pp. w14633.
- HAZELDINE, J., HAMPSON, P., OPOKU, F.A., FOSTER, M. and LORD, J.M., 2015. N-Formyl peptides drive mitochondrial damage associated molecular pattern induced neutrophil activation through ERK1/2 and P38 MAP kinase signalling pathways. *Injury*, **46**(6), pp. 975-984.

- HE, H.Q. and YE, R.D., 2017. The Formyl Peptide Receptors: Diversity of Ligands and Mechanism for Recognition. *Molecules (Basel, Switzerland)*, **22**(3), pp. 10.3390/molecules22030455.
- HE, W., BARAK, Y., HEVENER, A., OLSON, P., LIAO, D., LE, J., NELSON, M., ONG, E., OLEFSKY, J.M. and EVANS, R.M., 2003. Adipose-specific peroxisome proliferator-activated receptor gamma knockout causes insulin resistance in fat and liver but not in muscle. *Proceedings of the National Academy of Sciences of the United States of America*, **100**(26), pp. 15712-15717.
- HE, Z., ZHU, H.H., BAULER, T.J., WANG, J., CIARALDI, T., ALDERSON, N., LI, S., RAQUIL, M.A., JI, K., WANG, S., SHAO, J., HENRY, R.R., KING, P.D. and FENG, G.S., 2013. Nonreceptor tyrosine phosphatase Shp2 promotes adipogenesis through inhibition of p38 MAP kinase. *Proceedings of the National Academy of Sciences of the United States of America*, **110**(1), pp. E79-88.
- HECTOR, J., SCHWARZLOH, B., GOEHRING, J., STRATE, T.G., HESS, U.F., DEURETZBACHER, G., HANSEN-ALGENSTAEDT, N., BEIL, F.U. and ALGENSTAEDT, P., 2007. TNF-alpha alters visfatin and adiponectin levels in human fat. *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et metabolisme*, **39**(4), pp. 250-255.
- HEFFRON, S.P., PARIKH, A., VOLODARSKIY, A., REN-FIELDING, C., SCHWARTZBARD, A., NICHOLSON, J. and BANGALORE, S., 2016. Changes in Lipid Profile of Obese Patients Following Contemporary Bariatric Surgery: A Meta-Analysis. *The American Journal of Medicine*, **129**(9), pp. 952-959.
- HEGELE, R.A., CAO, H., HUFF, M.W. and ANDERSON, C.M., 2000. LMNA R482Q mutation in partial lipodystrophy associated with reduced plasma leptin concentration. *The Journal of clinical endocrinology and metabolism*, **85**(9), pp. 3089-3093.
- HEILBRONN, L., SMITH, S.R. and RAVUSSIN, E., 2004. Failure of fat cell proliferation, mitochondrial function and fat oxidation results in ectopic fat storage, insulin resistance and type II diabetes mellitus. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*, **28 Suppl 4**, pp. S12-21.
- HELLERSTEIN, M.K., 1999. De novo lipogenesis in humans: metabolic and regulatory aspects. *European journal of clinical nutrition*, **53 Suppl 1**, pp. S53-65.
- HENEGAR, C., TORDJMAN, J., ACHARD, V., LACASA, D., CREMER, I., GUERREMILLO, M., POITOU, C., BASDEVANT, A., STICH, V., VIGUERIE, N., LANGIN, D., BEDOSSA, P., ZUCKER, J.D. and CLEMENT, K., 2008. Adipose tissue transcriptomic signature highlights the pathological relevance of extracellular matrix in human obesity. *Genome biology*, **9**(1), pp. R14-2008-9-1-r14.

- HEPLER, C., VISHVANATH, L. and GUPTA, R.K., 2017. Sorting out adipocyte precursors and their role in physiology and disease. *Genes & development*, **31**(2), pp. 127-140.
- HERRANZ, P., DE LUCAS, R., PEREZ-ESPANA, L. and MAYOR, M., 2008. Lipodystrophy syndromes. *Dermatologic clinics*, **26**(4), pp. 569-78, ix.
- HERRERO, L., SHAPIRO, H., NAYER, A., LEE, J. and SHOELSON, S.E., 2010. Inflammation and adipose tissue macrophages in lipodystrophic mice. *Proceedings of the National Academy of Sciences of the United States of America*, **107**(1), pp. 240-245.
- HIGUCHI, M., DUSTING, G.J., PESHAVARIYA, H., JIANG, F., HSIAO, S.T., CHAN, E.C. and LIU, G.S., 2013. Differentiation of human adipose-derived stem cells into fat involves reactive oxygen species and Forkhead box O1 mediated upregulation of antioxidant enzymes. *Stem cells and development*, **22**(6), pp. 878-888.
- HIRSCH, J., FRIED, S.K., EDENS, N.K. and LEIBEL, R.L., 1989. The fat cell. *The Medical clinics of North America*, **73**(1), pp. 83-96.
- HO, I.C., KIM, J.H., ROONEY, J.W., SPIEGELMAN, B.M. and GLIMCHER, L.H., 1998. A potential role for the nuclear factor of activated T cells family of transcriptional regulatory proteins in adipogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, **95**(26), pp. 15537-15541.
- HOLST, J.J., 2007. The physiology of glucagon-like peptide 1. *Physiological Reviews*, **87**(4), pp. 1409-1439.
- HOPKINS, M. and BLUNDELL, J.E., 2016. Energy balance, body composition, sedentariness and appetite regulation: pathways to obesity. *Clinical science (London, England : 1979)*, **130**(18), pp. 1615-1628.
- HOPKINS, M., GIBBONS, C., CAUDWELL, P., HELLSTROM, P.M., NASLUND, E., KING, N.A., FINLAYSON, G. and BLUNDELL, J.E., 2014. The adaptive metabolic response to exercise-induced weight loss influences both energy expenditure and energy intake. *European journal of clinical nutrition*, **68**(5), pp. 581-586.
- HOPKINS, M., KING, N.A. and BLUNDELL, J.E., 2010. Acute and long-term effects of exercise on appetite control: is there any benefit for weight control? *Current opinion in clinical nutrition and metabolic care*, **13**(6), pp. 635-640.
- HORTON, J.D., SHIMOMURA, I., BROWN, M.S., HAMMER, R.E., GOLDSTEIN, J.L. and SHIMANO, H., 1998. Activation of cholesterol synthesis in preference to fatty acid synthesis in liver and adipose tissue of transgenic mice overproducing sterol regulatory element-binding protein-2. *The Journal of clinical investigation*, **101**(11), pp. 2331-2339.
- HOSOGAI, N., FUKUHARA, A., OSHIMA, K., MIYATA, Y., TANAKA, S., SEGAWA, K., FURUKAWA, S., TOCHINO, Y., KOMURO, R., MATSUDA, M. and

- SHIMOMURA, I., 2007. Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes*, **56**(4), pp. 901-911.
- HOTAMISLIGIL, G.S., SHARGILL, N.S. and SPIEGELMAN, B.M., 1993. Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science (New York, N.Y.)*, **259**(5091), pp. 87-91.
- HOTAMISLIGIL, G.S. and SPIEGELMAN, B.M., 1994. Tumor necrosis factor α : a key component of the obesity-diabetes link. *Diabetes*, **43**(11), pp. 1271-1278.
- HOTTA, K., FUNAHASHI, T., BODKIN, N.L., ORTMAYER, H.K., ARITA, Y., HANSEN, B.C. and MATSUZAWA, Y., 2001. Circulating concentrations of the adipocyte protein adiponectin are decreased in parallel with reduced insulin sensitivity during the progression to type 2 diabetes in rhesus monkeys. *Diabetes*, **50**(5), pp. 1126-1133.
- HOUSTIS, N., ROSEN, E.D. and LANDER, E.S., 2006. Reactive oxygen species have a causal role in multiple forms of insulin resistance. *Nature*, **440**(7086), pp. 944-948.
- HRUBY, A. and HU, F.B., 2015. The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*, **33**(7), pp. 673-689.
- HU, E., KIM, J.B., SARRAF, P. and SPIEGELMAN, B.M., 1996. Inhibition of adipogenesis through MAP kinase-mediated phosphorylation of PPAR γ . *Science (New York, N.Y.)*, **274**(5295), pp. 2100-2103.
- HU, E., LIANG, P. and SPIEGELMAN, B.M., 1996. AdipoQ is a novel adipose-specific gene dysregulated in obesity. *The Journal of biological chemistry*, **271**(18), pp. 10697-10703.
- HU, F.B., 2007. Obesity and mortality: watch your waist, not just your weight. *Archives of Internal Medicine*, **167**(9), pp. 875-876.
- HUANG, J., JIA, Y., FU, T., VISWAKARMA, N., BAI, L., RAO, M.S., ZHU, Y., BORENSZTAJN, J. and REDDY, J.K., 2012. Sustained activation of PPAR α by endogenous ligands increases hepatic fatty acid oxidation and prevents obesity in ob/ob mice. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **26**(2), pp. 628-638.
- HUANG, Z.H., MANICKAM, B., RYVKIN, V., ZHOU, X.J., FANTUZZI, G., MAZZONE, T. and SAM, S., 2013. PCOS is associated with increased CD11c expression and crown-like structures in adipose tissue and increased central abdominal fat depots independent of obesity. *The Journal of clinical endocrinology and metabolism*, **98**(1), pp. E17-24.
- HUSZAR, D., LYNCH, C.A., FAIRCHILD-HUNTRESS, V., DUNMORE, J.H., FANG, Q., BERKEMEIER, L.R., GU, W., KESTERSON, R.A., BOSTON, B.A., CONE, R.D.,

- SMITH, F.J., CAMPFIELD, L.A., BURN, P. and LEE, F., 1997. Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell*, **88**(1), pp. 131-141.
- IKEMOTO, S., THOMPSON, K.S., TAKAHASHI, M., ITAKURA, H., LANE, M.D. and EZAKI, O., 1995. High fat diet-induced hyperglycemia: prevention by low level expression of a glucose transporter (GLUT4) minigene in transgenic mice. *Proceedings of the National Academy of Sciences of the United States of America*, **92**(8), pp. 3096-3099.
- ISHIDA, H., MITAMURA, T., TAKAHASHI, Y., HISATOMI, A., FUKUHARA, Y., MURATO, K. and OHARA, K., 1997. Cataract development induced by repeated oral dosing with FK506 (tacrolimus) in adult rats. *Toxicology*, **123**(3), pp. 167-175.
- IWAKI-EGAWA, S., WATANABE, Y., KIKUYA, Y. and FUJIMOTO, Y., 1998. Dipeptidyl peptidase IV from human serum: purification, characterization, and N-terminal amino acid sequence. *Journal of Biochemistry*, **124**(2), pp. 428-433.
- IYER, A., FAIRLIE, D.P., PRINS, J.B., HAMMOCK, B.D. and BROWN, L., 2010. Inflammatory lipid mediators in adipocyte function and obesity. *Nature reviews.Endocrinology*, **6**(2), pp. 71-82.
- JACKSON, R.M., GRIESEL, B.A., GURLEY, J.M., SZWEDA, L.I. and OLSON, A.L., 2017. Glucose availability controls adipogenesis in mouse 3T3-L1 adipocytes via up-regulation of nicotinamide metabolism. *The Journal of biological chemistry*, **292**(45), pp. 18556-18564.
- JACQUES, C., HOLZENBERGER, M., MLADENOVIC, Z., SALVAT, C., PECCHI, E., BERENBAUM, F. and GOSSET, M., 2012. Proinflammatory actions of visfatin/nicotinamide phosphoribosyltransferase (Nampt) involve regulation of insulin signaling pathway and Nampt enzymatic activity. *The Journal of biological chemistry*, **287**(18), pp. 15100-15108.
- JAKICIC, J.M., MARCUS, B.H., LANG, W. and JANNEY, C., 2008. Effect of exercise on 24-month weight loss maintenance in overweight women. *Archives of Internal Medicine*, **168**(14), pp. 1550-9; discussion 1559-60.
- JAN, V., CERVERA, P., MAACHI, M., BAUDRIMONT, M., KIM, M., VIDAL, H., GIRARD, P.M., LEVAN, P., ROZENBAUM, W., LOMBES, A., CAPEAU, J. and BASTARD, J.P., 2004. Altered fat differentiation and adipocytokine expression are inter-related and linked to morphological changes and insulin resistance in HIV-1-infected lipodystrophic patients. *Antiviral Therapy*, **9**(4), pp. 555-564.
- JAUNOO, S.S. and SOUTHALL, P.J., 2010. Bariatric surgery. *International journal of surgery (London, England)*, **8**(2), pp. 86-89.
- JOHN, K., MARINO, J.S., SANCHEZ, E.R. and HINDS, T.D., Jr, 2016. The glucocorticoid receptor: cause of or cure for obesity? *American journal of physiology.Endocrinology and metabolism*, **310**(4), pp. E249-57.

- JOHNSTONE, S.A., HUBAISHY, I. and WAISMAN, D.M., 1993. Regulation of annexin I-dependent aggregation of phospholipid vesicles by protein kinase C. *The Biochemical journal*, **294** (Pt 3)(Pt 3), pp. 801-807.
- JOOSEN, A.M. and WESTERTERP, K.R., 2006. Energy expenditure during overfeeding. *Nutrition & metabolism*, **3**, pp. 25-7075-3-25.
- JUNG, S.H., PARK, H.S., KIM, K.S., CHOI, W.H., AHN, C.W., KIM, B.T., KIM, S.M., LEE, S.Y., AHN, S.M., KIM, Y.K., KIM, H.J., KIM, D.J. and LEE, K.W., 2008. Effect of weight loss on some serum cytokines in human obesity: increase in IL-10 after weight loss. *The Journal of nutritional biochemistry*, **19**(6), pp. 371-375.
- KAMAL, A.H., KIM, W.K., CHO, K., PARK, A., MIN, J.K., HAN, B.S., PARK, S.G., LEE, S.C. and BAE, K.H., 2013. Investigation of adipocyte proteome during the differentiation of brown preadipocytes. *Journal of proteomics*, **94**, pp. 327-336.
- KAMINSKA, A., KOPCZYNSKA, E., BRONISZ, A., ZMUDZINSKA, M., BIELINSKI, M., BORKOWSKA, A., TYRAKOWSKI, T. and JUNIK, R., 2010. An evaluation of visfatin levels in obese subjects. *Endokrynologia Polska*, **61**(2), pp. 169-173.
- KANDA, H., TATEYA, S., TAMORI, Y., KOTANI, K., HIASA, K., KITAZAWA, R., KITAZAWA, S., MIYACHI, H., MAEDA, S., EGASHIRA, K. and KASUGA, M., 2006. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *The Journal of clinical investigation*, **116**(6), pp. 1494-1505.
- KARELIS, A.D., ST-PIERRE, D.H., CONUS, F., RABASA-LHORET, R. and POEHLMAN, E.T., 2004. Metabolic and body composition factors in subgroups of obesity: what do we know? *The Journal of clinical endocrinology and metabolism*, **89**(6), pp. 2569-2575.
- KARKI, S., FARB, M.G., MYERS, S., APOVIAN, C., HESS, D.T. and GOKCE, N., 2015. Effect of Bariatric Weight Loss on the Adipose Lipolytic Transcriptome in Obese Humans. *Mediators of inflammation*, **2015**, pp. 106237.
- KARPE, F., DICKMANN, J.R. and FRAYN, K.N., 2011. Fatty acids, obesity, and insulin resistance: time for a reevaluation. *Diabetes*, **60**(10), pp. 2441-2449.
- KASER, S., KASER, A., SANDHOFER, A., EBENBICHLER, C.F., TILG, H. and PATSCH, J.R., 2003. Resistin messenger-RNA expression is increased by proinflammatory cytokines in vitro. *Biochemical and biophysical research communications*, **309**(2), pp. 286-290.
- KAWAI, M., NAMBA, N., MUSHIAKE, S., ETANI, Y., NISHIMURA, R., MAKISHIMA, M. and OZONO, K., 2007. Growth hormone stimulates adipogenesis of 3T3-L1 cells through activation of the Stat5A/5B-PPARgamma pathway. *Journal of Molecular Endocrinology*, **38**(1-2), pp. 19-34.

- KAWAMURA, Y., TANAKA, Y., KAWAMORI, R. and MAEDA, S., 2006. Overexpression of Kruppel-like factor 7 regulates adipocytokine gene expressions in human adipocytes and inhibits glucose-induced insulin secretion in pancreatic beta-cell line. *Molecular endocrinology (Baltimore, Md.)*, **20**(4), pp. 844-856.
- KEIRE, D.A., BOWERS, C.W., SOLOMON, T.E. and REEVE, J.R., Jr, 2002. Structure and receptor binding of PYY analogs. *Peptides*, **23**(2), pp. 305-321.
- KENNEDY, D.J., KUCHIBHOTLA, S., WESTFALL, K.M., SILVERSTEIN, R.L., MORTON, R.E. and FEBBRAIO, M., 2011. A CD36-dependent pathway enhances macrophage and adipose tissue inflammation and impairs insulin signalling. *Cardiovascular research*, **89**(3), pp. 604-613.
- KERN, P.A., RANGANATHAN, S., LI, C., WOOD, L. and RANGANATHAN, G., 2001. Adipose tissue tumor necrosis factor and interleukin-6 expression in human obesity and insulin resistance. *American journal of physiology. Endocrinology and metabolism*, **280**(5), pp. E745-51.
- KERN, P.A., SAGHIZADEH, M., ONG, J.M., BOSCH, R.J., DEEM, R. and SIMSOLO, R.B., 1995. The expression of tumor necrosis factor in human adipose tissue. Regulation by obesity, weight loss, and relationship to lipoprotein lipase. *The Journal of clinical investigation*, **95**(5), pp. 2111-2119.
- KERSTEN, S., 2001. Mechanisms of nutritional and hormonal regulation of lipogenesis. *EMBO reports*, **2**(4), pp. 282-286.
- KHAN, A.A., ALSAHLI, M.A. and RAHMANI, A.H., 2018. Myeloperoxidase as an Active Disease Biomarker: Recent Biochemical and Pathological Perspectives. *Medical sciences (Basel, Switzerland)*, **6**(2), pp. 10.3390/medsci6020033.
- KIM, J.Y., VAN DE WALL, E., LAPLANTE, M., AZZARA, A., TRUJILLO, M.E., HOFMANN, S.M., SCHRAW, T., DURAND, J.L., LI, H., LI, G., JELICKS, L.A., MEHLER, M.F., HUI, D.Y., DESHAIES, Y., SHULMAN, G.I., SCHWARTZ, G.J. and SCHERER, P.E., 2007. Obesity-associated improvements in metabolic profile through expansion of adipose tissue. *The Journal of clinical investigation*, **117**(9), pp. 2621-2637.
- KLEE, C.B., CROUCH, T.H. and KRINKS, M.H., 1979. Calcineurin: a calcium- and calmodulin-binding protein of the nervous system. *Proceedings of the National Academy of Sciences of the United States of America*, **76**(12), pp. 6270-6273.
- KOLDITZ, C.I. and LANGIN, D., 2010. Adipose tissue lipolysis. *Current opinion in clinical nutrition and metabolic care*, **13**(4), pp. 377-381.
- KOLEHMAINEN, M., VIDAL, H., ALHAVA, E. and UUSITUPA, M.I., 2001. Sterol regulatory element binding protein 1c (SREBP-1c) expression in human obesity. *Obesity research*, **9**(11), pp. 706-712.

- KOSICKA, A., CUNLIFFE, A.D., MACKENZIE, R., ZARIWALA, M.G., PERRETTI, M., FLOWER, R.J. and RENSHAW, D., 2013. Attenuation of plasma annexin A1 in human obesity. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **27**(1), pp. 368-378.
- KOUIDHI, S., BERRHOUMA, R., ROUISSI, K., JARBOUI, S., CLERGET-FROIDEVAUX, M.S., SEUGNET, I., BCHIR, F., DEMENEIX, B., GUISSOUMA, H. and ELGAAIED, A.B., 2013. Human subcutaneous adipose tissue Glut 4 mRNA expression in obesity and type 2 diabetes. *Acta Diabetologica*, **50**(2), pp. 227-232.
- KOVACS, P., HANSON, R.L., LEE, Y.H., YANG, X., KOBES, S., PERMANA, P.A., BOGARDUS, C. and BAIER, L.J., 2003. The role of insulin receptor substrate-1 gene (IRS1) in type 2 diabetes in Pima Indians. *Diabetes*, **52**(12), pp. 3005-3009.
- KOZAK, L.P., 2010. Brown fat and the myth of diet-induced thermogenesis. *Cell metabolism*, **11**(4), pp. 263-267.
- KRAAKMAN, M.J., MURPHY, A.J., JANDELEIT-DAHME, K. and KAMMOUN, H.L., 2014. Macrophage polarization in obesity and type 2 diabetes: weighing down our understanding of macrophage function? *Frontiers in immunology*, **5**, pp. 470.
- KRAMER, D.K., AL-KHALILI, L., GUIGAS, B., LENG, Y., GARCIA-ROVES, P.M. and KROOK, A., 2007. Role of AMP kinase and PPARdelta in the regulation of lipid and glucose metabolism in human skeletal muscle. *The Journal of biological chemistry*, **282**(27), pp. 19313-19320.
- KUBOTA, N., TERAUCHI, Y., YAMAUCHI, T., KUBOTA, T., MOROI, M., MATSUI, J., ETO, K., YAMASHITA, T., KAMON, J., SATOH, H., YANO, W., FROGUEL, P., NAGAI, R., KIMURA, S., KADOWAKI, T. and NODA, T., 2002. Disruption of adiponectin causes insulin resistance and neointimal formation. *The Journal of biological chemistry*, **277**(29), pp. 25863-25866.
- KUMADA, M., KIHARA, S., OUCHI, N., KOBAYASHI, H., OKAMOTO, Y., OHASHI, K., MAEDA, K., NAGARETANI, H., KISHIDA, K., MAEDA, N., NAGASAWA, A., FUNAHASHI, T. and MATSUZAWA, Y., 2004. Adiponectin specifically increased tissue inhibitor of metalloproteinase-1 through interleukin-10 expression in human macrophages. *Circulation*, **109**(17), pp. 2046-2049.
- KUPPER, T.S. and GROVES, R.W., 1995. The interleukin-1 axis and cutaneous inflammation. *The Journal of investigative dermatology*, **105**(1 Suppl), pp. 62S-66S.
- KWOK, K.H., LAM, K.S. and XU, A., 2016. Heterogeneity of white adipose tissue: molecular basis and clinical implications. *Experimental & molecular medicine*, **48**, pp. e215.
- KWON, H. and PESSIN, J.E., 2013. Adipokines mediate inflammation and insulin resistance. *Frontiers in endocrinology*, **4**, pp. 71.

- KWON, J.H., LEE, J.H., KIM, K.S., CHUNG, Y.W. and KIM, I.Y., 2012. Regulation of cytosolic phospholipase A2 phosphorylation by proteolytic cleavage of annexin A1 in activated mast cells. *Journal of immunology (Baltimore, Md.: 1950)*, **188**(11), pp. 5665-5673.
- LAFONTAN, M., 2014. Adipose tissue and adipocyte dysregulation. *Diabetes & metabolism*, **40**(1), pp. 16-28.
- LAGRAND, W.K., VISSER, C.A., HERMENS, W.T., NIESSEN, H.W., VERHEUGT, F.W., WOLBINK, G.J. and HACK, C.E., 1999. C-reactive protein as a cardiovascular risk factor: more than an epiphenomenon? *Circulation*, **100**(1), pp. 96-102.
- LAMERS, D., FAMULLA, S., WRONKOWITZ, N., HARTWIG, S., LEHR, S., OUWENS, D.M., ECKARDT, K., KAUFMAN, J.M., RYDEN, M., MÄLLER, S., HANISCH, F.G., RUIGE, J., ARNER, P., SELL, H. and ECKEL, J., 2011. Dipeptidyl peptidase 4 is a novel adipokine potentially linking obesity to the metabolic syndrome. *Diabetes*, **60**(7), pp. 1917-1925.
- LAURENCIKIENE, J. and RYDEN, M., 2012. Liver X receptors and fat cell metabolism. *International journal of obesity (2005)*, **36**(12), pp. 1494-1502.
- LAWRENCE, T. and GILROY, D.W., 2007. Chronic inflammation: a failure of resolution? *International journal of experimental pathology*, **88**(2), pp. 85-94.
- LE, Y., MURPHY, P.M. and WANG, J.M., 2002. Formyl-peptide receptors revisited. *Trends in immunology*, **23**(11), pp. 541-548.
- LEBIEN, T.W. and TEDDER, T.F., 2008. B lymphocytes: how they develop and function. *Blood*, **112**(5), pp. 1570-1580.
- LEE, B.C. and LEE, J., 2014. Cellular and molecular players in adipose tissue inflammation in the development of obesity-induced insulin resistance. *Biochimica et biophysica acta*, **1842**(3), pp. 446-462.
- LEE, H., LEE, Y.J., CHOI, H., KO, E.H. and KIM, J.W., 2009. Reactive oxygen species facilitate adipocyte differentiation by accelerating mitotic clonal expansion. *The Journal of biological chemistry*, **284**(16), pp. 10601-10609.
- LEE, J., 2013. Adipose tissue macrophages in the development of obesity-induced inflammation, insulin resistance and type 2 diabetes. *Archives of Pharmacal Research*, **36**(2), pp. 208-222.
- LEE, K., VILLENA, J.A., MOON, Y.S., KIM, K.H., LEE, S., KANG, C. and SUL, H.S., 2003. Inhibition of adipogenesis and development of glucose intolerance by soluble preadipocyte factor-1 (Pref-1). *The Journal of clinical investigation*, **111**(4), pp. 453-461.
- LEE, Y.H., MOTTILLO, E.P. and GRANNEMAN, J.G., 2014. Adipose tissue plasticity from WAT to BAT and in between. *Biochimica et biophysica acta*, **1842**(3), pp. 358-369.

- LEHNER, R. and KUKSIS, A., 1996. Biosynthesis of triacylglycerols. *Progress in lipid research*, **35**(2), pp. 169-201.
- LEIBEL, R.L., ROSENBAUM, M. and HIRSCH, J., 1995. Changes in energy expenditure resulting from altered body weight. *The New England journal of medicine*, **332**(10), pp. 621-628.
- LEIBOWITZ, S.F., HAMMER, N.J. and CHANG, K., 1981. Hypothalamic paraventricular nucleus lesions produce overeating and obesity in the rat. *Physiology & Behavior*, **27**(6), pp. 1031-1040.
- LEPPER, C. and FAN, C.M., 2010. Inducible lineage tracing of Pax7-descendant cells reveals embryonic origin of adult satellite cells. *Genesis (New York, N.Y.: 2000)*, **48**(7), pp. 424-436.
- LEVI, J., GRAY, S.L., SPECK, M., HUYNH, F.K., BABICH, S.L., GIBSON, W.T. and KIEFFER, T.J., 2011. Acute disruption of leptin signaling in vivo leads to increased insulin levels and insulin resistance. *Endocrinology*, **152**(9), pp. 3385-3395.
- LEY, R.E., TURNBAUGH, P.J., KLEIN, S. and GORDON, J.I., 2006. Microbial ecology: human gut microbes associated with obesity. *Nature*, **444**(7122), pp. 1022-1023.
- LI, D., YEA, S., LI, S., CHEN, Z., NARLA, G., BANCK, M., LABORDA, J., TAN, S., FRIEDMAN, J.M., FRIEDMAN, S.L. and WALSH, M.J., 2005. Kruppel-like factor-6 promotes preadipocyte differentiation through histone deacetylase 3-dependent repression of DLK1. *The Journal of biological chemistry*, **280**(29), pp. 26941-26952.
- LIAO, W., NGUYEN, M.T., YOSHIZAKI, T., FAVELYUKIS, S., PATSOURIS, D., IMAMURA, T., VERMA, I.M. and OLEFSKY, J.M., 2007. Suppression of PPAR-gamma attenuates insulin-stimulated glucose uptake by affecting both GLUT1 and GLUT4 in 3T3-L1 adipocytes. *American journal of physiology. Endocrinology and metabolism*, **293**(1), pp. E219-27.
- LINDEGAARD, K.K., JORGENSEN, N.B., JUST, R., HEEGAARD, P.M. and MADSBAD, S., 2015. Effects of Roux-en-Y gastric bypass on fasting and postprandial inflammation-related parameters in obese subjects with normal glucose tolerance and in obese subjects with type 2 diabetes. *Diabetology & metabolic syndrome*, **7**, pp. 12-015-0012-9. eCollection 2015.
- LINDGREN, C.M., NILSSON, A., ORHO-MELANDER, M., ALMGREN, P. and GROOP, L.C., 2001. Characterization of the annexin I gene and evaluation of its role in type 2 diabetes. *Diabetes*, **50**(10), pp. 2402-2405.
- LIU, C., HUANG, K., LI, G., WANG, P., LIU, C., GUO, C., SUN, Z. and PAN, J., 2017. Ascorbic acid promotes 3T3-L1 cells adipogenesis by attenuating ERK signaling to upregulate the collagen VI. *Nutrition & metabolism*, **14**, pp. 79-017-0234-y. eCollection 2017.

- LIU, J., FOX, C.S., HICKSON, D.A., MAY, W.D., HAIRSTON, K.G., CARR, J.J. and TAYLOR, H.A., 2010. Impact of abdominal visceral and subcutaneous adipose tissue on cardiometabolic risk factors: the Jackson Heart Study. *The Journal of clinical endocrinology and metabolism*, **95**(12), pp. 5419-5426.
- LIU, S., XU, R., GERIN, I., CAWTHORN, W.P., MACDOUGALD, O.A., CHEN, X.W., SALTIEL, A.R., KOENIG, R.J. and XU, B., 2014. SRA regulates adipogenesis by modulating p38/JNK phosphorylation and stimulating insulin receptor gene expression and downstream signaling. *PloS one*, **9**(4), pp. e95416.
- LIVAK, K.J. and SCHMITTGEN, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)*, **25**(4), pp. 402-408.
- LIZARBE, M.A., BARRASA, J.I., OLMO, N., GAVILANES, F. and TURNAY, J., 2013. Annexin-phospholipid interactions. Functional implications. *International journal of molecular sciences*, **14**(2), pp. 2652-2683.
- LO, J.C., LJUBICIC, S., LEIBIGER, B., KERN, M., LEIBIGER, I.B., MOEDE, T., KELLY, M.E., CHATTERJEE BHOWMICK, D., MURANO, I., COHEN, P., BANKS, A.S., KHANDEKAR, M.J., DIETRICH, A., FLIER, J.S., CINTI, S., BLUHER, M., DANIAL, N.N., BERGGREN, P.O. and SPIEGELMAN, B.M., 2014. Adipsin is an adipokine that improves beta cell function in diabetes. *Cell*, **158**(1), pp. 41-53.
- LOEBIG, M., KLEMENT, J., SCHMOLLER, A., BETZ, S., HEUCK, N., SCHWEIGER, U., PETERS, A., SCHULTES, B. and OLTMANN, K.M., 2010. Evidence for a relationship between VEGF and BMI independent of insulin sensitivity by glucose clamp procedure in a homogenous group healthy young men. *PloS one*, **5**(9), pp. e12610.
- LOFFREDA, S., YANG, S.Q., LIN, H.Z., KARP, C.L., BRENGMAN, M.L., WANG, D.J., KLEIN, A.S., BULKLEY, G.B., BAO, C., NOBLE, P.W., LANE, M.D. and DIEHL, A.M., 1998. Leptin regulates proinflammatory immune responses. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **12**(1), pp. 57-65.
- LOFTUS, T.M., KUHAJDA, F.P. and LANE, M.D., 1998. Insulin depletion leads to adipose-specific cell death in obese but not lean mice. *Proceedings of the National Academy of Sciences of the United States of America*, **95**(24), pp. 14168-14172.
- LOMONACO, R., ORTIZ-LOPEZ, C., ORSAK, B., WEBB, A., HARDIES, J., DARLAND, C., FINCH, J., GASTALDELLI, A., HARRISON, S., TIO, F. and CUSI, K., 2012. Effect of adipose tissue insulin resistance on metabolic parameters and liver histology in obese patients with nonalcoholic fatty liver disease. *Hepatology (Baltimore, Md.)*, **55**(5), pp. 1389-1397.
- LONGO, K.A., WRIGHT, W.S., KANG, S., GERIN, I., CHIANG, S.H., LUCAS, P.C., OPP, M.R. and MACDOUGALD, O.A., 2004. Wnt10b inhibits development of white

and brown adipose tissues. *The Journal of biological chemistry*, **279**(34), pp. 35503-35509.

LOOS, R.J., LINDGREN, C.M., LI, S., WHEELER, E., ZHAO, J.H., PROKOPENKO, I., INOUE, M., FREATHY, R.M., ATTWOOD, A.P., BECKMANN, J.S., BERNDT, S.I., PROSTATE, LUNG, COLORECTAL, AND OVARIAN (PLCO) CANCER SCREENING TRIAL, JACOBS, K.B., CHANOCK, S.J., HAYES, R.B., BERGMANN, S., BENNETT, A.J., BINGHAM, S.A., BOCHUD, M., BROWN, M., CAUCHI, S., CONNELL, J.M., COOPER, C., SMITH, G.D., DAY, I., DINA, C., DE, S., DERMITZAKIS, E.T., DONEY, A.S., ELLIOTT, K.S., ELLIOTT, P., EVANS, D.M., SADAF FAROOQI, I., FROGUEL, P., GHORI, J., GROVES, C.J., GWILLIAM, R., HADLEY, D., HALL, A.S., HATTERSLEY, A.T., HEBEBRAND, J., HEID, I.M., KORA, LAMINA, C., GIEGER, C., ILLIG, T., MEITINGER, T., WICHMANN, H.E., HERRERA, B., HINNEY, A., HUNT, S.E., JARVELIN, M.R., JOHNSON, T., JOLLEY, J.D., KARPE, F., KENIRY, A., KHAW, K.T., LUBEN, R.N., MANGINO, M., MARCHINI, J., MCARDLE, W.L., MCGINNIS, R., MEYRE, D., MUNROE, P.B., MORRIS, A.D., NESS, A.R., NEVILLE, M.J., NICA, A.C., ONG, K.K., O'RAHILLY, S., OWEN, K.R., PALMER, C.N., PAPADAKIS, K., POTTER, S., POUTA, A., QI, L., NURSES' HEALTH STUDY, RANDALL, J.C., RAYNER, N.W., RING, S.M., SANDHU, M.S., SCHERAG, A., SIMS, M.A., SONG, K., SORANZO, N., SPELIOTES, E.K., DIABETES GENETICS INITIATIVE, SYDDALL, H.E., TEICHMANN, S.A., TIMPSON, N.J., TOBIAS, J.H., UDA, M., SARDINIA STUDY, VOGEL, C.I., WALLACE, C., WATERWORTH, D.M., WEEDON, M.N., WELLCOME TRUST CASE CONTROL CONSORTIUM, WILLER, C.J., FUSION, WRAIGHT, YUAN, X., ZEGGINI, E., HIRSCHHORN, J.N., STRACHAN, D.P., OUWEHAND, W.H., CAULFIELD, M.J., SAMANI, N.J., FRAYLING, T.M., VOLLENWEIDER, P., WAEBER, G., MOOSER, V., DELOUKAS, P., MCCARTHY, M.I., WAREHAM, N.J., BARROSO, I., JACOBS, K.B., CHANOCK, S.J., HAYES, R.B., LAMINA, C., GIEGER, C., ILLIG, T., MEITINGER, T., WICHMANN, H.E., KRAFT, P., HANKINSON, S.E., HUNTER, D.J., HU, F.B., LYON, H.N., VOIGHT, B.F., RIDDERSTRALE, M., GROOP, L., SCHEET, P., SANNA, S., ABECASIS, G.R., ALBAI, G., NAGARAJA, R., SCHLESSINGER, D., JACKSON, A.U., TUOMILEHTO, J., COLLINS, F.S., BOEHNKE, M. and MOHLKE, K.L., 2008. Common variants near MC4R are associated with fat mass, weight and risk of obesity. *Nature genetics*, **40**(6), pp. 768-775.

LOPEZ, M., VARELA, L., VAZQUEZ, M.J., RODRIGUEZ-CUENCA, S., GONZALEZ, C.R., VELAGAPUDI, V.R., MORGAN, D.A., SCHOENMAKERS, E., AGASSANDIAN, K., LAGE, R., MARTINEZ DE MORENTIN, P.B., TOVAR, S., NOGUEIRAS, R., CARLING, D., LELLIOTT, C., GALLEGU, R., ORESIC, M., CHATTERJEE, K., SAHA, A.K., RAHMOUNI, K., DIEGUEZ, C. and VIDAL-PUIG, A., 2010. Hypothalamic AMPK and fatty acid metabolism mediate thyroid regulation of energy balance. *Nature medicine*, **16**(9), pp. 1001-1008.

LUCIANO, R., BARRACO, G.M., MURACA, M., OTTINO, S., SPREGHINI, M.R., SFORZA, R.W., RUSTICO, C., MORINO, G.S. and MANCO, M., 2015. Biomarkers of Alzheimer disease, insulin resistance, and obesity in childhood. *Pediatrics*, **135**(6), pp. 1074-1081.

- LUMENG, C.N., DEYOUNG, S.M. and SALTIEL, A.R., 2007. Macrophages block insulin action in adipocytes by altering expression of signaling and glucose transport proteins. *American journal of physiology. Endocrinology and metabolism*, **292**(1), pp. E166-74.
- MA, X., LEE, P., CHISHOLM, D.J. and JAMES, D.E., 2015. Control of adipocyte differentiation in different fat depots; implications for pathophysiology or therapy. *Frontiers in endocrinology*, **6**, pp. 1.
- MACDOUGALD, O.A., HWANG, C.S., FAN, H. and LANE, M.D., 1995. Regulated expression of the obese gene product (leptin) in white adipose tissue and 3T3-L1 adipocytes. *Proceedings of the National Academy of Sciences of the United States of America*, **92**(20), pp. 9034-9037.
- MACIA, L., DELACRE, M., ABBOUD, G., OUK, T.S., DELANOYE, A., VERWAERDE, C., SAULE, P. and WOLOWCZUK, I., 2006. Impairment of dendritic cell functionality and steady-state number in obese mice. *Journal of immunology (Baltimore, Md.: 1950)*, **177**(9), pp. 5997-6006.
- MACIEJEWSKI, M.L., ARTERBURN, D.E., VAN SCOYOC, L., SMITH, V.A., YANCY, W.S., Jr, WEIDENBACHER, H.J., LIVINGSTON, E.H. and OLSEN, M.K., 2016. Bariatric Surgery and Long-term Durability of Weight Loss. *JAMA surgery*, **151**(11), pp. 1046-1055.
- MADERNA, P., YONA, S., PERRETTI, M. and GODSON, C., 2005. Modulation of phagocytosis of apoptotic neutrophils by supernatant from dexamethasone-treated macrophages and annexin-derived peptide Ac(2-26). *Journal of immunology (Baltimore, Md.: 1950)*, **174**(6), pp. 3727-3733.
- MAEDA, N., SHIMOMURA, I., KISHIDA, K., NISHIZAWA, H., MATSUDA, M., NAGARETANI, H., FURUYAMA, N., KONDO, H., TAKAHASHI, M., ARITA, Y., KOMURO, R., OUCHI, N., KIHARA, S., TOCHINO, Y., OKUTOMI, K., HORIE, M., TAKEDA, S., AOYAMA, T., FUNAHASHI, T. and MATSUZAWA, Y., 2002. Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Nature medicine*, **8**(7), pp. 731-737.
- MAKKI, K., FROGUEL, P. and WOLOWCZUK, I., 2013. Adipose tissue in obesity-related inflammation and insulin resistance: cells, cytokines, and chemokines. *ISRN inflammation*, **2013**, pp. 139239.
- MANDRUP, S. and LANE, M.D., 1997. Regulating adipogenesis. *The Journal of biological chemistry*, **272**(9), pp. 5367-5370.
- MANDRUP-POULSEN, T., 1996. The role of interleukin-1 in the pathogenesis of IDDM. *Diabetologia*, **39**(9), pp. 1005-1029.

- MANNA, P. and JAIN, S.K., 2015. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metabolic syndrome and related disorders*, **13**(10), pp. 423-444.
- MANTZOROS, C.S., 2012. Leptin in relation to the lipodystrophy-associated metabolic syndrome. *Diabetes & metabolism journal*, **36**(3), pp. 181-189.
- MARROQUI, L., GONZALEZ, A., NECO, P., CABALLERO-GARRIDO, E., VIEIRA, E., RIPOLL, C., NADAL, A. and QUESADA, I., 2012. Role of leptin in the pancreatic beta-cell: effects and signaling pathways. *Journal of Molecular Endocrinology*, **49**(1), pp. R9-17.
- MASKREY, B.H., WHITFIELD, P.D., MEGSON, I.L. and ROSSI, A.G., 2011. Mechanisms of resolution of inflammation: A focus on cardiovascular disease. *Arteriosclerosis, Thrombosis, and Vascular Biology*, **31**(5), pp. 1001-1006.
- MASSON, O., CHAVEY, C., DRAY, C., MEULLE, A., DAVIAUD, D., QUILLIOT, D., MULLER, C., VALET, P. and LIAUDET-COOPMAN, E., 2009. LRP1 receptor controls adipogenesis and is up-regulated in human and mouse obese adipose tissue. *PLoS one*, **4**(10), pp. e7422.
- MATHIS, D., 2013. Immunological goings-on in visceral adipose tissue. *Cell metabolism*, **17**(6), pp. 851-859.
- MATTHEWS, V.B., ALLEN, T.L., RISIS, S., CHAN, M.H., HENSTRIDGE, D.C., WATSON, N., ZAFFINO, L.A., BABB, J.R., BOON, J., MEIKLE, P.J., JOWETT, J.B., WATT, M.J., JANSSON, J.O., BRUCE, C.R. and FEBBRAIO, M.A., 2010. Interleukin-6-deficient mice develop hepatic inflammation and systemic insulin resistance. *Diabetologia*, **53**(11), pp. 2431-2441.
- MAURY, E. and BRICHARD, S.M., 2010. Adipokine dysregulation, adipose tissue inflammation and metabolic syndrome. *Molecular and cellular endocrinology*, **314**(1), pp. 1-16.
- MCALLISTER, E.J., DHURANDHAR, N.V., KEITH, S.W., ARONNE, L.J., BARGER, J., BASKIN, M., BENCA, R.M., BIGGIO, J., BOGGIANO, M.M., EISENMANN, J.C., ELOBEID, M., FONTAINE, K.R., GLUCKMAN, P., HANLON, E.C., KATZMARZYK, P., PIETROBELLI, A., REDDEN, D.T., RUDEN, D.M., WANG, C., WATERLAND, R.A., WRIGHT, S.M. and ALLISON, D.B., 2009. Ten putative contributors to the obesity epidemic. *Critical reviews in food science and nutrition*, **49**(10), pp. 868-913.
- MEDZHITOV, R., 2008. Origin and physiological roles of inflammation. *Nature*, **454**(7203), pp. 428-435.
- MEEK, C.L., LEWIS, H.B., REIMANN, F., GRIBBLE, F.M. and PARK, A.J., 2016. The effect of bariatric surgery on gastrointestinal and pancreatic peptide hormones. *Peptides*, **77**, pp. 28-37.

- MEHLEM, A., HAGBERG, C.E., MUHL, L., ERIKSSON, U. and FALKEVALL, A., 2013. Imaging of neutral lipids by oil red O for analyzing the metabolic status in health and disease. *Nature protocols*, **8**(6), pp. 1149-1154.
- MELKI, V., HULLIN, F., MAZARGUIL, H., FAUVEL, J., RAGAB-THOMAS, J.M. and CHAP, H., 1994. Annexin I as a potential inhibitor of insulin receptor protein tyrosine kinase. *Biochemical and biophysical research communications*, **203**(2), pp. 813-819.
- METZ, C.N. and TRACEY, K.J., 2005. It takes nerve to dampen inflammation. *Nature immunology*, **6**(8), pp. 756-757.
- MILES, P.D., BARAK, Y., HE, W., EVANS, R.M. and OLEFSKY, J.M., 2000. Improved insulin-sensitivity in mice heterozygous for PPAR-gamma deficiency. *The Journal of clinical investigation*, **105**(3), pp. 287-292.
- MINOKOSHI, Y., KIM, Y.B., PERONI, O.D., FRYER, L.G., MULLER, C., CARLING, D. and KAHN, B.B., 2002. Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature*, **415**(6869), pp. 339-343.
- MISRA, A. and GARG, A., 2003. Clinical features and metabolic derangements in acquired generalized lipodystrophy: case reports and review of the literature. *Medicine*, **82**(2), pp. 129-146.
- MOEHLECKE, M., CANANI, L.H., SILVA, L.O., TRINDADE, M.R., FRIEDMAN, R. and LEITAO, C.B., 2016. Determinants of body weight regulation in humans. *Archives of endocrinology and metabolism*, **60**(2), pp. 152-162.
- MOITRA, J., MASON, M.M., OLIVE, M., KRYLOV, D., GAVRILOVA, O., MARCUS-SAMUELS, B., FEIGENBAUM, L., LEE, E., AOYAMA, T., ECKHAUS, M., REITMAN, M.L. and VINSON, C., 1998. Life without white fat: a transgenic mouse. *Genes & development*, **12**(20), pp. 3168-3181.
- MONAJEMI, H., ZHANG, L., LI, G., JENINGA, E.H., CAO, H., MAAS, M., BROUWER, C.B., KALKHOVEN, E., STROES, E., HEGELE, R.A. and LEFF, T., 2007. Familial partial lipodystrophy phenotype resulting from a single-base mutation in deoxyribonucleic acid-binding domain of peroxisome proliferator-activated receptor-gamma. *The Journal of clinical endocrinology and metabolism*, **92**(5), pp. 1606-1612.
- MONTEZ, J.M., SOUKAS, A., ASILMAZ, E., FAYZIKHODJAEVA, G., FANTUZZI, G. and FRIEDMAN, J.M., 2005. Acute leptin deficiency, leptin resistance, and the physiologic response to leptin withdrawal. *Proceedings of the National Academy of Sciences of the United States of America*, **102**(7), pp. 2537-2542.
- MOON, H.S., DALAMAGA, M., KIM, S.Y., POLYZOS, S.A., HAMNVIK, O.P., MAGKOS, F., PARUTHI, J. and MANTZOROS, C.S., 2013. Leptin's role in lipodystrophic and nonlipodystrophic insulin-resistant and diabetic individuals. *Endocrine reviews*, **34**(3), pp. 377-412.

- MOON, Y.S., SMAS, C.M., LEE, K., VILLENA, J.A., KIM, K.H., YUN, E.J. and SUL, H.S., 2002. Mice lacking paternally expressed Pref-1/Dlk1 display growth retardation and accelerated adiposity. *Molecular and cellular biology*, **22**(15), pp. 5585-5592.
- MORAND, E.F., HUTCHINSON, P., HARGREAVES, A., GOULDING, N.J., BOYCE, N.W. and HOLDSWORTH, S.R., 1995. Detection of intracellular lipocortin 1 in human leukocyte subsets. *Clinical immunology and immunopathology*, **76**(2), pp. 195-202.
- MORENO-CASTELLANOS, N., GUZMAN-RUIZ, R., CANO, D.A., MADRAZO-ATUTXA, A., PEINADO, J.R., PEREIRA-CUNILL, J.L., GARCIA-LUNA, P.P., MORALES-CONDE, S., SOCAS-MACIAS, M., VAZQUEZ-MARTINEZ, R., LEAL-CERRO, A. and MALAGON, M.M., 2016. The Effects of Bariatric Surgery-Induced Weight Loss on Adipose Tissue in Morbidly Obese Women Depends on the Initial Metabolic Status. *Obesity Surgery*, **26**(8), pp. 1757-1767.
- MORI, H., HANADA, R., HANADA, T., AKI, D., MASHIMA, R., NISHINAKAMURA, H., TORISU, T., CHIEN, K.R., YASUKAWA, H. and YOSHIMURA, A., 2004. Socs3 deficiency in the brain elevates leptin sensitivity and confers resistance to diet-induced obesity. *Nature medicine*, **10**(7), pp. 739-743.
- MORI, T., SAKAUE, H., IGUCHI, H., GOMI, H., OKADA, Y., TAKASHIMA, Y., NAKAMURA, K., NAKAMURA, T., YAMAUCHI, T., KUBOTA, N., KADOWAKI, T., MATSUKI, Y., OGAWA, W., HIRAMATSU, R. and KASUGA, M., 2005. Role of Kruppel-like factor 15 (KLF15) in transcriptional regulation of adipogenesis. *The Journal of biological chemistry*, **280**(13), pp. 12867-12875.
- MRAZ, M. and HALUZIK, M., 2014. The role of adipose tissue immune cells in obesity and low-grade inflammation. *The Journal of endocrinology*, **222**(3), pp. R113-27.
- MUELLER, C., WEAVER, V., VANDEN HEUVEL, J.P., AUGUST, A. and CANTORNA, M.T., 2003. Peroxisome proliferator-activated receptor gamma ligands attenuate immunological symptoms of experimental allergic asthma. *Archives of Biochemistry and Biophysics*, **418**(2), pp. 186-196.
- MULLER, M.J., ENDERLE, J., POURHASSAN, M., BRAUN, W., EGGELING, B., LAGERPUSCH, M., GLUER, C.C., KEHAYIAS, J.J., KIOSZ, D. and BOSY-WESTPHAL, A., 2015. Metabolic adaptation to caloric restriction and subsequent refeeding: the Minnesota Starvation Experiment revisited. *The American Journal of Clinical Nutrition*, **102**(4), pp. 807-819.
- MURAILLE, E., LEO, O. and MOSER, M., 2014. TH1/TH2 paradigm extended: macrophage polarization as an unappreciated pathogen-driven escape mechanism? *Frontiers in immunology*, **5**, pp. 603.
- MURAKAMI, K., BUJO, H., UNOKI, H. and SAITO, Y., 2007. Effect of PPARalpha activation of macrophages on the secretion of inflammatory cytokines in cultured adipocytes. *European journal of pharmacology*, **561**(1-3), pp. 206-213.

- MURPHY, K.G. and BLOOM, S.R., 2006. Gut hormones and the regulation of energy homeostasis. *Nature*, **444**(7121), pp. 854-859.
- MURRI, M., INSENER, M., BERNAL-LOPEZ, M.R., PEREZ-MARTINEZ, P., ESCOBAR-MORREALE, H.F. and TINAHONES, F.J., 2013. Proteomic analysis of visceral adipose tissue in pre-obese patients with type 2 diabetes. *Molecular and cellular endocrinology*, **376**(1-2), pp. 99-106.
- NAGELL, C.F., WETTERGREN, A., PEDERSEN, J.F., MORTENSEN, D. and HOLST, J.J., 2004. Glucagon-like peptide-2 inhibits antral emptying in man, but is not as potent as glucagon-like peptide-1. *Scandinavian Journal of Gastroenterology*, **39**(4), pp. 353-358.
- NAPOLITANO, A., LOWELL, B.B., DAMM, D., LEIBEL, R.L., RAVUSSIN, E., JIMERSON, D.C., LESEM, M.D., VAN DYKE, D.C., DALY, P.A. and CHATIS, P., 1994. Concentrations of adipsin in blood and rates of adipsin secretion by adipose tissue in humans with normal, elevated and diminished adipose tissue mass. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*, **18**(4), pp. 213-218.
- NASLUND, E., BARKELING, B., KING, N., GUTNIAK, M., BLUNDELL, J.E., HOLST, J.J., ROSSNER, S. and HELLSTROM, P.M., 1999. Energy intake and appetite are suppressed by glucagon-like peptide-1 (GLP-1) in obese men. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*, **23**(3), pp. 304-311.
- NATARAJAN, R., HAGMAN, S., HAMALAINEN, M., LEPPANEN, T., DASTIDAR, P., MOILANEN, E. and ELOVAARA, I., 2015. Adipsin Is Associated with Multiple Sclerosis: A Follow-Up Study of Adipokines. *Multiple sclerosis international*, **2015**, pp. 371734.
- NAUSHEEN, S., SHAH, I.H., PEZESHKI, A., SIGALET, D.L. and CHELIKANI, P.K., 2013. Effects of sleeve gastrectomy and ileal transposition, alone and in combination, on food intake, body weight, gut hormones, and glucose metabolism in rats. *American journal of physiology. Endocrinology and metabolism*, **305**(4), pp. E507-18.
- NAWROCKI, A.R., RAJALA, M.W., TOMAS, E., PAJVANI, U.B., SAHA, A.K., TRUMBAUER, M.E., PANG, Z., CHEN, A.S., RUDERMAN, N.B., CHEN, H., ROSSETTI, L. and SCHERER, P.E., 2006. Mice lacking adiponectin show decreased hepatic insulin sensitivity and reduced responsiveness to peroxisome proliferator-activated receptor gamma agonists. *The Journal of biological chemistry*, **281**(5), pp. 2654-2660.
- NEAL, J.W. and CLIPSTONE, N.A., 2002. Calcineurin mediates the calcium-dependent inhibition of adipocyte differentiation in 3T3-L1 cells. *The Journal of biological chemistry*, **277**(51), pp. 49776-49781.

- NEEL, J.V., 1999. The "thrifty genotype" in 1998. *Nutrition reviews*, **57**(5 Pt 2), pp. S2-9.
- NEELS, J.G., 2013. A role for 5-lipoxygenase products in obesity-associated inflammation and insulin resistance. *Adipocyte*, **2**(4), pp. 262-265.
- NETTO, B.D., BETTINI, S.C., CLEMENTE, A.P., FERREIRA, J.P., BORITZA, K., SOUZA SDE, F., VON DER HEYDE, M.E., EARTHMAN, C.P. and DAMASO, A.R., 2015. Roux-en-Y gastric bypass decreases pro-inflammatory and thrombotic biomarkers in individuals with extreme obesity. *Obesity Surgery*, **25**(6), pp. 1010-1018.
- NGUYEN, M.T., FAVELYUKIS, S., NGUYEN, A.K., REICHART, D., SCOTT, P.A., JENN, A., LIU-BRYAN, R., GLASS, C.K., NEELS, J.G. and OLEFSKY, J.M., 2007. A subpopulation of macrophages infiltrates hypertrophic adipose tissue and is activated by free fatty acids via Toll-like receptors 2 and 4 and JNK-dependent pathways. *The Journal of biological chemistry*, **282**(48), pp. 35279-35292.
- NIELSEN, T.S., JESSEN, N., JORGENSEN, J.O., MOLLER, N. and LUND, S., 2014. Dissecting adipose tissue lipolysis: molecular regulation and implications for metabolic disease. *Journal of Molecular Endocrinology*, **52**(3), pp. R199-222.
- NIJHUIS, J., RENSEN, S.S., SLAATS, Y., VAN DIELEN, F.M., BUURMAN, W.A. and GREVE, J.W., 2009. Neutrophil activation in morbid obesity, chronic activation of acute inflammation. *Obesity (Silver Spring, Md.)*, **17**(11), pp. 2014-2018.
- NING, X., HE, J., SHI, X. and YANG, G., 2016. Regulation of Adipogenesis by Quinine through the ERK/S6 Pathway. *International journal of molecular sciences*, **17**(4), pp. 504.
- NISHIMOTO, S., FUKUDA, D., HIGASHIKUNI, Y., TANAKA, K., HIRATA, Y., MURATA, C., KIM-KANEYAMA, J.R., SATO, F., BANDO, M., YAGI, S., SOEKI, T., HAYASHI, T., IMOTO, I., SAKAUE, H., SHIMABUKURO, M. and SATA, M., 2016. Obesity-induced DNA released from adipocytes stimulates chronic adipose tissue inflammation and insulin resistance. *Science advances*, **2**(3), pp. e1501332.
- NISHIMURA, S., MANABE, I. and NAGAI, R., 2009. Adipose tissue inflammation in obesity and metabolic syndrome. *Discovery medicine*, **8**(41), pp. 55-60.
- NORRELUND, H., NAIR, K.S., JORGENSEN, J.O., CHRISTIANSEN, J.S. and MOLLER, N., 2001. The protein-retaining effects of growth hormone during fasting involve inhibition of muscle-protein breakdown. *Diabetes*, **50**(1), pp. 96-104.
- NOUSHMEHR, H., D'AMICO, E., FARILLA, L., HUI, H., WAWROWSKY, K.A., MLYNARSKI, W., DORIA, A., ABUMRAD, N.A. and PERFETTI, R., 2005. Fatty acid translocase (FAT/CD36) is localized on insulin-containing granules in human pancreatic beta-cells and mediates fatty acid effects on insulin secretion. *Diabetes*, **54**(2), pp. 472-481.

- NTAMBI, J.M. and TAKOVA, T., 1996. Role of Ca²⁺ in the early stages of murine adipocyte differentiation as evidenced by calcium mobilizing agents. *Differentiation; research in biological diversity*, **60**(3), pp. 151-158.
- NUGENT, C., PRINS, J.B., WHITEHEAD, J.P., SAVAGE, D., WENTWORTH, J.M., CHATTERJEE, V.K. and O'RAHILLY, S., 2001. Potentiation of glucose uptake in 3T3-L1 adipocytes by PPAR gamma agonists is maintained in cells expressing a PPAR gamma dominant-negative mutant: evidence for selectivity in the downstream responses to PPAR gamma activation. *Molecular endocrinology (Baltimore, Md.)*, **15**(10), pp. 1729-1738.
- NUTTALL, F.Q., 2015. Body Mass Index: Obesity, BMI, and Health: A Critical Review. *Nutrition today*, **50**(3), pp. 117-128.
- NYE, C.K., HANSON, R.W. and KALHAN, S.C., 2008. Glyceroneogenesis is the dominant pathway for triglyceride glycerol synthesis in vivo in the rat. *The Journal of biological chemistry*, **283**(41), pp. 27565-27574.
- OBICI, S., ZHANG, B.B., KARKANIAS, G. and ROSSETTI, L., 2002. Hypothalamic insulin signaling is required for inhibition of glucose production. *Nature medicine*, **8**(12), pp. 1376-1382.
- OBREGON, M.J., 2014. Adipose tissues and thyroid hormones. *Frontiers in physiology*, **5**, pp. 479.
- OFEL, F., HUREL, S., NEWKIRK, J., SOPWITH, M. and TAYLOR, R., 1996. Effects of an engineered human anti-TNF-alpha antibody (CDP571) on insulin sensitivity and glycemic control in patients with NIDDM. *Diabetes*, **45**(7), pp. 881-885.
- OHNISHI, M., TOKUDA, M., MASAKI, T., FUJIMURA, T., TAI, Y., ITANO, T., MATSUI, H., ISHIDA, T., KONISHI, R. and TAKAHARA, J., 1995. Involvement of annexin-I in glucose-induced insulin secretion in rat pancreatic islets. *Endocrinology*, **136**(6), pp. 2421-2426.
- OISHI, Y., MANABE, I., TOBE, K., TSUSHIMA, K., SHINDO, T., FUJII, K., NISHIMURA, G., MAEMURA, K., YAMAUCHI, T., KUBOTA, N., SUZUKI, R., KITAMURA, T., AKIRA, S., KADOWAKI, T. and NAGAI, R., 2005. Kruppel-like transcription factor KLF5 is a key regulator of adipocyte differentiation. *Cell metabolism*, **1**(1), pp. 27-39.
- OKI, K., YAMANE, K., KAMEI, N., NOJIMA, H. and KOHNO, N., 2007. Circulating visfatin level is correlated with inflammation, but not with insulin resistance. *Clinical endocrinology*, **67**(5), pp. 796-800.
- OLLER DO NASCIMENTO, C.M., RIBEIRO, E.B. and OYAMA, L.M., 2009. Metabolism and secretory function of white adipose tissue: effect of dietary fat. *Anais da Academia Brasileira de Ciencias*, **81**(3), pp. 453-466.

- OLSON, A.L., 2012. Regulation of GLUT4 and Insulin-Dependent Glucose Flux. *ISRN molecular biology*, **2012**, pp. 856987.
- ORAL, E.A., SIMHA, V., RUIZ, E., ANDEWELT, A., PREMKUMAR, A., SNELL, P., WAGNER, A.J., DEPAOLI, A.M., REITMAN, M.L., TAYLOR, S.I., GORDEN, P. and GARG, A., 2002. Leptin-replacement therapy for lipodystrophy. *The New England journal of medicine*, **346**(8), pp. 570-578.
- OSBORN, O. and OLEFSKY, J.M., 2012. The cellular and signaling networks linking the immune system and metabolism in disease. *Nature medicine*, **18**(3), pp. 363-374.
- OUCHI, N., KIHARA, S., ARITA, Y., MAEDA, K., KURIYAMA, H., OKAMOTO, Y., HOTTA, K., NISHIDA, M., TAKAHASHI, M., NAKAMURA, T., YAMASHITA, S., FUNAHASHI, T. and MATSUZAWA, Y., 1999. Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation*, **100**(25), pp. 2473-2476.
- OUCHI, N., KIHARA, S., ARITA, Y., NISHIDA, M., MATSUYAMA, A., OKAMOTO, Y., ISHIGAMI, M., KURIYAMA, H., KISHIDA, K., NISHIZAWA, H., HOTTA, K., MURAGUCHI, M., OHMOTO, Y., YAMASHITA, S., FUNAHASHI, T. and MATSUZAWA, Y., 2001. Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages. *Circulation*, **103**(8), pp. 1057-1063.
- OUCHI, N., KIHARA, S., ARITA, Y., OKAMOTO, Y., MAEDA, K., KURIYAMA, H., HOTTA, K., NISHIDA, M., TAKAHASHI, M., MURAGUCHI, M., OHMOTO, Y., NAKAMURA, T., YAMASHITA, S., FUNAHASHI, T. and MATSUZAWA, Y., 2000. Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF-kappaB signaling through a cAMP-dependent pathway. *Circulation*, **102**(11), pp. 1296-1301.
- OUCHI, N., PARKER, J.L., LUGUS, J.J. and WALSH, K., 2011. Adipokines in inflammation and metabolic disease. *Nature reviews.Immunology*, **11**(2), pp. 85-97.
- OZATA, M., OZDEMIR, I.C. and LICINIO, J., 1999. Human leptin deficiency caused by a missense mutation: multiple endocrine defects, decreased sympathetic tone, and immune system dysfunction indicate new targets for leptin action, greater central than peripheral resistance to the effects of leptin, and spontaneous correction of leptin-mediated defects. *The Journal of clinical endocrinology and metabolism*, **84**(10), pp. 3686-3695.
- PACENTI, M., BARZON, L., FAVARETTO, F., FINCATI, K., ROMANO, S., MILAN, G., VETTOR, R. and PALU, G., 2006. Microarray analysis during adipogenesis identifies new genes altered by antiretroviral drugs. *AIDS (London, England)*, **20**(13), pp. 1691-1705.
- PALANIVEL, R., MAIDA, A., LIU, Y. and SWEENEY, G., 2006. Regulation of insulin signalling, glucose uptake and metabolism in rat skeletal muscle cells upon prolonged exposure to resistin. *Diabetologia*, **49**(1), pp. 183-190.

- PAN, B., KONG, J., JIN, J., KONG, J., HE, Y., DONG, S., JI, L., LIU, D., HE, D., KONG, L., JIN, D.K., WILLARD, B., PENNATHUR, S. and ZHENG, L., 2016. A novel anti-inflammatory mechanism of high density lipoprotein through up-regulating annexin A1 in vascular endothelial cells. *Biochimica et biophysica acta*, **1861**(6), pp. 501-512.
- PARDINA, E., FERRER, R., BAENA-FUSTEGUERAS, J.A., RIVERO, J., LECUBE, A., FORT, J.M., VARGAS, V., CATALAN, R. and PEINADO-ONSURBE, J., 2012. Only C-reactive protein, but not TNF-alpha or IL6, reflects the improvement in inflammation after bariatric surgery. *Obesity Surgery*, **22**(1), pp. 131-139.
- PARDINI, V.C., VICTORIA, I.M., ROCHA, S.M., ANDRADE, D.G., ROCHA, A.M., PIERONI, F.B., MILAGRES, G., PURISCH, S. and VELHO, G., 1998. Leptin levels, beta-cell function, and insulin sensitivity in families with congenital and acquired generalized lipoatrophic diabetes. *The Journal of clinical endocrinology and metabolism*, **83**(2), pp. 503-508.
- PARRA-ROJAS, I., DEL MORAL-HERNANDEZ, O., SALGADO-BERNABE, A.B., GUZMAN-GUZMAN, I.P., SALGADO-GOYTIA, L. and MUNOZ-VALLE, J.F., 2013. Adenovirus-36 seropositivity and its relation with obesity and metabolic profile in children. *International journal of endocrinology*, **2013**, pp. 463194.
- PATEL, D.M., AHMAD, S.F., WEISS, D.G., GERKE, V. and KUZNETSOV, S.A., 2011. Annexin A1 is a new functional linker between actin filaments and phagosomes during phagocytosis. *Journal of cell science*, **124**(Pt 4), pp. 578-588.
- PATEL, H.B., KORNERUP, K.N., SAMPAIO, A.L., D'ACQUISTO, F., SEED, M.P., GIROL, A.P., GRAY, M., PITZALIS, C., OLIANI, S.M. and PERRETTI, M., 2012. The impact of endogenous annexin A1 on glucocorticoid control of inflammatory arthritis. *Annals of the Rheumatic Diseases*, **71**(11), pp. 1872-1880.
- PATEL, N.G., HOLDER, J.C., SMITH, S.A., KUMAR, S. and EGGO, M.C., 2003. Differential regulation of lipogenesis and leptin production by independent signaling pathways and rosiglitazone during human adipocyte differentiation. *Diabetes*, **52**(1), pp. 43-50.
- PATIL, M., SHARMA, B.K., ELATTAR, S., CHANG, J., KAPIL, S., YUAN, J. and SATYANARAYANA, A., 2017. Id1 Promotes Obesity by Suppressing Brown Adipose Thermogenesis and White Adipose Browning. *Diabetes*, **66**(6), pp. 1611-1625.
- PEACEY, S.R., WRIGHT, D., AYE, M. and MOISEY, R., 2012. Glucocorticoid replacement therapy and fibrinolysis in patients with hypopituitarism. *Clinical endocrinology*, **77**(1), pp. 94-98.
- PEETERS, A., BARENDREGT, J.J., WILLEKENS, F., MACKENBACH, J.P., AL MAMUN, A., BONNEUX, L. and NEDCOM, THE NETHERLANDS EPIDEMIOLOGY AND DEMOGRAPHY COMPRESSION OF MORBIDITY RESEARCH GROUP, 2003. Obesity in adulthood and its consequences for life expectancy: a life-table analysis. *Annals of Internal Medicine*, **138**(1), pp. 24-32.

- PERMANA, P.A., NAIR, S., LEE, Y.H., LUCZY-BACHMAN, G., VOZAROVA DE COURTEN, B. and TATARANNI, P.A., 2004. Subcutaneous abdominal preadipocyte differentiation in vitro inversely correlates with central obesity. *American journal of physiology. Endocrinology and metabolism*, **286**(6), pp. E958-62.
- PERRETTI, M. and D'ACQUISTO, F., 2009. Annexin A1 and glucocorticoids as effectors of the resolution of inflammation. *Nature reviews. Immunology*, **9**(1), pp. 62-70.
- PERRETTI, M. and FLOWER, R.J., 1996. Measurement of lipocortin 1 levels in murine peripheral blood leukocytes by flow cytometry: modulation by glucocorticoids and inflammation. *British journal of pharmacology*, **118**(3), pp. 605-610.
- PERRETTI, M., FLOWER, R.J. and GOULDING, N.J., 1993. The ability of murine leukocytes to bind lipocortin 1 is lost during acute inflammation. *Biochemical and biophysical research communications*, **192**(2), pp. 345-350.
- PERRETTI, M. and SOLITO, E., 2004. Annexin 1 and neutrophil apoptosis. *Biochemical Society transactions*, **32**(Pt3), pp. 507-510.
- PERUSSE, L., CHAGNON, Y.C., WEISNAGEL, S.J., RANKINEN, T., SNYDER, E., SANDS, J. and BOUCHARD, C., 2001. The human obesity gene map: the 2000 update. *Obesity research*, **9**(2), pp. 135-169.
- PETRI, M.H., LAGUNA-FERNANDEZ, A., GONZALEZ-DIEZ, M., PAULSSON-BERNE, G., HANSSON, G.K. and BACK, M., 2015. The role of the FPR2/ALX receptor in atherosclerosis development and plaque stability. *Cardiovascular research*, **105**(1), pp. 65-74.
- PETROVIC, N., WALDEN, T.B., SHABALINA, I.G., TIMMONS, J.A., CANNON, B. and NEDERGAARD, J., 2010. Chronic peroxisome proliferator-activated receptor gamma (PPARgamma) activation of epididymally derived white adipocyte cultures reveals a population of thermogenically competent, UCP1-containing adipocytes molecularly distinct from classic brown adipocytes. *The Journal of biological chemistry*, **285**(10), pp. 7153-7164.
- PFLUGER, P.T., KABRA, D.G., AICHLER, M., SCHRIEVER, S.C., PFUHLMANN, K., GARCIA, V.C., LEHTI, M., WEBER, J., KUTSCHKE, M., ROZMAN, J., ELROD, J.W., HEVENER, A.L., FEUCHTINGER, A., HRABE DE ANGELIS, M., WALCH, A., ROLLMANN, S.M., ARONOW, B.J., MULLER, T.D., PEREZ-TILVE, D., JASTROCH, M., DE LUCA, M., MOLKENTIN, J.D. and TSCHOP, M.H., 2015. Calcineurin Links Mitochondrial Elongation with Energy Metabolism. *Cell metabolism*, **22**(5), pp. 838-850.
- PHAN, J. and REUE, K., 2005. Lipin, a lipodystrophy and obesity gene. *Cell metabolism*, **1**(1), pp. 73-83.
- PICARD, F. and AUWERX, J., 2002. PPAR(gamma) and glucose homeostasis. *Annual Review of Nutrition*, **22**, pp. 167-197.

- PIETRANI, N.T., FERREIRA, C.N., RODRIGUES, K.F., PERUCCI, L.O., CARNEIRO, F.S., BOSCO, A.A., OLIVEIRA, M.C., PEREIRA, S.S., TEIXEIRA, A.L., ALVAREZ-LEITE, J.I., FERREIRA, A.V., SOUSA, L.P. and GOMES, K.B., 2018. Proresolving protein Annexin A1: The role in type 2 diabetes mellitus and obesity. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*, **103**, pp. 482-489.
- PLATA-SALAMAN, C.R., 1994. Meal patterns in response to the intracerebroventricular administration of interleukin-1 beta in rats. *Physiology & Behavior*, **55**(4), pp. 727-733.
- PLUM, L., SCHUBERT, M. and BRUNING, J.C., 2005. The role of insulin receptor signaling in the brain. *Trends in endocrinology and metabolism: TEM*, **16**(2), pp. 59-65.
- POLETO, A.C., DAVID-SILVA, A., YAMAMOTO, A.P., MACHADO, U.F. and FURUYA, D.T., 2015. Reduced Slc2a4/GLUT4 expression in subcutaneous adipose tissue of monosodium glutamate obese mice is recovered after atorvastatin treatment. *Diabetology & metabolic syndrome*, **7**, pp. 18-015-0015-6. eCollection 2015.
- PRADHAN, A.D., MANSON, J.E., RIFAI, N., BURING, J.E. and RIDKER, P.M., 2001. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *Jama*, **286**(3), pp. 327-334.
- PRICOLA, K.L., KUHN, N.Z., HALEEM-SMITH, H., SONG, Y. and TUAN, R.S., 2009. Interleukin-6 maintains bone marrow-derived mesenchymal stem cell stemness by an ERK1/2-dependent mechanism. *Journal of cellular biochemistry*, **108**(3), pp. 577-588.
- PRUSTY, D., PARK, B.H., DAVIS, K.E. and FARMER, S.R., 2002. Activation of MEK/ERK signaling promotes adipogenesis by enhancing peroxisome proliferator-activated receptor gamma (PPARgamma) and C/EBPalpha gene expression during the differentiation of 3T3-L1 preadipocytes. *The Journal of biological chemistry*, **277**(48), pp. 46226-46232.
- PUPJALIS, D., GOETSCH, J., KOTTAS, D.J., GERKE, V. and RESCHER, U., 2011. Annexin A1 released from apoptotic cells acts through formyl peptide receptors to dampen inflammatory monocyte activation via JAK/STAT/SOCS signalling. *EMBO molecular medicine*, **3**(2), pp. 102-114.
- PURVIS, G.S.D., COLLINO, M., LOIOLA, R.A., BARAGETTI, A., CHIAZZA, F., BROVELLI, M., SHEIKH, M.H., COLLOTTA, D., CENTO, A., MASTROCOLA, R., ARAGNO, M., CUTRIN, J.C., REUTELINGSPERGER, C., GRIGORE, L., CATAPANO, A.L., YAQOOB, M.M., NORATA, G.D., SOLITO, E. and THIEMERMANN, C., 2019. Identification of AnnexinA1 as an Endogenous Regulator of RhoA, and Its Role in the Pathophysiology and Experimental Therapy of Type-2 Diabetes. *Frontiers in immunology*, **10**, pp. 571.
- PURVIS, G.S.D., SOLITO, E. and THIEMERMANN, C., 2019. Annexin-A1: Therapeutic Potential in Microvascular Disease. *Frontiers in immunology*, **10**, pp. 938.

- RABIET, M.J., HUET, E. and BOULAY, F., 2007. The N-formyl peptide receptors and the anaphylatoxin C5a receptors: an overview. *Biochimie*, **89**(9), pp. 1089-1106.
- RAJALA, M.W., OBICI, S., SCHERER, P.E. and ROSSETTI, L., 2003. Adipose-derived resistin and gut-derived resistin-like molecule-beta selectively impair insulin action on glucose production. *The Journal of clinical investigation*, **111**(2), pp. 225-230.
- RAKOTOARIVELO, V., LACRAZ, G., MAYHUE, M., BROWN, C., ROTTEMBOURG, D., FRADETTE, J., ILANGUMARAN, S., MENENDEZ, A., LANGLOIS, M.F. and RAMANATHAN, S., 2018. Inflammatory Cytokine Profiles in Visceral and Subcutaneous Adipose Tissues of Obese Patients Undergoing Bariatric Surgery Reveal Lack of Correlation With Obesity or Diabetes. *EBioMedicine*, **30**, pp. 237-247.
- RAMIREZ-ZACARIAS, J.L., CASTRO-MUNOZLEDO, F. and KURI-HARCUC, W., 1992. Quantitation of adipose conversion and triglycerides by staining intracytoplasmic lipids with Oil red O. *Histochemistry*, **97**(6), pp. 493-497.
- RAMOS-LOPEZ, O., SAMBLAS, M., MILAGRO, F.I., RIEZU-BOJ, J.I., CRUJEIRAS, A.B., MARTINEZ, J.A. and PROJECT, M., 2018. Circadian gene methylation profiles are associated with obesity, metabolic disturbances and carbohydrate intake. *Chronobiology international*, **35**(7), pp. 969-981.
- RANKIN, W. and WITTERT, G., 2015. Anti-obesity drugs. *Current opinion in lipidology*, **26**(6), pp. 536-543.
- RAVUSSIN, E., LILLIOJA, S., KNOWLER, W.C., CHRISTIN, L., FREYMOND, D., ABBOTT, W.G., BOYCE, V., HOWARD, B.V. and BOGARDUS, C., 1988. Reduced rate of energy expenditure as a risk factor for body-weight gain. *The New England journal of medicine*, **318**(8), pp. 467-472.
- REDDY, N.L., TAN, B.K., BARBER, T.M. and RANDEVA, H.S., 2014. Brown adipose tissue: endocrine determinants of function and therapeutic manipulation as a novel treatment strategy for obesity. *BMC obesity*, **1**, pp. 13-014-0013-5. eCollection 2014.
- REIDY, S.P. and WEBER, J., 2000. Leptin: an essential regulator of lipid metabolism. *Comparative biochemistry and physiology. Part A, Molecular & integrative physiology*, **125**(3), pp. 285-298.
- REN, D., COLLINGWOOD, T.N., REBAR, E.J., WOLFFE, A.P. and CAMP, H.S., 2002. PPARgamma knockdown by engineered transcription factors: exogenous PPARgamma2 but not PPARgamma1 reactivates adipogenesis. *Genes & development*, **16**(1), pp. 27-32.
- REVOLLO, J.R., KORNER, A., MILLS, K.F., SATOH, A., WANG, T., GARTEN, A., DASGUPTA, B., SASAKI, Y., WOLBERGER, C., TOWNSEND, R.R., MILBRANDT, J., KIESS, W. and IMAI, S., 2007. Nampt/PBEF/Visfatin regulates insulin secretion in beta cells as a systemic NAD biosynthetic enzyme. *Cell metabolism*, **6**(5), pp. 363-375.

- RIBEIRO, M.O., CARVALHO, S.D., SCHULTZ, J.J., CHIELLINI, G., SCANLAN, T.S., BIANCO, A.C. and BRENT, G.A., 2001. Thyroid hormone--sympathetic interaction and adaptive thermogenesis are thyroid hormone receptor isoform--specific. *The Journal of clinical investigation*, **108**(1), pp. 97-105.
- RICCO, R.C., RICCO, R.G., QUELUZ, M.C., DE PAULA, M.T.S., ATIQUÉ, P.V., CUSTODIO, R.J., TOURINHO FILHO, H., DEL ROIO LIBERATORI, R., Jr and MARTINELLI, C.E., Jr, 2018. IGF-1R mRNA expression is increased in obese children. *Growth hormone & IGF research : official journal of the Growth Hormone Research Society and the International IGF Research Society*, **39**, pp. 1-5.
- RICKLIN, D., HAJISHENGALLIS, G., YANG, K. and LAMBRIS, J.D., 2010. Complement: a key system for immune surveillance and homeostasis. *Nature immunology*, **11**(9), pp. 785-797.
- RICQUIER, D., 2006. Fundamental mechanisms of thermogenesis. *Comptes rendus biologies*, **329**(8), pp. 578-86; discussion 653-5.
- RISTOW, M., MULLER-WIELAND, D., PFEIFFER, A., KRONE, W. and KAHN, C.R., 1998. Obesity associated with a mutation in a genetic regulator of adipocyte differentiation. *The New England journal of medicine*, **339**(14), pp. 953-959.
- ROBIDOUX, J., MARTIN, T.L. and COLLINS, S., 2004. Beta-adrenergic receptors and regulation of energy expenditure: a family affair. *Annual Review of Pharmacology and Toxicology*, **44**, pp. 297-323.
- RODER, M.E., PORTE, D., Jr, SCHWARTZ, R.S. and KAHN, S.E., 1998. Disproportionately elevated proinsulin levels reflect the degree of impaired B cell secretory capacity in patients with noninsulin-dependent diabetes mellitus. *The Journal of clinical endocrinology and metabolism*, **83**(2), pp. 604-608.
- ROJANO-RODRIGUEZ, M.E., VALENZUELA-SALAZAR, C., CARDENAS-LAILSON, L.E., ROMERO LOERA, L.S., TORRES-OLALDE, M. and MORENO-PORTILLO, M., 2014. C-reactive protein level in morbidly obese patients before and after bariatric surgery. *Revista de gastroenterologia de Mexico*, **79**(2), pp. 90-95.
- RONTI, T., LUPATTELLI, G. and MANNARINO, E., 2006. The endocrine function of adipose tissue: an update. *Clinical endocrinology*, **64**(4), pp. 355-365.
- ROSEBERRY, A.G., LIU, H., JACKSON, A.C., CAI, X. and FRIEDMAN, J.M., 2004. Neuropeptide Y-mediated inhibition of proopiomelanocortin neurons in the arcuate nucleus shows enhanced desensitization in ob/ob mice. *Neuron*, **41**(5), pp. 711-722.
- ROSEN, E., EGUCHI, J. and XU, Z., 2009. Transcriptional targets in adipocyte biology. *Expert opinion on therapeutic targets*, **13**(8), pp. 975-986.
- ROSEN, E.D. and SPIEGELMAN, B.M., 2014. What we talk about when we talk about fat. *Cell*, **156**(1-2), pp. 20-44.

- ROSEN, E.D. and SPIEGELMAN, B.M., 2006. Adipocytes as regulators of energy balance and glucose homeostasis. *Nature*, **444**(7121), pp. 847-853.
- ROSENBAUM, M., MURPHY, E.M., HEYMSFIELD, S.B., MATTHEWS, D.E. and LEIBEL, R.L., 2002. Low dose leptin administration reverses effects of sustained weight-reduction on energy expenditure and circulating concentrations of thyroid hormones. *The Journal of clinical endocrinology and metabolism*, **87**(5), pp. 2391-2394.
- ROSS, S.E., HEMATI, N., LONGO, K.A., BENNETT, C.N., LUCAS, P.C., ERICKSON, R.L. and MACDOUGALD, O.A., 2000. Inhibition of adipogenesis by Wnt signaling. *Science (New York, N.Y.)*, **289**(5481), pp. 950-953.
- ROTHWELL, N.J. and STOCK, M.J., 1979. A role for brown adipose tissue in diet-induced thermogenesis. *Nature*, **281**(5726), pp. 31-35.
- ROTTER, V., NAGAEV, I. and SMITH, U., 2003. Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor-alpha, overexpressed in human fat cells from insulin-resistant subjects. *The Journal of biological chemistry*, **278**(46), pp. 45777-45784.
- RUAN, H., MILES, P.D., LADD, C.M., ROSS, K., GOLUB, T.R., OLEFSKY, J.M. and LODISH, H.F., 2002. Profiling gene transcription in vivo reveals adipose tissue as an immediate target of tumor necrosis factor-alpha: implications for insulin resistance. *Diabetes*, **51**(11), pp. 3176-3188.
- RUPNICK, M.A., PANIGRAHY, D., ZHANG, C.Y., DALLABRIDA, S.M., LOWELL, B.B., LANGER, R. and FOLKMAN, M.J., 2002. Adipose tissue mass can be regulated through the vasculature. *Proceedings of the National Academy of Sciences of the United States of America*, **99**(16), pp. 10730-10735.
- RUSNAK, F. and MERTZ, P., 2000. Calcineurin: form and function. *Physiological Reviews*, **80**(4), pp. 1483-1521.
- RYDEN, M. and ARNER, P., 2007. Tumour necrosis factor-alpha in human adipose tissue -- from signalling mechanisms to clinical implications. *Journal of internal medicine*, **262**(4), pp. 431-438.
- SADDI-ROSA, P., OLIVEIRA, C.S., GIUFFRIDA, F.M. and REIS, A.F., 2010. Visfatin, glucose metabolism and vascular disease: a review of evidence. *Diabetology & metabolic syndrome*, **2**, pp. 21-5996-2-21. eCollection 2010.
- SAELY, C.H., GEIGER, K. and DREXEL, H., 2012. Brown versus white adipose tissue: a mini-review. *Gerontology*, **58**(1), pp. 15-23.
- SAKAGUCHI, M., FUJISAKA, S., CAI, W., WINNAY, J.N., KONISHI, M., O'NEILL, B.T., LI, M., GARCIA-MARTIN, R., TAKAHASHI, H., HU, J., KULKARNI, R.N. and KAHN, C.R., 2017. Adipocyte Dynamics and Reversible Metabolic Syndrome in Mice

with an Inducible Adipocyte-Specific Deletion of the Insulin Receptor. *Cell metabolism*, **25**(2), pp. 448-462.

SAKAMOTO, T., REPASKY, W.T., UCHIDA, K., HIRATA, A. and HIRATA, F., 1996. Modulation of cell death pathways to apoptosis and necrosis of H₂O₂-treated rat thymocytes by lipocortin I. *Biochemical and biophysical research communications*, **220**(3), pp. 643-647.

SALEH, J., SNIDERMAN, A.D. and CIANFLONE, K., 1999. Regulation of Plasma fatty acid metabolism. *Clinica chimica acta; international journal of clinical chemistry*, **286**(1-2), pp. 163-180.

SAMARAS, K., BOTELHO, N.K., CHISHOLM, D.J. and LORD, R.V., 2010. Subcutaneous and visceral adipose tissue gene expression of serum adipokines that predict type 2 diabetes. *Obesity (Silver Spring, Md.)*, **18**(5), pp. 884-889.

SAMS, V.G., BLACKLEDGE, C., WIJAYATUNGA, N., BARLOW, P., MANCINI, M., MANCINI, G. and MOUSTAID-MOUSSA, N., 2016. Effect of bariatric surgery on systemic and adipose tissue inflammation. *Surgical endoscopy*, **30**(8), pp. 3499-3504.

SANTOS-ALVAREZ, J., GOBERNA, R. and SANCHEZ-MARGALET, V., 1999. Human leptin stimulates proliferation and activation of human circulating monocytes. *Cellular immunology*, **194**(1), pp. 6-11.

SAPONARO, C., GAGGINI, M., CARLI, F. and GASTALDELLI, A., 2015. The Subtle Balance between Lipolysis and Lipogenesis: A Critical Point in Metabolic Homeostasis. *Nutrients*, **7**(11), pp. 9453-9474.

SARJEANT, K. and STEPHENS, J.M., 2012. Adipogenesis. *Cold Spring Harbor perspectives in biology*, **4**(9), pp. a008417.

SCHELLER, J., CHALARIS, A., SCHMIDT-ARRAS, D. and ROSE-JOHN, S., 2011. The pro- and anti-inflammatory properties of the cytokine interleukin-6. *Biochimica et biophysica acta*, **1813**(5), pp. 878-888.

SCHIRRA, J., KATSCHINSKI, M., WEIDMANN, C., SCHAFER, T., WANK, U., ARNOLD, R. and GOKE, B., 1996. Gastric emptying and release of incretin hormones after glucose ingestion in humans. *The Journal of clinical investigation*, **97**(1), pp. 92-103.

SCHWARTZ, D.R. and LAZAR, M.A., 2011. Human resistin: found in translation from mouse to man. *Trends in endocrinology and metabolism: TEM*, **22**(7), pp. 259-265.

SCHWARTZ, M.W., SEELEY, R.J., ZELTSER, L.M., DREWNOWSKI, A., RAVUSSIN, E., REDMAN, L.M. and LEIBEL, R.L., 2017. Obesity Pathogenesis: An Endocrine Society Scientific Statement. *Endocrine reviews*, **38**(4), pp. 267-296.

- SCHWARTZ, M.W., WOODS, S.C., PORTE, D., Jr, SEELEY, R.J. and BASKIN, D.G., 2000. Central nervous system control of food intake. *Nature*, **404**(6778), pp. 661-671.
- SEALE, P., BJORK, B., YANG, W., KAJIMURA, S., CHIN, S., KUANG, S., SCIME, A., DEVARAKONDA, S., CONROE, H.M., ERDJUMENT-BROMAGE, H., TEMPST, P., RUDNICKI, M.A., BEIER, D.R. and SPIEGELMAN, B.M., 2008. PRDM16 controls a brown fat/skeletal muscle switch. *Nature*, **454**(7207), pp. 961-967.
- SEALE, P., CONROE, H.M., ESTALL, J., KAJIMURA, S., FRONTINI, A., ISHIBASHI, J., COHEN, P., CINTI, S. and SPIEGELMAN, B.M., 2011. Prdm16 determines the thermogenic program of subcutaneous white adipose tissue in mice. *The Journal of clinical investigation*, **121**(1), pp. 96-105.
- SEALE, P., KAJIMURA, S., YANG, W., CHIN, S., ROHAS, L.M., ULDRY, M., TAVERNIER, G., LANGIN, D. and SPIEGELMAN, B.M., 2007. Transcriptional control of brown fat determination by PRDM16. *Cell metabolism*, **6**(1), pp. 38-54.
- SEAQUIST, E.R., DAMBERG, G.S., TKAC, I. and GRUETTER, R., 2001. The effect of insulin on in vivo cerebral glucose concentrations and rates of glucose transport/metabolism in humans. *Diabetes*, **50**(10), pp. 2203-2209.
- SECHER, A., JELSING, J., BAQUERO, A.F., HECKSHER-SORENSEN, J., COWLEY, M.A., DALBOGE, L.S., HANSEN, G., GROVE, K.L., PYKE, C., RAUN, K., SCHAFFER, L., TANG-CHRISTENSEN, M., VERMA, S., WITGEN, B.M., VRANG, N. and BJERRE KNUDSEN, L., 2014. The arcuate nucleus mediates GLP-1 receptor agonist liraglutide-dependent weight loss. *The Journal of clinical investigation*, **124**(10), pp. 4473-4488.
- SEGAL, K.R., LANDT, M. and KLEIN, S., 1996. Relationship between insulin sensitivity and plasma leptin concentration in lean and obese men. *Diabetes*, **45**(7), pp. 988-991.
- SEGAL, K.R., PRESTA, E. and GUTIN, B., 1984. Thermic effect of food during graded exercise in normal weight and obese men. *The American Journal of Clinical Nutrition*, **40**(5), pp. 995-1000.
- SELL, H., BLUHER, M., KLOTING, N., SCHLICH, R., WILLEMS, M., RUPPE, F., KNOEFEL, W.T., DIETRICH, A., FIELDING, B.A., ARNER, P., FRAYN, K.N. and ECKEL, J., 2013. Adipose dipeptidyl peptidase-4 and obesity: correlation with insulin resistance and depot-specific release from adipose tissue in vivo and in vitro. *Diabetes care*, **36**(12), pp. 4083-4090.
- SEMENZA, G.L., 2003. Targeting HIF-1 for cancer therapy. *Nature reviews.Cancer*, **3**(10), pp. 721-732.
- SENA, A., GRISHINA, I., THAI, A., GOULART, L., MACAL, M., FENTON, A., LI, J., PRINDIVILLE, T., OLIANI, S.M., DANDEKAR, S., GOULART, L. and

SANKARAN-WALTERS, S., 2013. Dysregulation of anti-inflammatory annexin A1 expression in progressive Crohns Disease. *PLoS one*, **8**(10), pp. e76969.

SENN, J.J., KLOVER, P.J., NOWAK, I.A., ZIMMERS, T.A., KONIARIS, L.G., FURLANETTO, R.W. and MOONEY, R.A., 2003. Suppressor of cytokine signaling-3 (SOCS-3), a potential mediator of interleukin-6-dependent insulin resistance in hepatocytes. *The Journal of biological chemistry*, **278**(16), pp. 13740-13746.

SHAO, X., WANG, M., WEI, X., DENG, S., FU, N., PENG, Q., JIANG, Y., YE, L., XIE, J. and LIN, Y., 2016. Peroxisome Proliferator-Activated Receptor-gamma: Master Regulator of Adipogenesis and Obesity. *Current stem cell research & therapy*, **11**(3), pp. 282-289.

SHAUL, M.E., BENNETT, G., STRISSEL, K.J., GREENBERG, A.S. and OBIN, M.S., 2010. Dynamic, M2-like remodeling phenotypes of CD11c⁺ adipose tissue macrophages during high-fat diet--induced obesity in mice. *Diabetes*, **59**(5), pp. 1171-1181.

SHAULIAN, E. and KARIN, M., 2002. AP-1 as a regulator of cell life and death. *Nature cell biology*, **4**(5), pp. E131-6.

SHEN, S. and WONG, C.H., 2016. Bugging inflammation: role of the gut microbiota. *Clinical & translational immunology*, **5**(4), pp. e72.

SHENHAR-TSARFATY, S., SHERF-DAGAN, S., BERMAN, G., WEBB, M., RAZIEL, A., KEIDAR, A., GOITEIN, D., SAKRAN, N., ZWANG, E., SHAPIRA, I., ZELTSER, D., BERLINER, S., ROGOWSKI, O., SHIBOLET, O. and ZELBER-SAGI, S., 2018. Obesity-related acetylcholinesterase elevation is reversed following laparoscopic sleeve gastrectomy. *International journal of obesity (2005)*, .

SHI, H., HALVORSEN, Y.D., ELLIS, P.N., WILKISON, W.O. and ZEMEL, M.B., 2000. Role of intracellular calcium in human adipocyte differentiation. *Physiological genomics*, **3**(2), pp. 75-82.

SHIMOMURA, I., HAMMER, R.E., IKEMOTO, S., BROWN, M.S. and GOLDSTEIN, J.L., 1999. Leptin reverses insulin resistance and diabetes mellitus in mice with congenital lipodystrophy. *Nature*, **401**(6748), pp. 73-76.

SHIMOMURA, I., SHIMANO, H., HORTON, J.D., GOLDSTEIN, J.L. and BROWN, M.S., 1997. Differential expression of exons 1a and 1c in mRNAs for sterol regulatory element binding protein-1 in human and mouse organs and cultured cells. *The Journal of clinical investigation*, **99**(5), pp. 838-845.

SHIN, M.K., JANG, Y.H., YOO, H.J., KANG, D.W., PARK, M.H., KIM, M.K., SONG, J.H., KIM, S.D., MIN, G., YOU, H.K., CHOI, K.Y., BAE, Y.S. and MIN DO, S., 2011. N-formyl-methionyl-leucyl-phenylalanine (fMLP) promotes osteoblast differentiation via the N-formyl peptide receptor 1-mediated signaling pathway in human mesenchymal stem cells from bone marrow. *The Journal of biological chemistry*, **286**(19), pp. 17133-17143.

- SHIOMI, Y., YAMAUCHI, T., IWABU, M., OKADA-IWABU, M., NAKAYAMA, R., ORIKAWA, Y., YOSHIOKA, Y., TANAKA, K., UEKI, K. and KADOWAKI, T., 2015. A Novel Peroxisome Proliferator-activated Receptor (PPAR)alpha Agonist and PPARgamma Antagonist, Z-551, Ameliorates High-fat Diet-induced Obesity and Metabolic Disorders in Mice. *The Journal of biological chemistry*, **290**(23), pp. 14567-14581.
- SHOELSON, S.E., LEE, J. and GOLDFINE, A.B., 2006. Inflammation and insulin resistance. *The Journal of clinical investigation*, **116**(7), pp. 1793-1801.
- SICA, A., WANG, J.M., COLOTTA, F., DEJANA, E., MANTOVANI, A., OPPENHEIM, J.J., LARSEN, C.G., ZACHARIAE, C.O. and MATSUSHIMA, K., 1990. Monocyte chemotactic and activating factor gene expression induced in endothelial cells by IL-1 and tumor necrosis factor. *Journal of immunology (Baltimore, Md.: 1950)*, **144**(8), pp. 3034-3038.
- SIERRA-HONIGMANN, M.R., NATH, A.K., MURAKAMI, C., GARCIA-CARDENA, G., PAPAPETROPOULOS, A., SESSA, W.C., MADGE, L.A., SCHECHNER, J.S., SCHWABB, M.B., POLVERINI, P.J. and FLORES-RIVEROS, J.R., 1998. Biological action of leptin as an angiogenic factor. *Science (New York, N.Y.)*, **281**(5383), pp. 1683-1686.
- SILLAKIVI, T., SUUMANN, J., KIRSIMAGI, U. and PEETSALU, A., 2013. Plasma levels of gastric biomarkers in patients after bariatric surgery: biomarkers after bariatric surgery. *Hepato-gastroenterology*, **60**(128), pp. 2129-2132.
- SILVA, J.E., 2003. The thermogenic effect of thyroid hormone and its clinical implications. *Annals of Internal Medicine*, **139**(3), pp. 205-213.
- SIMONSON, D.C., TAPPY, L., JEQUIER, E., FELBER, J.P. and DEFRONZO, R.A., 1988. Normalization of carbohydrate-induced thermogenesis by fructose in insulin-resistant states. *The American Journal of Physiology*, **254**(2 Pt 1), pp. E201-7.
- SIN, D.D. and SUTHERLAND, E.R., 2008. Obesity and the lung: 4. Obesity and asthma. *Thorax*, **63**(11), pp. 1018-1023.
- SINGER, K., MORRIS, D.L., OATMEN, K.E., WANG, T., DELPROPOSTO, J., MERGIAN, T., CHO, K.W. and LUMENG, C.N., 2013. Neuropeptide Y is produced by adipose tissue macrophages and regulates obesity-induced inflammation. *PloS one*, **8**(3), pp. e57929.
- SIRIWARDHANA, N., KALUPAHANA, N.S., CEKANOVA, M., LEMIEUX, M., GREER, B. and MOUSTAID-MOUSSA, N., 2013. Modulation of adipose tissue inflammation by bioactive food compounds. *The Journal of nutritional biochemistry*, **24**(4), pp. 613-623.

- SISSONS, J.G., WEST, R.J., FALLOWS, J., WILLIAMS, D.G., BOUCHER, B.J., AMOS, N. and PETERS, D.K., 1976. The complement abnormalities of lipodystrophy. *The New England journal of medicine*, **294**(9), pp. 461-465.
- SMAS, C.M., CHEN, L. and SUL, H.S., 1997. Cleavage of membrane-associated pref-1 generates a soluble inhibitor of adipocyte differentiation. *Molecular and cellular biology*, **17**(2), pp. 977-988.
- SMAS, C.M., KACHINSKAS, D., LIU, C.M., XIE, X., DIRCKS, L.K. and SUL, H.S., 1998. Transcriptional control of the pref-1 gene in 3T3-L1 adipocyte differentiation. Sequence requirement for differentiation-dependent suppression. *The Journal of biological chemistry*, **273**(48), pp. 31751-31758.
- SMAS, C.M. and SUL, H.S., 1993. Pref-1, a protein containing EGF-like repeats, inhibits adipocyte differentiation. *Cell*, **73**(4), pp. 725-734.
- SOLITO, E., KAMAL, A., RUSSO-MARIE, F., BUCKINGHAM, J.C., MARULLO, S. and PERRETTI, M., 2003. A novel calcium-dependent proapoptotic effect of annexin 1 on human neutrophils. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **17**(11), pp. 1544-1546.
- SONG, N.J., KIM, S., JANG, B.H., CHANG, S.H., YUN, U.J., PARK, K.M., WAKI, H., LI, D.Y., TONTONOZ, P. and PARK, K.W., 2016. Small Molecule-Induced Complement Factor D (Adipsin) Promotes Lipid Accumulation and Adipocyte Differentiation. *PloS one*, **11**(9), pp. e0162228.
- SOTORNIK, R., BRASSARD, P., MARTIN, E., YALE, P., CARPENTIER, A.C. and ARDILOUZE, J.L., 2012. Update on adipose tissue blood flow regulation. *American journal of physiology. Endocrinology and metabolism*, **302**(10), pp. E1157-70.
- SOUKAS, A., SOCCI, N.D., SAATKAMP, B.D., NOVELLI, S. and FRIEDMAN, J.M., 2001. Distinct transcriptional profiles of adipogenesis in vivo and in vitro. *The Journal of biological chemistry*, **276**(36), pp. 34167-34174.
- SPADARO, O., CAMELL, C.D., BOSURGI, L., NGUYEN, K.Y., YOUM, Y.H., ROTHLIN, C.V. and DIXIT, V.D., 2017. IGF1 Shapes Macrophage Activation in Response to Immunometabolic Challenge. *Cell reports*, **19**(2), pp. 225-234.
- SPADARO, O., GOLDBERG, E.L., CAMELL, C.D., YOUM, Y.H., KOPCHICK, J.J., NGUYEN, K.Y., BARTKE, A., SUN, L.Y. and DIXIT, V.D., 2016. Growth Hormone Receptor Deficiency Protects against Age-Related NLRP3 Inflammasome Activation and Immune Senescence. *Cell reports*, **14**(7), pp. 1571-1580.
- SPALDING, K.L., ARNER, E., WESTERMARK, P.O., BERNARD, S., BUCHHOLZ, B.A., BERGMANN, O., BLOMQVIST, L., HOFFSTEDT, J., NASLUND, E., BRITTON, T., CONCHA, H., HASSAN, M., RYDEN, M., FRISEN, J. and ARNER, P., 2008. Dynamics of fat cell turnover in humans. *Nature*, **453**(7196), pp. 783-787.

SPIEGELMAN, B.M. and FLIER, J.S., 2001. Obesity and the regulation of energy balance. *Cell*, **104**(4), pp. 531-543.

SPITE, M., CLARIA, J. and SERHAN, C.N., 2014. Resolvins, specialized proresolving lipid mediators, and their potential roles in metabolic diseases. *Cell metabolism*, **19**(1), pp. 21-36.

SPRECKLEY, E. and MURPHY, K.G., 2015. The L-Cell in Nutritional Sensing and the Regulation of Appetite. *Frontiers in nutrition*, **2**, pp. 23.

STANDEVEN, K.F., HESS, K., CARTER, A.M., RICE, G.I., CORDELL, P.A., BALMFORTH, A.J., LU, B., SCOTT, D.J., TURNER, A.J., HOOPER, N.M. and GRANT, P.J., 2011. Neprilysin, obesity and the metabolic syndrome. *International journal of obesity (2005)*, **35**(8), pp. 1031-1040.

STANLEY, T.L., ZANNI, M.V., JOHNSEN, S., RASHEED, S., MAKIMURA, H., LEE, H., KHOR, V.K., AHIMA, R.S. and GRINSPOON, S.K., 2011. TNF-alpha antagonism with etanercept decreases glucose and increases the proportion of high molecular weight adiponectin in obese subjects with features of the metabolic syndrome. *The Journal of clinical endocrinology and metabolism*, **96**(1), pp. E146-50.

STARKIE, R., OSTROWSKI, S.R., JAUFFRED, S., FEBBRAIO, M. and PEDERSEN, B.K., 2003. Exercise and IL-6 infusion inhibit endotoxin-induced TNF-alpha production in humans. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **17**(8), pp. 884-886.

STEFANOWICZ, M., NIKOLAJUK, A., MATULEWICZ, N. and KARCZEWSKA-KUPCZEWSKA, M., 2018. Adipose tissue, but not skeletal muscle, sirtuin 1 expression is decreased in obesity and related to insulin sensitivity. *Endocrine*, **60**(2), pp. 263-271.

STEPHENS, J.M., MORRISON, R.F. and PILCH, P.F., 1996. The expression and regulation of STATs during 3T3-L1 adipocyte differentiation. *The Journal of biological chemistry*, **271**(18), pp. 10441-10444.

STEPPAN, C.M., BAILEY, S.T., BHAT, S., BROWN, E.J., BANERJEE, R.R., WRIGHT, C.M., PATEL, H.R., AHIMA, R.S. and LAZAR, M.A., 2001. The hormone resistin links obesity to diabetes. *Nature*, **409**(6818), pp. 307-312.

STEPPAN, C.M., WANG, J., WHITEMAN, E.L., BIRNBAUM, M.J. and LAZAR, M.A., 2005. Activation of SOCS-3 by resistin. *Molecular and cellular biology*, **25**(4), pp. 1569-1575.

STEVENS, G.A., SINGH, G.M., LU, Y., DANAEI, G., LIN, J.K., FINUCANE, M.M., BAHALIM, A.N., MCINTIRE, R.K., GUTIERREZ, H.R., COWAN, M., PACIOREK, C.J., FARZADFAR, F., RILEY, L., EZZATI, M. and GLOBAL BURDEN OF METABOLIC RISK FACTORS OF CHRONIC DISEASES COLLABORATING GROUP (BODY MASS INDEX), 2012. National, regional, and global trends in adult

overweight and obesity prevalences. *Population health metrics*, **10**(1), pp. 22-7954-10-22.

STRABLE, M.S. and NTAMBI, J.M., 2010. Genetic control of de novo lipogenesis: role in diet-induced obesity. *Critical reviews in biochemistry and molecular biology*, **45**(3), pp. 199-214.

STRADER, A.D., VAHL, T.P., JANDACEK, R.J., WOODS, S.C., D'ALESSIO, D.A. and SEELEY, R.J., 2005. Weight loss through ileal transposition is accompanied by increased ileal hormone secretion and synthesis in rats. *American journal of physiology. Endocrinology and metabolism*, **288**(2), pp. E447-53.

STRAWFORD, A., ANTELO, F., CHRISTIANSEN, M. and HELLERSTEIN, M.K., 2004. Adipose tissue triglyceride turnover, de novo lipogenesis, and cell proliferation in humans measured with 2H₂O. *American journal of physiology. Endocrinology and metabolism*, **286**(4), pp. E577-88.

STRISSEL, K.J., STANCHEVA, Z., MIYOSHI, H., PERFIELD, J.W.,^{2nd}, DEFURIA, J., JICK, Z., GREENBERG, A.S. and OBIN, M.S., 2007. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes*, **56**(12), pp. 2910-2918.

SUBRAMANIAN, M., OZCAN, L., GHORPADE, D.S., FERRANTE, A.W., Jr and TABAS, I., 2015. Suppression of Adaptive Immune Cell Activation Does Not Alter Innate Immune Adipose Inflammation or Insulin Resistance in Obesity. *PloS one*, **10**(8), pp. e0135842.

SUBRAMANYAM, L., SIMHA, V. and GARG, A., 2010. Overlapping syndrome with familial partial lipodystrophy, Dunnigan variety and cardiomyopathy due to amino-terminal heterozygous missense lamin A/C mutations. *Clinical genetics*, **78**(1), pp. 66-73.

SUGIMOTO, M.A., SOUSA, L.P., PINHO, V., PERRETTI, M. and TEIXEIRA, M.M., 2016. Resolution of Inflammation: What Controls Its Onset? *Frontiers in immunology*, **7**, pp. 160.

SUGIMOTO, M.A., VAGO, J.P., TEIXEIRA, M.M. and SOUSA, L.P., 2016. Annexin A1 and the Resolution of Inflammation: Modulation of Neutrophil Recruitment, Apoptosis, and Clearance. *Journal of immunology research*, **2016**, pp. 8239258.

SULLIVAN, M., KARLSSON, J., SJOSTROM, L., BACKMAN, L., BENGTSSON, C., BOUCHARD, C., DAHLGREN, S., JONSSON, E., LARSSON, B. and LINDSTEDT, S., 1993. Swedish obese subjects (SOS)--an intervention study of obesity. Baseline evaluation of health and psychosocial functioning in the first 1743 subjects examined. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*, **17**(9), pp. 503-512.

SUN, G., BISHOP, J., KHALILI, S., VASDEV, S., GILL, V., PACE, D., FITZPATRICK, D., RANDELL, E., XIE, Y.G. and ZHANG, H., 2007. Serum visfatin

concentrations are positively correlated with serum triacylglycerols and down-regulated by overfeeding in healthy young men. *The American Journal of Clinical Nutrition*, **85**(2), pp. 399-404.

SUN, K., WERNSTEDT ASTERHOLM, I., KUSMINSKI, C.M., BUENO, A.C., WANG, Z.V., POLLARD, J.W., BREKKEN, R.A. and SCHERER, P.E., 2012. Dichotomous effects of VEGF-A on adipose tissue dysfunction. *Proceedings of the National Academy of Sciences of the United States of America*, **109**(15), pp. 5874-5879.

SUNDBOM, M., HOLDSTOCK, C., ENGSTROM, B.E. and KARLSSON, F.A., 2007. Early changes in ghrelin following Roux-en-Y gastric bypass: influence of vagal nerve functionality? *Obesity Surgery*, **17**(3), pp. 304-310.

SUZUKI, K., SIMPSON, K.A., MINNION, J.S., SHILLITO, J.C. and BLOOM, S.R., 2010. The role of gut hormones and the hypothalamus in appetite regulation. *Endocrine journal*, **57**(5), pp. 359-372.

SVENSSON, M.K., JANSSON, P.A., PERSSON, A.L., SJOSTRAND, M. and ERIKSSON, J.W., 2011. Atropine improves insulin sensitivity in both lean and abdominally obese subjects. *The Journal of clinical endocrinology and metabolism*, **96**(11), pp. E1843-7.

SZENDROEDI, J. and RODEN, M., 2008. Mitochondrial fitness and insulin sensitivity in humans. *Diabetologia*, **51**(12), pp. 2155-2167.

TAKASHIMA, Y., ERA, T., NAKAO, K., KONDO, S., KASUGA, M., SMITH, A.G. and NISHIKAWA, S., 2007. Neuroepithelial cells supply an initial transient wave of MSC differentiation. *Cell*, **129**(7), pp. 1377-1388.

TAKEMURA, Y., OUCHI, N., SHIBATA, R., APRAHAMIAN, T., KIRBER, M.T., SUMMER, R.S., KIHARA, S. and WALSH, K., 2007. Adiponectin modulates inflammatory reactions via calreticulin receptor-dependent clearance of early apoptotic bodies. *The Journal of clinical investigation*, **117**(2), pp. 375-386.

TAKENOUCHI, T., TAKAYAMA, Y. and TAKEZAWA, T., 2004. Co-treatment with dexamethasone and octanoate induces adipogenesis in 3T3-L1 cells. *Cell biology international*, **28**(3), pp. 209-216.

TALUKDAR, S., OH DA, Y., BANDYOPADHYAY, G., LI, D., XU, J., MCNELIS, J., LU, M., LI, P., YAN, Q., ZHU, Y., OFRECIO, J., LIN, M., BRENNER, M.B. and OLEFSKY, J.M., 2012. Neutrophils mediate insulin resistance in mice fed a high-fat diet through secreted elastase. *Nature medicine*, **18**(9), pp. 1407-1412.

TAN, T.M., SALEM, V., TROKE, R.C., ALSAFI, A., FIELD, B.C., DE SILVA, A., MISRA, S., BAYNES, K.C., DONALDSON, M., MINNION, J., GHATEI, M.A., GODSLAND, I.F. and BLOOM, S.R., 2014. Combination of peptide YY3-36 with GLP-1(7-36) amide causes an increase in first-phase insulin secretion after IV glucose. *The Journal of clinical endocrinology and metabolism*, **99**(11), pp. E2317-24.

- TANAKA, T., YOSHIDA, N., KISHIMOTO, T. and AKIRA, S., 1997. Defective adipocyte differentiation in mice lacking the C/EBPbeta and/or C/EBPdelta gene. *The EMBO journal*, **16**(24), pp. 7432-7443.
- TANG, W., ZEVE, D., SUH, J.M., BOSNAKOVSKI, D., KYBA, M., HAMMER, R.E., TALLQUIST, M.D. and GRAFF, J.M., 2008. White fat progenitor cells reside in the adipose vasculature. *Science (New York, N.Y.)*, **322**(5901), pp. 583-586.
- TAO, Y.X., 2005. Molecular mechanisms of the neural melanocortin receptor dysfunction in severe early onset obesity. *Molecular and cellular endocrinology*, **239**(1-2), pp. 1-14.
- TAPPY, L., 1996. Thermic effect of food and sympathetic nervous system activity in humans. *Reproduction, nutrition, development*, **36**(4), pp. 391-397.
- TAPPY, L. and JEQUIER, E., 1993. Fructose and dietary thermogenesis. *The American Journal of Clinical Nutrition*, **58**(5 Suppl), pp. 766S-770S.
- TCHERNOF, A., NOLAN, A., SITES, C.K., ADES, P.A. and POEHLMAN, E.T., 2002. Weight loss reduces C-reactive protein levels in obese postmenopausal women. *Circulation*, **105**(5), pp. 564-569.
- THIRONE, A.C., HUANG, C. and KLIP, A., 2006. Tissue-specific roles of IRS proteins in insulin signaling and glucose transport. *Trends in endocrinology and metabolism: TEM*, **17**(2), pp. 72-78.
- TILG, H. and MOSCHEN, A.R., 2006. Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nature reviews.Immunology*, **6**(10), pp. 772-783.
- TOMAS, E., TSAO, T.S., SAHA, A.K., MURREY, H.E., ZHANG CC, C., ITANI, S.I., LODISH, H.F. and RUDERMAN, N.B., 2002. Enhanced muscle fat oxidation and glucose transport by ACRP30 globular domain: acetyl-CoA carboxylase inhibition and AMP-activated protein kinase activation. *Proceedings of the National Academy of Sciences of the United States of America*, **99**(25), pp. 16309-16313.
- TONG, Q., DALGIN, G., XU, H., TING, C.N., LEIDEN, J.M. and HOTAMISLIGIL, G.S., 2000. Function of GATA transcription factors in preadipocyte-adipocyte transition. *Science (New York, N.Y.)*, **290**(5489), pp. 134-138.
- TONG, Q., TSAI, J., TAN, G., DALGIN, G. and HOTAMISLIGIL, G.S., 2005. Interaction between GATA and the C/EBP family of transcription factors is critical in GATA-mediated suppression of adipocyte differentiation. *Molecular and cellular biology*, **25**(2), pp. 706-715.
- TONTONOZ, P., HU, E. and SPIEGELMAN, B.M., 1994. Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor. *Cell*, **79**(7), pp. 1147-1156.

TONTONOZ, P., KIM, J.B., GRAVES, R.A. and SPIEGELMAN, B.M., 1993. ADD1: a novel helix-loop-helix transcription factor associated with adipocyte determination and differentiation. *Molecular and cellular biology*, **13**(8), pp. 4753-4759.

TOWBIN, H., STAEBELIN, T. and GORDON, J., 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences of the United States of America*, **76**(9), pp. 4350-4354.

TRAN, K.V., GEALEKMAN, O., FRONTINI, A., ZINGARETTI, M.C., MORRONI, M., GIORDANO, A., SMORLESI, A., PERUGINI, J., DE MATTEIS, R., SBARBATI, A., CORVERA, S. and CINTI, S., 2012. The vascular endothelium of the adipose tissue gives rise to both white and brown fat cells. *Cell metabolism*, **15**(2), pp. 222-229.

TRAYHURN, P., 2013. Hypoxia and adipose tissue function and dysfunction in obesity. *Physiological Reviews*, **93**(1), pp. 1-21.

TRAYHURN, P. and BEATTIE, J.H., 2001. Physiological role of adipose tissue: white adipose tissue as an endocrine and secretory organ. *The Proceedings of the Nutrition Society*, **60**(3), pp. 329-339.

TRAYHURN, P. and WOOD, I.S., 2004. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *The British journal of nutrition*, **92**(3), pp. 347-355.

TREMME, M., GERDTHAM, U.G., NILSSON, P.M. and SAHA, S., 2017. Economic Burden of Obesity: A Systematic Literature Review. *International journal of environmental research and public health*, **14**(4), pp. 435. doi: 10.3390/ijerph14040435. Print 2017 Apr.

TRUJILLO, M.E. and SCHERER, P.E., 2006. Adipose tissue-derived factors: impact on health and disease. *Endocrine reviews*, **27**(7), pp. 762-778.

TRUJILLO, M.E. and SCHERER, P.E., 2005. Adiponectin--journey from an adipocyte secretory protein to biomarker of the metabolic syndrome. *Journal of internal medicine*, **257**(2), pp. 167-175.

TSIGOS, C., PAPANICOLAOU, D.A., KYROU, I., DEFENSOR, R., MITSIADIS, C.S. and CHROUSOS, G.P., 1997. Dose-dependent effects of recombinant human interleukin-6 on glucose regulation. *The Journal of clinical endocrinology and metabolism*, **82**(12), pp. 4167-4170.

TSUCHIDA, A., YAMAUCHI, T., TAKEKAWA, S., HADA, Y., ITO, Y., MAKI, T. and KADOWAKI, T., 2005. Peroxisome proliferator-activated receptor (PPAR)alpha activation increases adiponectin receptors and reduces obesity-related inflammation in adipose tissue: comparison of activation of PPARalpha, PPARgamma, and their combination. *Diabetes*, **54**(12), pp. 3358-3370.

- TZANAVARI, T., GIANNOGONAS, P. and KARALIS, K.P., 2010. TNF-alpha and obesity. *Current directions in autoimmunity*, **11**, pp. 145-156.
- UNGER, R.H., CLARK, G.O., SCHERER, P.E. and ORCI, L., 2010. Lipid homeostasis, lipotoxicity and the metabolic syndrome. *Biochimica et biophysica acta*, **1801**(3), pp. 209-214.
- UYSAL, K.T., WIESBROCK, S.M., MARINO, M.W. and HOTAMISLIGIL, G.S., 1997. Protection from obesity-induced insulin resistance in mice lacking TNF-alpha function. *Nature*, **389**(6651), pp. 610-614.
- VAGO, J.P., NOGUEIRA, C.R., TAVARES, L.P., SORIANI, F.M., LOPES, F., RUSSO, R.C., PINHO, V., TEIXEIRA, M.M. and SOUSA, L.P., 2012. Annexin A1 modulates natural and glucocorticoid-induced resolution of inflammation by enhancing neutrophil apoptosis. *Journal of leukocyte biology*, **92**(2), pp. 249-258.
- VAHL, T.P., DRAZEN, D.L., SEELEY, R.J., D'ALESSIO, D.A. and WOODS, S.C., 2010. Meal-anticipatory glucagon-like peptide-1 secretion in rats. *Endocrinology*, **151**(2), pp. 569-575.
- VAN DEN BERG, S.M., VAN DAM, A.D., RENSEN, P.C., DE WINTHER, M.P. and LUTGENS, E., 2017. Immune Modulation of Brown(ing) Adipose Tissue in Obesity. *Endocrine reviews*, **38**(1), pp. 46-68.
- VAN DEN HOEK, A.M., TEUSINK, B., VOSHOL, P.J., HAVEKES, L.M., ROMIJN, J.A. and PIJL, H., 2008. Leptin deficiency per se dictates body composition and insulin action in ob/ob mice. *Journal of neuroendocrinology*, **20**(1), pp. 120-127.
- VAN GAAL, L.F., MERTENS, I.L. and DE BLOCK, C.E., 2006. Mechanisms linking obesity with cardiovascular disease. *Nature*, **444**(7121), pp. 875-880.
- VAN HARMELEN, V., REYNISDOTTIR, S., CIANFLONE, K., DEGERMAN, E., HOFFSTEDT, J., NILSELL, K., SNIDERMAN, A. and ARNER, P., 1999. Mechanisms involved in the regulation of free fatty acid release from isolated human fat cells by acylation-stimulating protein and insulin. *The Journal of biological chemistry*, **274**(26), pp. 18243-18251.
- VAN MALDERGEM, L., MAGRE, J., KHALLOUF, T.E., GEDDE-DAHL, T., Jr, DELEPINE, M., TRYGSTAD, O., SEEMANOVA, E., STEPHENSON, T., ALBOTT, C.S., BONNICI, F., PANZ, V.R., MEDINA, J.L., BOGALHO, P., HUET, F., SAVASTA, S., VERLOES, A., ROBERT, J.J., LORET, H., DE Kerdanet, M., TUBIANA-RUFI, N., MEGARBANE, A., MAASSEN, J., POLAK, M., LACOMBE, D., KAHN, C.R., SILVEIRA, E.L., D'ABRONZO, F.H., GRIGORESCU, F., LATHROP, M., CAPEAU, J. and O'RAHILLY, S., 2002. Genotype-phenotype relationships in Berardinelli-Seip congenital lipodystrophy. *Journal of medical genetics*, **39**(10), pp. 722-733.

- VANTYGHM, M.C., VINCENT-DESPLANQUES, D., DEFRANCE-FAIVRE, F., CAPEAU, J., FERMON, C., VALAT, A.S., LASCOLS, O., HECART, A.C., PIGNY, P., DELEMER, B., VIGOUROUX, C. and WEMEAU, J.L., 2008. Fertility and obstetrical complications in women with LMNA-related familial partial lipodystrophy. *The Journal of clinical endocrinology and metabolism*, **93**(6), pp. 2223-2229.
- VELIJA-ASIMI, Z., IZETBEGOVIC, S., KARAMEHIC, J., CORIC, J., PANJETA, M., MACIC-DZANKOVIC, A., DJELILOVIC-VRANIC, J. and DIZDAREVIC-BOSTANDZIC, A., 2013. The effects of dipeptidyl peptidase-4 inhibitors in treatment of obese patients with type 2 diabetes. *Medical archives (Sarajevo, Bosnia and Herzegovina)*, **67**(5), pp. 365-367.
- VERDICH, C., FLINT, A., GUTZWILLER, J.P., NASLUND, E., BEGLINGER, C., HELLSTROM, P.M., LONG, S.J., MORGAN, L.M., HOLST, J.J. and ASTRUP, A., 2001. A meta-analysis of the effect of glucagon-like peptide-1 (7-36) amide on ad libitum energy intake in humans. *The Journal of clinical endocrinology and metabolism*, **86**(9), pp. 4382-4389.
- VERWAERDE, C., DELANOYE, A., MACIA, L., TAILLEUX, A. and WOLOWCZUK, I., 2006. Influence of high-fat feeding on both I and antigen-experienced T-cell immune response in DO10.11 mice. *Scandinavian Journal of Immunology*, **64**(5), pp. 457-466.
- VIANA, E.C., ARAUJO-DASILIO, K.L., MIGUEL, G.P., BRESSAN, J., LEMOS, E.M., MOYSES, M.R., DE ABREU, G.R., DE AZEVEDO, J.L., CARVALHO, P.S., PASSOS-BUENO, M.R., ERRERA, F.I. and BISSOLI, N.S., 2013. Gastric bypass and sleeve gastrectomy: the same impact on IL-6 and TNF-alpha. Prospective clinical trial. *Obesity Surgery*, **23**(8), pp. 1252-1261.
- VILLARROYA, F., DOMINGO, P. and GIRALT, M., 2010. Drug-induced lipotoxicity: lipodystrophy associated with HIV-1 infection and antiretroviral treatment. *Biochimica et biophysica acta*, **1801**(3), pp. 392-399.
- VILSBOLL, T. and HOLST, J.J., 2004. Incretins, insulin secretion and Type 2 diabetes mellitus. *Diabetologia*, **47**(3), pp. 357-366.
- VINSON, C.R., SIGLER, P.B. and MCKNIGHT, S.L., 1989. Scissors-grip model for DNA recognition by a family of leucine zipper proteins. *Science (New York, N.Y.)*, **246**(4932), pp. 911-916.
- VIRTANEN, K.A., LIDELL, M.E., ORAVA, J., HEGLIND, M., WESTERGREN, R., NIEMI, T., TAITTONEN, M., LAINE, J., SAVISTO, N.J., ENERBACK, S. and NUUTILA, P., 2009. Functional brown adipose tissue in healthy adults. *The New England journal of medicine*, **360**(15), pp. 1518-1525.
- VIRTUE, S. and VIDAL-PUIG, A., 2010. Adipose tissue expandability, lipotoxicity and the Metabolic Syndrome--an allostatic perspective. *Biochimica et biophysica acta*, **1801**(3), pp. 338-349.

VISSER, M., BOUTER, L.M., MCQUILLAN, G.M., WENER, M.H. and HARRIS, T.B., 1999. Elevated C-reactive protein levels in overweight and obese adults. *Jama*, **282**(22), pp. 2131-2135.

VOZAROVA, B., WEYER, C., HANSON, K., TATARANNI, P.A., BOGARDUS, C. and PRATLEY, R.E., 2001. Circulating interleukin-6 in relation to adiposity, insulin action, and insulin secretion. *Obesity research*, **9**(7), pp. 414-417.

VROEGRIJK, I.O., VAN KLINKEN, J.B., VAN DIEPEN, J.A., VAN DEN BERG, S.A., FEBBRAIO, M., STEINBUSCH, L.K., GLATZ, J.F., HAVEKES, L.M., VOSHOL, P.J., RENSEN, P.C., VAN DIJK, K.W. and VAN HARMELEN, V., 2013. CD36 is important for adipocyte recruitment and affects lipolysis. *Obesity (Silver Spring, Md.)*, **21**(10), pp. 2037-2045.

WAKI, H. and TONTONOZ, P., 2007. Endocrine functions of adipose tissue. *Annual review of pathology*, **2**, pp. 31-56.

WALDEN, T.B., HANSEN, I.R., TIMMONS, J.A., CANNON, B. and NEDERGAARD, J., 2012. Recruited vs. nonrecruited molecular signatures of brown, "brite," and white adipose tissues. *American journal of physiology. Endocrinology and metabolism*, **302**(1), pp. E19-31.

WALLACE, A.M., MCMAHON, A.D., PACKARD, C.J., KELLY, A., SHEPHERD, J., GAW, A. and SATTAR, N., 2001. Plasma leptin and the risk of cardiovascular disease in the west of Scotland coronary prevention study (WOSCOPS). *Circulation*, **104**(25), pp. 3052-3056.

WALLENIUS, V., WALLENIUS, K., AHREN, B., RUDLING, M., CARLSTEN, H., DICKSON, S.L., OHLSSON, C. and JANSSEN, J.O., 2002. Interleukin-6-deficient mice develop mature-onset obesity. *Nature medicine*, **8**(1), pp. 75-79.

WANG, B., WOOD, I.S. and TRAYHURN, P., 2007. Dysregulation of the expression and secretion of inflammation-related adipokines by hypoxia in human adipocytes. *Pflugers Archiv : European journal of physiology*, **455**(3), pp. 479-492.

WANG, D., WANG, Q., YAN, G., QIAO, Y., ZHU, B., LIU, B. and TANG, C., 2018. Hypoxia induces lactate secretion and glycolytic efflux by downregulating mitochondrial pyruvate carrier levels in human umbilical vein endothelial cells. *Molecular medicine reports*, **18**(2), pp. 1710-1717.

WANG, D., ZHOU, Y., LEI, W., ZHANG, K., SHI, J., HU, Y., SHU, G. and SONG, J., 2009. Signal transducer and activator of transcription 3 (STAT3) regulates adipocyte differentiation via peroxisome-proliferator-activated receptor gamma (PPARgamma). *Biology of the cell*, **102**(1), pp. 1-12.

WANG, G.L., SHI, X., SALISBURY, E., SUN, Y., ALBRECHT, J.H., SMITH, R.G. and TIMCHENKO, N.A., 2006. Cyclin D3 maintains growth-inhibitory activity of

C/EBPalpha by stabilizing C/EBPalpha-cdk2 and C/EBPalpha-Brm complexes. *Molecular and cellular biology*, **26**(7), pp. 2570-2582.

WANG, H., XIE, Z. and SCOTT, R.E., 1996. Differentiation modulates the balance of positive and negative Jun/AP-1 DNA binding activities to regulate cellular proliferative potential: different effects in nontransformed and transformed cells. *The Journal of cell biology*, **135**(4), pp. 1151-1162.

WANG, N.D., FINEGOLD, M.J., BRADLEY, A., OU, C.N., ABDELSAYED, S.V., WILDE, M.D., TAYLOR, L.R., WILSON, D.R. and DARLINGTON, G.J., 1995. Impaired energy homeostasis in C/EBP alpha knockout mice. *Science (New York, N.Y.)*, **269**(5227), pp. 1108-1112.

WANG, Q.A., TAO, C., GUPTA, R.K. and SCHERER, P.E., 2013. Tracking adipogenesis during white adipose tissue development, expansion and regeneration. *Nature medicine*, **19**(10), pp. 1338-1344.

WANG, Y., CHENG, Y.S., YIN, X.Q., YU, G. and JIA, B.L., 2018. Anxa2 gene silencing attenuates obesity-induced insulin resistance by suppressing NF-small ka, CyrillicB signaling pathway. *American journal of physiology. Cell physiology*, .

WANG, Y., MAK, J.C.W., LEE, M.Y.K., XU, A. and IP, M.S.M., 2018. Low-Frequency Intermittent Hypoxia Promotes Subcutaneous Adipogenic Differentiation. *Oxidative medicine and cellular longevity*, **2018**, pp. 4501757.

WANG, Y., NISHINA, P.M. and NAGGERT, J.K., 2009. Degradation of IRS1 leads to impaired glucose uptake in adipose tissue of the type 2 diabetes mouse model TALLYHO/Jng. *The Journal of endocrinology*, **203**(1), pp. 65-74.

WANG, Z. and NAKAYAMA, T., 2010. Inflammation, a link between obesity and cardiovascular disease. *Mediators of inflammation*, **2010**, pp. 535918.

WANG, Z.V. and SCHERER, P.E., 2016. Adiponectin, the past two decades. *Journal of molecular cell biology*, **8**(2), pp. 93-100.

WARNE, J.P., JOHN, C.D., CHRISTIAN, H.C., MORRIS, J.F., FLOWER, R.J., SUGDEN, D., SOLITO, E., GILLIES, G.E. and BUCKINGHAM, J.C., 2006. Gene deletion reveals roles for annexin A1 in the regulation of lipolysis and IL-6 release in epididymal adipose tissue. *American journal of physiology. Endocrinology and metabolism*, **291**(6), pp. E1264-73.

WEIGERT, N., SCHAFFER, K., SCHUSDZIARRA, V., CLASSEN, M. and SCHEPP, W., 1996. Gastrin secretion from primary cultures of rabbit antral G cells: stimulation by inflammatory cytokines. *Gastroenterology*, **110**(1), pp. 147-154.

WEISBERG, S.P., MCCANN, D., DESAI, M., ROSENBAUM, M., LEIBEL, R.L. and FERRANTE, A.W., Jr, 2003. Obesity is associated with macrophage accumulation in adipose tissue. *The Journal of clinical investigation*, **112**(12), pp. 1796-1808.

- WEISS, M.J. and ORKIN, S.H., 1995. GATA transcription factors: key regulators of hematopoiesis. *Experimental hematology*, **23**(2), pp. 99-107.
- WESTERTEP, K.R., 2004. Diet induced thermogenesis. *Nutrition & metabolism*, **1**(1), pp. 5.
- WHITE, M.F., 2002. IRS proteins and the common path to diabetes. *American journal of physiology. Endocrinology and metabolism*, **283**(3), pp. E413-22.
- WIECKOWSKA, A., PAPOUCHADO, B.G., LI, Z., LOPEZ, R., ZEIN, N.N. and FELDSTEIN, A.E., 2008. Increased hepatic and circulating interleukin-6 levels in human nonalcoholic steatohepatitis. *The American Journal of Gastroenterology*, **103**(6), pp. 1372-1379.
- WINER, D.A., WINER, S., SHEN, L., WADIA, P.P., YANTHA, J., PALTSER, G., TSUI, H., WU, P., DAVIDSON, M.G., ALONSO, M.N., LEONG, H.X., GLASSFORD, A., CAIMOL, M., KENKEL, J.A., TEDDER, T.F., MCLAUGHLIN, T., MIKLOS, D.B., DOSCH, H.M. and ENGLEMAN, E.G., 2011. B cells promote insulin resistance through modulation of T cells and production of pathogenic IgG antibodies. *Nature medicine*, **17**(5), pp. 610-617.
- WOLF, A.M., WOLF, D., RUMPOLD, H., ENRICH, B. and TILG, H., 2004. Adiponectin induces the anti-inflammatory cytokines IL-10 and IL-1RA in human leukocytes. *Biochemical and biophysical research communications*, **323**(2), pp. 630-635.
- WOLFSDORF, J., SADEGHI-NEJAD, A. and SENIOR, B., 2002. Leptin-replacement therapy in lipodystrophy. *The New England journal of medicine*, **346**(25), pp. 2008-9; author reply 2009-10.
- WOLOSZYNEK, J.C., HU, Y. and PHAM, C.T., 2012. Cathepsin G-regulated release of formyl peptide receptor agonists modulate neutrophil effector functions. *The Journal of biological chemistry*, **287**(41), pp. 34101-34109.
- WON, J.H., KANG, N.N., AUH, C.K. and PARK, Y.M., 2003. The surface receptor is involved in annexin I-stimulated insulin secretion in MIN6N8a cells. *Biochemical and biophysical research communications*, **307**(2), pp. 389-394.
- WOOD, I.S., DE HEREDIA, F.P., WANG, B. and TRAYHURN, P., 2009. Cellular hypoxia and adipose tissue dysfunction in obesity. *The Proceedings of the Nutrition Society*, **68**(4), pp. 370-377.
- WOODS, S.C., LUTZ, T.A., GEARY, N. and LANGHANS, W., 2006. Pancreatic signals controlling food intake; insulin, glucagon and amylin. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, **361**(1471), pp. 1219-1235.
- WRIGHT, H.L., MOOTS, R.J., BUCKNALL, R.C. and EDWARDS, S.W., 2010. Neutrophil function in inflammation and inflammatory diseases. *Rheumatology (Oxford, England)*, **49**(9), pp. 1618-1631.

- WU, D., MOLOFSKY, A.B., LIANG, H.E., RICARDO-GONZALEZ, R.R., JOUIHAN, H.A., BANDO, J.K., CHAWLA, A. and LOCKSLEY, R.M., 2011. Eosinophils sustain adipose alternatively activated macrophages associated with glucose homeostasis. *Science (New York, N.Y.)*, **332**(6026), pp. 243-247.
- WU, J., SRINIVASAN, S.V., NEUMANN, J.C. and LINGREL, J.B., 2005. The KLF2 transcription factor does not affect the formation of preadipocytes but inhibits their differentiation into adipocytes. *Biochemistry*, **44**(33), pp. 11098-11105.
- XIAO, Y., JUNFENG, H., TIANHONG, L., LU, W., SHULIN, C., YU, Z., XIAOHUA, L., WEIXIA, J., SHENG, Z., YANYUN, G., GUO, L. and MIN, L., 2006. Cathepsin K in adipocyte differentiation and its potential role in the pathogenesis of obesity. *The Journal of clinical endocrinology and metabolism*, **91**(11), pp. 4520-4527.
- XU, B., JU, Y. and SONG, G., 2014. Role of p38, ERK1/2, focal adhesion kinase, RhoA/ROCK and cytoskeleton in the adipogenesis of human mesenchymal stem cells. *Journal of bioscience and bioengineering*, **117**(5), pp. 624-631.
- XU, F.P., CHEN, M.S., WANG, Y.Z., YI, Q., LIN, S.B., CHEN, A.F. and LUO, J.D., 2004. Leptin induces hypertrophy via endothelin-1-reactive oxygen species pathway in cultured neonatal rat cardiomyocytes. *Circulation*, **110**(10), pp. 1269-1275.
- XU, J. and LIAO, K., 2004. Protein kinase B/AKT 1 plays a pivotal role in insulin-like growth factor-1 receptor signaling induced 3T3-L1 adipocyte differentiation. *The Journal of biological chemistry*, **279**(34), pp. 35914-35922.
- XU, X., GRIJALVA, A., SKOWRONSKI, A., VAN EIJK, M., SERLIE, M.J. and FERRANTE, A.W., Jr, 2013. Obesity activates a program of lysosomal-dependent lipid metabolism in adipose tissue macrophages independently of classic activation. *Cell metabolism*, **18**(6), pp. 816-830.
- XUAN, W., QU, Q., ZHENG, B., XIONG, S. and FAN, G.H., 2015. The chemotaxis of M1 and M2 macrophages is regulated by different chemokines. *Journal of leukocyte biology*, **97**(1), pp. 61-69.
- YAHAGI, N., SHIMANO, H., HASTY, A.H., MATSUZAKA, T., IDE, T., YOSHIKAWA, T., AMEMIYA-KUDO, M., TOMITA, S., OKAZAKI, H., TAMURA, Y., IIZUKA, Y., OHASHI, K., OSUGA, J., HARADA, K., GOTODA, T., NAGAI, R., ISHIBASHI, S. and YAMADA, N., 2002. Absence of sterol regulatory element-binding protein-1 (SREBP-1) ameliorates fatty livers but not obesity or insulin resistance in Lep(ob)/Lep(ob) mice. *The Journal of biological chemistry*, **277**(22), pp. 19353-19357.
- YAMATO, E., IKEGAMI, H., TAKEKAWA, K., FUJISAWA, T., NAKAGAWA, Y., HAMADA, Y., UEDA, H. and OGIHARA, T., 1997. Tissue-specific and glucose-dependent expression of receptor genes for glucagon and glucagon-like peptide-1 (GLP-1). *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et metabolisme*, **29**(2), pp. 56-59.

YAMAUCHI, T., KAMON, J., WAKI, H., IMAI, Y., SHIMOZAWA, N., HIOKI, K., UCHIDA, S., ITO, Y., TAKAKUWA, K., MATSUI, J., TAKATA, M., ETO, K., TERAUCHI, Y., KOMEDA, K., TSUNODA, M., MURAKAMI, K., OHNISHI, Y., NAITOH, T., YAMAMURA, K., UHEYAMA, Y., FROGUEL, P., KIMURA, S., NAGAI, R. and KADOWAKI, T., 2003. Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis. *The Journal of biological chemistry*, **278**(4), pp. 2461-2468.

YAMAUCHI, T., KAMON, J., WAKI, H., TERAUCHI, Y., KUBOTA, N., HARA, K., MORI, Y., IDE, T., MURAKAMI, K., TSUBOYAMA-KASAOKA, N., EZAKI, O., AKANUMA, Y., GAVRILOVA, O., VINSON, C., REITMAN, M.L., KAGECHIKA, H., SHUDO, K., YODA, M., NAKANO, Y., TOBE, K., NAGAI, R., KIMURA, S., TOMITA, M., FROGUEL, P. and KADOWAKI, T., 2001. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipodystrophy and obesity. *Nature medicine*, **7**(8), pp. 941-946.

YAN, E., KO, E., LUONG, V., WANG, H.J., ROMANOVA, M. and LI, Z., 2008. Long-term changes in weight loss and obesity-related comorbidities after Roux-en-Y gastric bypass: a primary care experience. *American Journal of Surgery*, **195**(1), pp. 94-98.

YANG, S., LIU, A., WEIDENHAMMER, A., COOKSEY, R.C., MCCLAIN, D., KIM, M.K., AGUILERA, G., ABEL, E.D. and CHUNG, J.H., 2009. The role of mPer2 clock gene in glucocorticoid and feeding rhythms. *Endocrinology*, **150**(5), pp. 2153-2160.

YANG, T.T., XIONG, Q., ENSLEN, H., DAVIS, R.J. and CHOW, C.W., 2002. Phosphorylation of NFATc4 by p38 mitogen-activated protein kinases. *Molecular and cellular biology*, **22**(11), pp. 3892-3904.

YANG, W.S., LEE, W.J., FUNAHASHI, T., TANAKA, S., MATSUZAWA, Y., CHAO, C.L., CHEN, C.L., TAI, T.Y. and CHUANG, L.M., 2001. Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *The Journal of clinical endocrinology and metabolism*, **86**(8), pp. 3815-3819.

YANG, Y.H., MORAND, E.F., GETTING, S.J., PAUL-CLARK, M., LIU, D.L., YONA, S., HANNON, R., BUCKINGHAM, J.C., PERRETTI, M. and FLOWER, R.J., 2004. Modulation of inflammation and response to dexamethasone by Annexin 1 in antigen-induced arthritis. *Arthritis and Rheumatism*, **50**(3), pp. 976-984.

YANG, Y.H., SONG, W., DEANE, J.A., KAO, W., OOI, J.D., NGO, D., KITCHING, A.R., MORAND, E.F. and HICKEY, M.J., 2013. Deficiency of annexin A1 in CD4+ T cells exacerbates T cell-dependent inflammation. *Journal of immunology (Baltimore, Md.: 1950)*, **190**(3), pp. 997-1007.

YANG, Y.H., TOH, M.L., CLYNE, C.D., LEECH, M., AEBERLI, D., XUE, J., DACUMOS, A., SHARMA, L. and MORAND, E.F., 2006. Annexin 1 negatively regulates IL-6 expression via effects on p38 MAPK and MAPK phosphatase-1. *Journal of immunology (Baltimore, Md.: 1950)*, **177**(11), pp. 8148-8153.

- YAZBECK, R., HOWARTH, G.S. and ABBOTT, C.A., 2009. Dipeptidyl peptidase inhibitors, an emerging drug class for inflammatory disease? *Trends in pharmacological sciences*, **30**(11), pp. 600-607.
- YE, J., 2009. Emerging role of adipose tissue hypoxia in obesity and insulin resistance. *International journal of obesity (2005)*, **33**(1), pp. 54-66.
- YE, R.D., BOULAY, F., WANG, J.M., DAHLGREN, C., GERARD, C., PARMENTIER, M., SERHAN, C.N. and MURPHY, P.M., 2009. International Union of Basic and Clinical Pharmacology. LXXIII. Nomenclature for the formyl peptide receptor (FPR) family. *Pharmacological reviews*, **61**(2), pp. 119-161.
- YEH, W.C., CAO, Z., CLASSON, M. and MCKNIGHT, S.L., 1995. Cascade regulation of terminal adipocyte differentiation by three members of the C/EBP family of leucine zipper proteins. *Genes & development*, **9**(2), pp. 168-181.
- YIN, J., GAO, Z., HE, Q., ZHOU, D., GUO, Z. and YE, J., 2009. Role of hypoxia in obesity-induced disorders of glucose and lipid metabolism in adipose tissue. *American journal of physiology. Endocrinology and metabolism*, **296**(2), pp. E333-42.
- YONA, S., BUCKINGHAM, J.C., PERRETTI, M. and FLOWER, R.J., 2004. Stimulus-specific defect in the phagocytic pathways of annexin 1 null macrophages. *British journal of pharmacology*, **142**(5), pp. 890-898.
- YOUSSEIF, A., EMMANUEL, J., KARRA, E., MILLET, Q., ELKALAAWY, M., JENKINSON, A.D., HASHEMI, M., ADAMO, M., FINER, N., FIENNES, A.G., WITHERS, D.J. and BATTERHAM, R.L., 2014. Differential effects of laparoscopic sleeve gastrectomy and laparoscopic gastric bypass on appetite, circulating acyl-ghrelin, peptide YY3-36 and active GLP-1 levels in non-diabetic humans. *Obesity Surgery*, **24**(2), pp. 241-252.
- YU, S., RAO, S. and REDDY, J.K., 2003. Peroxisome proliferator-activated receptors, fatty acid oxidation, steatohepatitis and hepatocarcinogenesis. *Current Molecular Medicine*, **3**(6), pp. 561-572.
- YUDKIN, J.S., STEHOUWER, C.D., EMEIS, J.J. and COPPACK, S.W., 1999. C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arteriosclerosis, Thrombosis, and Vascular Biology*, **19**(4), pp. 972-978.
- ZABOLOTNY, J.M., BENCE-HANULEC, K.K., STRICKER-KRONGRAD, A., HAJ, F., WANG, Y., MINOKOSHI, Y., KIM, Y.B., ELMQUIST, J.K., TARTAGLIA, L.A., KAHN, B.B. and NEEL, B.G., 2002. PTP1B regulates leptin signal transduction in vivo. *Developmental cell*, **2**(4), pp. 489-495.
- ZARKESH-ESFAHANI, H., POCKLEY, G., METCALFE, R.A., BIDLINGMAIER, M., WU, Z., AJAMI, A., WEETMAN, A.P., STRASBURGER, C.J. and ROSS, R.J., 2001.

High-dose leptin activates human leukocytes via receptor expression on monocytes. *Journal of immunology (Baltimore, Md.: 1950)*, **167**(8), pp. 4593-4599.

ZECHNER, R., ZIMMERMANN, R., EICHMANN, T.O., KOHLWEIN, S.D., HAEMMERLE, G., LASS, A. and MADEO, F., 2012. FAT SIGNALS--lipases and lipolysis in lipid metabolism and signaling. *Cell metabolism*, **15**(3), pp. 279-291.

ZENG, J., DENG, S., WANG, Y., LI, P., TANG, L. and PANG, Y., 2017. Specific Inhibition of Acyl-CoA Oxidase-1 by an Acetylenic Acid Improves Hepatic Lipid and Reactive Oxygen Species (ROS) Metabolism in Rats Fed a High Fat Diet. *The Journal of biological chemistry*, **292**(9), pp. 3800-3809.

ZHANG, H., XIE, H., ZHAO, Q., XIE, G.Q., WU, X.P., LIAO, E.Y. and LUO, X.H., 2010. Relationships between serum adiponectin, apelin, leptin, resistin, visfatin levels and bone mineral density, and bone biochemical markers in post-menopausal Chinese women. *Journal of endocrinological investigation*, **33**(10), pp. 707-711.

ZHANG, Z., HUANG, L., ZHAO, W. and RIGAS, B., 2010. Annexin 1 induced by anti-inflammatory drugs binds to NF-kappaB and inhibits its activation: anticancer effects in vitro and in vivo. *Cancer research*, **70**(6), pp. 2379-2388.

ZHONG, J., RAO, X., DEIULIIS, J., BRAUNSTEIN, Z., NARULA, V., HAZEY, J., MIKAMI, D., NEEDLEMAN, B., SATOSKAR, A.R. and RAJAGOPALAN, S., 2013. A potential role for dendritic cell/macrophage-expressing DPP4 in obesity-induced visceral inflammation. *Diabetes*, **62**(1), pp. 149-157.

ZHOU, Q., GE, Q., DING, Y., QU, H., WEI, H., WU, R., YAO, L., WEI, Q., FENG, Z., LONG, J. and DENG, H., 2018. Relationship between serum adipon and the first phase of glucose-stimulated insulin secretion in individuals with different glucose tolerance. *Journal of diabetes investigation*, **9**(5), pp. 1128-1134.

ZHU, L., HAN, M.B., GAO, Y., WANG, H., DAI, L., WEN, Y. and NA, L.X., 2015. Curcumin triggers apoptosis via upregulation of Bax/Bcl-2 ratio and caspase activation in SW872 human adipocytes. *Molecular medicine reports*, **12**(1), pp. 1151-1156.

ZHUGE, F., NI, Y., NAGASHIMADA, M., NAGATA, N., XU, L., MUKAIDA, N., KANEKO, S. and OTA, T., 2016. DPP-4 Inhibition by Linagliptin Attenuates Obesity-Related Inflammation and Insulin Resistance by Regulating M1/M2 Macrophage Polarization. *Diabetes*, **65**(10), pp. 2966-2979.

ZIMMERMANN, R., STRAUSS, J.G., HAEMMERLE, G., SCHOISWOHL, G., BIRNER-GRUENBERGER, R., RIEDERER, M., LASS, A., NEUBERGER, G., EISENHABER, F., HERMETTER, A. and ZECHNER, R., 2004. Fat mobilization in adipose tissue is promoted by adipose triglyceride lipase. *Science (New York, N.Y.)*, **306**(5700), pp. 1383-1386.

Appendix

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

Content removed on data protection grounds

IRAS project ID	220666
-----------------	--------

Appendix A - List of Documents

Content removed on data protection grounds

IRAS project ID	220666
-----------------	--------

Appendix B - Summary of HRA Assessment

Content removed on data protection grounds

Content removed on data protection grounds

IRAS project ID	220666
-----------------	--------

Content removed on data protection grounds

Plasma Annexin A1 levels and insulin sensitivity during weight loss through Bariatric Surgery

Abstract

Recently, the adipose tissue has been acknowledged to be more than just a site of lipid storage as it contributes to metabolic homeostasis through the secretion of adipose tissue specific proteins; the “adipokines”. Obesity refers to excess lipid storage in the adipose tissue often leading to ectopic deposition in other organs affecting their physiological function. Bariatric surgery; otherwise known as weight loss surgery is thought to be an effective option for individuals who are morbidly obese or have a BMI of over 40. However, individuals often exhibit alterations in the levels of adipokines and insulin sensitivity leading to insulin resistance, despite weight loss. Therefore, the aim of this project is to investigate the levels of an anti-inflammatory protein AnxA1 during weight loss through bariatric surgery. AnxA1 is thought to regulate the function of insulin and preliminary data suggest it decreases the levels of pro-inflammatory markers which may contribute to effective resolution of inflammation and development of diabetes type 2 in these patients.

Introduction

Obesity is a growing pan-endemic health problem in both developing and developed countries. It is a complex disease which results from an increase in fat mass or increased adiposity and is defined as body mass index (BMI) of over 30kg/m² or a waist circumference more than 94cm for men and 80cm for women (Goktas, Moustaid-Moussa et al. 2013). It is a chronic disease as it evolves over the years and low grade inflammation causes metabolic changes. Therefore, it is a major risk factor of developing common diseases such as type 2 diabetes, hypertension and cardiovascular disease, thereby significantly increasing total mortality (Lindegard, Jorgensen et al. 2015). For example; insulin resistance in obesity is thought to be associated to adipose tissue fibrosis, hypoxia and infiltration of immune cells and inflammation leading to cellular hypertrophy, triggering cellular stress processes impairing adipocyte responses to insulin (Moreno-Castellanos, Guzman-Ruiz et al. 2016). Despite recognising the relationship of obesity and the onset of these diseases, the biological link to the pathogenesis is not well understood.

The first recommended treatment for obesity is weight loss through low calorie diets and regular physical exercise. However, lifestyle interventions have low observance and effectiveness in morbidly obese individuals with a BMI of over 40kg/m² (Goktas, Moustaid-Moussa et al. 2013). Bariatric surgery is considered to be the best choice for these individuals as it leads to weight loss through restriction and/or malabsorption of nutrients (Adami, Scopinaro et al. 2016). It is an effective and durable treatment in comparison to non-surgical interventions such as life style changes mentioned above and pharmacological therapies. After surgery individuals lose approximately 30% of their body weight and decrease the overall inflammatory response (Goktas, Moustaid-Moussa et al. 2013). In addition to weight loss, bariatric surgery reduces insulin resistance, cardiovascular disease and mortality. However, some patients experience remission of insulin resistance and the onset of diabetes (Adami, Scopinaro et al. 2016). Bariatric surgery causes changes in the adipose tissue biology that impact systemic glucose utilisation thereby affecting insulin sensitivity independent of weight loss (Moreno-Castellanos, Guzman-Ruiz et al. 2016). Metabolic changes observed after bariatric surgery independent of weight loss are thought to result from altered hormonal patterns; i.e upregulation of gastrointestinal insulinotrophic polypeptide (GIP), glucagon-like peptide 1 (GLP-1) and PYY that cause inhibition of appetite, mediating satiety and stimulate insulin production (Goktas, Moustaid-Moussa et al. 2013).

Adipose tissue is a site for storing triglycerides and fats in periods of excess energy intake and releasing them during periods of famine to provide energy. Additionally, it provides mechanical protection for organs and thermal insulation for the organism. It was considered to be a mere site for fat storage but now it is acknowledged as a “metabolically active” endocrine organ. It secretes hormones, cytokines and chemokines, collectively known as ‘adipokines’ that participate in appetite and satiety, insulin secretion and sensitivity, regulating energy homeostasis, inflammation, endothelial function other biological functions (Yang, Lee et al. 2001). Adipokines are detectable both in the plasma and adipose tissue and are active factors that modulate the effect of obesity and related co-morbidities. Dysregulated production of these adipokines in obesity is thought to contribute to the development of chronic low-grade inflammation leading to the onset of associated diseases (Lindegaard, Jorgensen et al. 2015).

Leptin (adipose tissue specific) is a satiety hormone that regulates energy balance by inhibiting hunger, insulin sensitivity and β -cell apoptosis (Goktas, Moustaid-Moussa et al. 2013). The plasma levels of leptin are directly proportional to body fat (Oswal, Yeo 2010). Adiponectin (adipose tissue specific) is involved in regulating glucose levels and fatty acids metabolism in insulin sensitive tissues. It is an anti-inflammatory, anti-apoptotic protein which stimulates energy expenditure and insulin production and sensitivity (Sams, Blackledge et al. 2016). The plasma levels of adiponectin inversely correlate with body fat, insulin resistance and dyslipidaemia. Adiponectin is activated during adipogenesis; however, a feedback inhibition on its production may impose the development of obesity (Yang, Lee et al. 2001). Weight loss through bariatric surgery has shown to have an inverse relationship of plasma adiponectin and weight loss (Sams, Blackledge et al. 2016). As individuals lose weight, there is an increase in plasma adiponectin levels and contributing to the resolution of co-morbidities. In addition, adipose tissue secretes universal cytokines including TNF- α , IL-1 β and IL-6. TNF- α is involved in systemic inflammation and is one of the cytokines that make up the acute phase reaction. The plasma levels of TNF- α directly correlate with body fat and is produced by dysfunctional adipocytes which is thought to impair insulin signalling and secretion (Adami, Scopinaro et al. 2016). IL-1 β is a potent pro-inflammatory cytokine that is upregulated in obesity and functionally it induces prostaglandin synthesis, neutrophil influx and T cell activation (Goktas, Moustaid-Moussa et al. 2013). Furthermore, IL-1 β secreted from dysfunctional adipocytes is thought to play a role in apoptosis, β cell development and destruction, increasing glycaemic deterioration leading to the onset of diabetes type 2.

The protein of interest in this project is AnxA1; it is a calcium dependent phospholipid binding protein which is produced and secreted by a variety of cell types including adipocytes, neutrophils and macrophages. It exhibits anti-inflammatory properties as it mediates the detachment of neutrophils from the vascular cells through mediating L-selectin shedding, inhibits the activity of phospholipase A2 (inhibiting production of eicosanoids) and inhibits activation of pro-inflammatory CD4⁺ T cell lineages (Rescher, Gerke 2004)(Kosicka, Cunliffe et al. 2013). Plasma AnxA1 levels are decreased in obese in comparison to normal weight individuals showing an inverse relationship (Kosicka, Cunliffe et al. 2013). Interestingly, knockout models have shown opposing results with some reporting increased susceptibility

to weight gain and insulin resistance and others increased weight loss (Akasheh, Pini et al. 2013, Warne, John et al. 2006).

Aims and Hypotheses:

The primary aim of this pilot project is to investigate the changes in AnxA1 and other plasma inflammatory biomarkers, gut hormones and insulin sensitivity in relation to weight loss in individuals undergoing bariatric surgery.

Studies:

- 1) Changes in body weight of patients over 3 months
- 2) The effect of weight loss on Plasma AnxA1 levels by types of bariatric procedures
- 3) The plasma levels of inflammatory biomarkers and their correlation with insulin sensitivity
- 4) The plasma levels of Gut hormones and their correlation with AnxA1 and inflammatory biomarkers

Hypotheses:

- 1) Weight loss results in increased plasma AnxA1 levels leading to improved insulin sensitivity leading to effective resolution of inflammation and decreasing the likely hood of developing co-morbidities.
- 2) Plasma AnxA1 levels directly correlate with anti-inflammatory and inversely correlate with pro-inflammatory markers during weight loss.
- 3) AnxA1 positively correlates insulin sensitivity and gut hormones


Experimental design and methods

Study population

Participants will be recruited from the bariatric surgical clinic at Heartlands Hospital, Birmingham. Written informed consent will be taken from up to 30 adult Obese participants who meet the criteria for bariatric surgery (Sleeve gastrectomy and Roux-en-Y gastric bypass); Adult subjects Male/Female, aged between 18-75 years, BMI of >40 if do not have type 2 diabetes and >35kg/m² if known to have type 2 diabetes. Type 1 Diabetic, have significant

mental health problems or patients receiving long term steroids will be excluded from the study (figure 1).

Figure 1: Flow diagram of the exclusion criteria and allocation into subpopulations (Non-Diabetic and Diabetic).



Recruitment

Patients attending the weight management clinics and identified for bariatric surgery will be approached for participation in the study. They will be provided with the patient information sheet, the purpose of the study will be explained and any questions and apprehensions will be discussed. Interested patients will be contacted 48 hours post the initial meeting and if they are willing to take part in the study, they will be invited for a screening visit. After obtaining the informed consent, patients will be screened for eligibility and baseline visit will be agreed.

Study design

Eligible participants will have clinical, biochemical, inflammatory markers and gut hormones measured at baseline and at each of the follow up appointments. A subset of participants who provide informed consent will undergo Oral Glucose Tolerance Test (OGTT) to assess insulin sensitivity. Similarly, the test will be carried out at baseline and at each of the follow up appointments.

Clinical and Biochemical Measures

The following clinical information will be taken from all participants: age, gender, past medical history, treatment, duration of diabetes, weight, height, BMI and systolic and diastolic blood pressure pre-and postoperatively. 5mL of whole blood sample will be collected via venepuncture at the same time and sent to the local routine lab within the hospital for the analysis of biochemical measurements. These include fasting glucose and lipids (total cholesterol, low density lipoprotein-cholesterol (LDL-C), high density lipoprotein-cholesterol (HDL-C) and triglycerides). Insulin and C-peptide will be measured using ELISA as described below. The clinical assessment and collection of blood samples will be done by a trained professional.

Inflammatory markers and Gut hormones

In addition, 10mL of whole blood will be collected in purple top anticoagulant-treated tubes (EDTA-treated) via venepuncture by a trained professional. The samples will be centrifuged for 10 minutes at 1,000-2,000xg to separate into serum and plasma. The supernatant (plasma) will be collected, aliquoted (0.5mL) and stored at -80C.

The plasma will be used to measure range of inflammatory markers including; CRP, AnxA1, IL-1 β , TNF- α , Leptin and adiponectin and Gut hormones; PYY and GLP-1. The biomarkers will be measured using commercially available sandwich ELISA kits following the manufacturer's methodology at Coventry University.

Typical Sandwich ELISA methodology:

The capture antibody is coated onto a solid support; 96 well plate. The antigen is incubated with a primary antibody, followed by a wash with an appropriate buffer to remove any excess, unbound antigens. The primary antibody is incubated with an enzyme labelled secondary

antibody. The most widely used enzymes for detection are Horse radish peroxidase (HRP) and alkaline phosphatase (ALP). The enzymes are incubated with their appropriate substrates; P-Nitrophenyl-phosphate for ALP and hydrogen peroxide for HRP, and the chemical reaction result in change of colour. The amount of colour produced is quantified using a spectrophotometer plate reader at a target wavelength and is proportional to the amount of antibody/antigen present.

Oral Glucose Tolerance Test:

The OGTT is specific to this study, therefore, only a subset of participants will undergo this test. Participants who consent participating in an OGTT will be made aware that they need to abstain from ingesting food and fast from the previous night (fast >10 hours) will be required. 2.7mL of fasting blood sample will be collected by venepuncture in a grey top tube (containing sodium fluoride; an antiglycolytic agent) and stored on ice. It will be recorded as 0 minutes. Participants will be given 75g anhydrous glucose in the form of a drink which they will need to consume within 5 minutes. Second blood sample (2.7mL) will be collected by venepuncture at +120 minutes and stored on ice. If any glucose is lost by vomiting, the test will be discontinued. The samples will be sent the routine hospital lab for analysis of blood glucose, insulin and C-peptide levels. The OGTT will be carried out by a trained professional.

Measurement of insulin sensitivity

The homeostasis model assessment (HOMA) will be used to estimate steady state β -cell function (% β) and insulin sensitivity (%S) preoperatively and postoperatively.

The following equations are going to be used to calculate %S and % β :

$$\text{HOMA}\% \beta = \frac{20 \times \text{Insulin}}{\text{Glucose} - 3.5} \quad (\text{Glucose in Molar Units mmol/L})$$

$$\text{HOMA}\% S = \frac{\text{Glucose} \times \text{Insulin}}{22.25} \quad (\text{Glucose in Molar Units mmol/L})$$

Data collection

All data will be collected on a standard clinical records folder and then entered on to a secure database. All subjects will be allocated a study number and no identifiable information will be

transferred on to the data base. Access to the data will be restricted to the study team and will be password protected.

Statistical analysis

The statistical analysis will be carried out using one way Anova, Tukeys post hoc test and non-parametric spearman rank correlation. Inflammatory Biomarkers and Gut hormones will be analysed by ANOVA. The analysis will be performed using Graphpad and the level of significance will be set at $P < 0.05$.

Confidentiality

All the patients' data will be coded so that no personal identifiers are on the same database as the results from biochemical, gut hormones or inflammatory biomarker analysis. All databases will be password protected and kept on passworded computers in locked rooms. Hard copy of the original data will be kept in sealed enveloped in locked filing cabinets. Only the people directly involved in the project will have access to any of the results from the project.

Benefits of the project

The aim of this project is to investigate the relationship between the levels of AnxA1 and weight loss and the results of this pilot project will contribute to our understanding of the link between anti-inflammatory markers and obesity. Therefore, AnxA1 could be used as a biomarker and could aid in the development of effective therapeutic strategies to help overcome obesity associated inflammation.

The outcome of the pilot project can lead to investigating the area further using a bigger cohort and investigating a range of different markers. Furthermore, different populations of individuals can be recruited including those who have undergone surgery, have had effective weight loss, but have relapsed diabetes type 2.

Resources and costs

This project is funded by the Knowledge Exchange Unit at Coventry University through their SPIDER project. The main resources required for this project will be ELISA Kits to measure different biomarkers. Each kit is approximately £500 (96 tests).

References

ADAMI, G.F., SCOPINARO, N. and CORDERA, R., 2016. Adipokine Pattern After Bariatric Surgery: Beyond the Weight Loss. *Obesity Surgery*, **26**(11), pp. 2793-2801.

AKASHEH, R.T., PINI, M., PANG, J. and FANTUZZI, G., 2013. Increased adiposity in annexin A1-deficient mice. *PLoS one*, **8**(12), pp. e82608.

GOKTAS, Z., MOUSTAID-MOUSSA, N., SHEN, C.L., BOYLAN, M., MO, H. and WANG, S., 2013. Effects of bariatric surgery on adipokine-induced inflammation and insulin resistance. *Frontiers in endocrinology*, **4**, pp. 69.

KOSICKA, A., CUNLIFFE, A.D., MACKENZIE, R., ZARIWALA, M.G., PERRETTI, M., FLOWER, R.J. and RENSHAW, D., 2013. Attenuation of plasma annexin A1 in human obesity. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **27**(1), pp. 368-378.

LINDEGAARD, K.K., JORGENSEN, N.B., JUST, R., HEEGAARD, P.M. and MADSBAD, S., 2015. Effects of Roux-en-Y gastric bypass on fasting and postprandial inflammation-related parameters in obese subjects with normal glucose tolerance and in obese subjects with type 2 diabetes. *Diabetology & metabolic syndrome*, **7**, pp. 12-015-0012-9. eCollection 2015.

MORENO-CASTELLANOS, N., GUZMAN-RUIZ, R., CANO, D.A., MADRAZO-ATUTXA, A., PEINADO, J.R., PEREIRA-CUNILL, J.L., GARCIA-LUNA, P.P., MORALES-CONDE, S., SOCAS-MACIAS, M., VAZQUEZ-MARTINEZ, R., LEAL-CERRO, A. and MALAGON, M.M., 2016. The Effects of Bariatric Surgery-Induced Weight Loss on Adipose Tissue in Morbidly Obese Women Depends on the Initial Metabolic Status. *Obesity Surgery*, **26**(8), pp. 1757-1767.

OSWAL, A. and YEO, G., 2010. Leptin and the control of body weight: a review of its diverse central targets, signaling mechanisms, and role in the pathogenesis of obesity. *Obesity (Silver Spring, Md.)*, **18**(2), pp. 221-229.

RESCHER, U. and GERKE, V., 2004. Annexins--unique membrane binding proteins with diverse functions. *Journal of cell science*, **117**(Pt 13), pp. 2631-2639.

SAMS, V.G., BLACKLEDGE, C., WIJAYATUNGA, N., BARLOW, P., MANCINI, M., MANCINI, G. and MOUSTAID-MOUSSA, N., 2016. Effect of bariatric surgery on systemic and adipose tissue inflammation. *Surgical endoscopy*, **30**(8), pp. 3499-3504.

WARNE, J.P., JOHN, C.D., CHRISTIAN, H.C., MORRIS, J.F., FLOWER, R.J., SUGDEN, D., SOLITO, E., GILLIES, G.E. and BUCKINGHAM, J.C., 2006. Gene deletion reveals roles for annexin A1 in the regulation of lipolysis and IL-6 release in epididymal adipose tissue. *American journal of physiology. Endocrinology and metabolism*, **291**(6), pp. E1264-73.

YANG, W.S., LEE, W.J., FUNAHASHI, T., TANAKA, S., MATSUZAWA, Y., CHAO, C.L., CHEN, C.L., TAI, T.Y. and CHUANG, L.M., 2001. Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *The Journal of clinical endocrinology and metabolism*, **86**(8), pp. 3815-3819.



Coventry University
 Priory St,
 Coventry,
 CV1 5FB



Content removed on data protection grounds

Patient Information Sheet

Plasma Annexin A1 levels and insulin sensitivity during weight loss through Bariatric Surgery

CU Ethics Reference Number: [P50481](#)

IRAS Ethics Reference Number: [220666](#)

Sehar Sajid, Professor Derek Renshaw, Dr Srikanth Bellary, Helen Jenner

You are invited to take part in a study regarding obesity. Thank you for considering helping with our research. This form explains what you will be asked to do. If you have any questions

about this, please ask me. The project is being carried out as part of my PhD under the supervision of Dr S Bellary and Professor D Renshaw.

By signing this form, you agree to take part in the study. However, please note that you are free to stop taking part at any time. Blood samples and data collected with your consent prior to withdrawal will be used in this study. You will be given a copy of the consent form to keep for your records.

Information about the project/Purpose of the Project

My PhD project involves investigating the role of an anti-inflammatory protein; Annexin A1 in obesity associated inflammation. Obesity alters the level of inflammation and biomarkers in human blood. To further my project and our understanding of the relationship between inflammatory biomarkers and obesity, I am proposing a study in which I will be investigating the levels of Annexin A1 during weight loss through bariatric surgery and whether it increases following weight loss. Additionally, I will be correlating it with inflammatory biomarkers, gut hormones and insulin sensitivity.

Why have I been chosen?

We are looking for volunteers who will be undergoing bariatric surgery and I understand you are scheduled for the surgery soon therefore, your involvement and participation in this project will be highly valuable.

Do I have to take part?

If you don't want to take part, you don't have to give a reason, and it won't affect the care you receive. If you do want to take part now, but change your mind later, you can pull out of the study at any time. If you agree to take part in this study, you will be asked to sign the Consent Form on the last page of this document. This document is 5 pages long, including the Consent Form. Please make sure you have read and understood all the pages. You can discuss it with your family, friends or GP if you wish. You will be given a copy of both the Participant Information Sheet and the Consent Form to keep.

What do I have to do?

You will be asked to visit the clinic at Heartlands Hospital on 3 separate occasions; 1) before your surgery which will be around the time of your pre-operative assessment 2) 6 weeks after your surgery and 3) 3 months after your surgery. To minimise the inconvenience for you, the visits will be arranged in advance with you to coincide with your clinical visits.

During each of the visits which is expected to last between 1-1.5 hours, you will undergo basic clinical assessment in which your height, weight and blood pressure will be measured. A venous blood sample will be taken for biochemical analysis such as glucose, insulin and lipids by a trained member of staff. A separate venous blood sample will be collected for analysis of inflammatory markers and gut hormones to meet the aims of this project.

If you agree to take part and give your consent, you will be asked if you would like to participate in an additional test; the oral glucose tolerance test (OGGT). If you agree to take part, you will need to undergo this test on the 3 separate visits mentioned above. This test takes 2 hours to complete, so your visit is expected to last up to 3.5 hours. It involves coming into the clinic after an overnight fast, where you will have blood taken followed by a drink containing 75g glucose. The blood test will be repeated after 2 hours. The blood samples will be measured for glucose, insulin and c-peptide measurement. You may experience some side effects which include nausea, stomach discomfort, diarrhoea and/or constipation as a result but you will be monitored throughout. This test is designed to test your body's ability to use sugar and can be used to diagnose prediabetes, diabetes and the other markers of insulin resistance. The participation in this test is optional and will not affect your decision to take part in the study.

Some individuals can feel dizzy or faint when they have needles inserted into veins. If you have previously fainted whilst giving blood, then please speak to me before you take part in the study. All participants will complete a standard laboratory health screening and risk assessment form prior to the sessions.

What are the benefits of taking part?

There are no extrinsic benefits for taking part in this study, however the information gathered will be important for future scientific studies which investigate role of Annexin A1 in adiposity.

Data protection and confidentiality

Your consent to participate in this study will be confidential. I assure you that complete confidentiality of data gathered in this investigation will be maintained. Your identity will not be made public without your prior consent. The samples will be transported to Coventry University in a secure boot of a car in an icebox and stored in locked -80°C Freezers. To ensure anonymity and confidentiality, the data will be coded and the original data will be stored in sealed envelopes in locked cupboards and passworded computers. Only myself and my supervisors will have access to the data collected and it will be destroyed within 5 years. At the expiry date, the samples will be incinerated and the sealed envelopes will be destroyed securely by shredding, by the Faculty Research Support Officer. The results of the project may be published in scientific journals and will be used in my Thesis which will be available through the Coventry University Library. A copy of the results can be made available upon request by contacting me once the project has been completed.

All information which is collected during the course of the research will be kept strictly confidential and comply with the principles of the Data Protection Act (1998). Please note that only the experimental team will have access to any data with your name on it. Any data stored electronically will use subject codes to retain anonymity. All data subsequently presented by these staff will be anonymous such that only these two staff will be able to determine that the data was gained from you. This study has been approved by the Coventry University – Faculty of Health and Life Science Ethics Committee and NHS ethics through IRAS. If you have any queries about this research, please ask;

Who should you talk to if you have questions

Content removed on data protection grounds

Thank you,

Sehar Sajid

**Coventry University**Priory St,
Coventry,

CV1 5FB

Patient Identification Number:

CONSENT FORM

Title of Project: Plasma Annexin A1 levels and insulin sensitivity during weight loss through Bariatric Surgery

CU Ethics Reference Number: P50481

IRAS Ethics Reference Number: 220666

Name of Researchers: Dr S. Bellary

Professor D Renshaw

Sehar Sajid

Helen Jenner

Please initial all boxes

1. I confirm that I have read and understand the information sheet dated **01/08/2017** (version **2**) for the above study. I have had the opportunity to consider the information, ask questions and have had these answered satisfactorily.

☐

2. I understand that my participation is voluntary and that I am free to withdraw at any time without giving any reason, without my medical care or legal rights being affected.

☐

3. I understand that relevant sections of my medical notes and data collected during the study, may be looked at by individuals from Coventry University, from regulatory authorities or from the NHS Trust, where it is relevant to my taking part in this research. I give permission for these individuals to have access to my records.

☐

4. I allow the researcher to inform my GP about my participation in the study

☐

5. I agree to take part in the Oral Glucose Tolerance Test

☐

6. I agree to withdrawal of blood samples

☐

7. I agree to take part in the above study.



_____	_____	_____
Name of Participant	Date	Signature

_____	_____	_____
Name of Person	Date	Signature

taking consent.